


Collectivity in Ultraperipheral Pb + Pb Collisions at the Large Hadron Collider

Wenbin Zhao¹, Chun Shen^{1,2}, and Björn Schenke³

¹*Department of Physics and Astronomy, Wayne State University, Detroit, Michigan 48201, USA*

²*RIKEN BNL Research Center, Brookhaven National Laboratory, Upton, New York 11973, USA*

³*Physics Department, Brookhaven National Laboratory, Upton, New York 11973, USA*

 (Received 24 March 2022; revised 27 September 2022; accepted 8 November 2022; published 13 December 2022)

We present the first full $(3 + 1)$ D dynamical simulations of ultraperipheral Pb + Pb collisions at the Large Hadron Collider. Extrapolating from $p + \text{Pb}$ collisions, we explore whether a quasireal photon γ^* interacting with the lead nucleus in an ultraperipheral collision can create a many-body system exhibiting fluid behavior. Assuming strong final-state interactions, we provide model results for charged hadron multiplicity, identified particle mean transverse momenta, and charged hadron anisotropic flow coefficients, and compare them with experimental data from the ALICE and ATLAS Collaborations. The elliptic flow hierarchy between $p + \text{Pb}$ and $\gamma^* + \text{Pb}$ collisions is dominated by the difference in longitudinal flow decorrelations and reproduces the experimental data well. We have demonstrated that our theoretical framework provides a quantitative tool to study particle production and collectivity for all system sizes, ranging from central heavy-ion collisions to small asymmetric collision systems at the Relativistic Heavy-Ion Collider and the Large Hadron Collider and even at the future Electron-Ion Collider.

DOI: [10.1103/PhysRevLett.129.252302](https://doi.org/10.1103/PhysRevLett.129.252302)

Introduction.—Ultrarelativistic collisions of heavy ions create and allow for the study of a novel state of matter, the quark-gluon plasma (QGP), which exhibits the degrees of freedom of the fundamental building blocks of visible matter. Precise measurements of the emergent collectivity of the produced matter were performed at the Relativistic Heavy Ion Collider (RHIC) and the Large Hadron Collider (LHC). Based on the success of hydrodynamic descriptions, extensive phenomenological studies have revealed that the created QGP fireball behaves like a nearly perfect fluid with very small specific shear viscosity (see Refs. [1–3]).

Rather strikingly, many features of collective expansion have also been observed in very small collision systems, such as $p + \text{Au}$, $d + \text{Au}$, $^3\text{He} + \text{Au}$ at RHIC [4,5], and $p + p$ and $p + \text{Pb}$ collisions at the LHC [6–8]. The flowlike signals in the small systems can be interpreted as a result of the hydrodynamic response of the QGP medium to the initial collision geometry [9–16]. Alternatively, the color glass condensate effective theory has also predicted a significant amount of correlations in the initial state of these small collision systems, that can mimic collective behavior to a certain degree [17–21]. Theoretical frameworks including both initial-state correlations and final-state interactions have been used to better understand the origin of the observed collective behavior in these small systems [22–24].

Recently, intriguing experimental results on two-particle azimuthal correlations in ultraperipheral Pb + Pb collisions (UPCs) at the LHC [25] have appeared. UPCs have appreciable rates of photonuclear interactions [26,27],

and the ATLAS measurements of such photonuclear ($\gamma^* + \text{Pb}$) interactions in Pb + Pb UPCs indicate the persistence of collective phenomena with the strength of correlations comparable to that observed in proton-proton and proton-lead collisions in similar multiplicity ranges [25].

Quantitative understanding of the many-body dynamics in UPC events poses big challenges to the theory community. First, it is known that in asymmetric systems boost invariance is strongly broken [28–32]. Because of the largely different incoming energies between the quasireal photon γ^* and the Pb nucleus, the $\gamma^* + \text{Pb}$ collision is highly asymmetric and the violation of boost invariance is expected to be even greater than that in $p + \text{Pb}$ collisions. The rapidity decorrelation of the collision geometry thus plays an essential role when computing and measuring the magnitudes of anisotropic flow coefficients in $\gamma^* + \text{Pb}$ and $p + \text{Pb}$ collisions. Second, the small sizes of the collision systems are pushing the limit of the applicability of the causal relativistic viscous hydrodynamic description [33–36]. Resolving these challenges would bring new exciting opportunities to study collectivity in small systems at the upcoming Electron-Ion Collider, where one has experimental control on the virtuality of the colliding photon and can use it as a dial to change the collision system size.

In the present Letter, we explore the collective QGP signatures in $\gamma^* + \text{Pb}$ collisions at the LHC by employing a full $(3 + 1)$ D dynamical framework with hydrodynamics and hadronic transport [37]. This framework was shown to provide a unified and quantitative description of particle

production in proton-proton, proton-nucleus, and nucleus-nucleus collisions across center-of-mass energies of a few GeV to several TeV [37]. We calibrate this model framework with the $p + \text{Pb}$ measurements at a center-of-mass energy $\sqrt{s} = 5.02$ TeV and then make predictions for $\gamma^* + \text{Pb}$ collisions in the UPC Pb + Pb events.

Our study provides the first quantitative predictions of the anisotropic flow hierarchy between $\gamma^* + \text{Pb}$ and $p + \text{Pb}$ collisions from a final-state-dominated theoretical framework. We also present results for the full rapidity dependence of particle yields and predictions for the photon virtuality dependence of midrapidity elliptic flow.

Methodology.—In photon-nucleus, $\gamma^* + \text{Pb}$, collisions, the virtual photon state may be decomposed into a set of vector meson (VM) states, like ρ , ω , and ϕ in the vector meson dominance picture [38]. Here, we treat the virtual photon as a vector meson with a lifetime longer than the time of interactions in the low virtuality regime, $Q^2 \sim \Lambda_{\text{QCD}}^2 - 1 \text{ GeV}^2$ [39]. Under these assumptions, the photon-nucleus interaction in UPCs proceeds as a vector meson-nucleus collision at an energy lower than that of the associated nucleus-nucleus collisions. Following Ref. [27], we consider fluctuating kinematics of the photon-nucleus systems in ultraperipheral A + A collisions. The probability distribution of the center-of-mass collision energy is

$$P(\sqrt{s_{\gamma^*N}}) \propto \frac{1}{\sqrt{s_{\gamma^*N}}} \left\{ w_R^{AA} K_0(w_R^{AA}) K_1(w_R^{AA}) - \frac{(w_R^{AA})^2}{2} [K_1^2(w_R^{AA}) - K_0^2(w_R^{AA})] \right\}, \quad (1)$$

where $w_R^{AA} = 2kR_A/\gamma_L$ with $k = s_{\gamma^*N}/(2\sqrt{s_{\text{NN}}})$ and $\gamma_L = \sqrt{s_{\text{NN}}}/(2m_N)$. For the Pb nucleus, we use a Woods-Saxon radius of $R_A = 6.62$ fm, and $K_0(w)$ and $K_1(w)$ are the modified Bessel functions. In Pb + Pb UPCs at $\sqrt{s_{\text{NN}}} = 5.02$ TeV, the $\sqrt{s_{\gamma^*N}}$ of the $\gamma^* + \text{Pb}$ collisions fluctuates from 0 up to ~ 894 GeV [25,27,40]. To make theoretical predictions with fluctuating $\sqrt{s_{\gamma^*N}}$, we employ the Monte Carlo 3D-GLAUBER initial-state model [37,41] to provide dynamical source terms of energy, momentum, and net baryon density for the subsequent relativistic viscous hydrodynamics evolution. This model can quantitatively reproduce particle production in heavy-ion and asymmetric collisions at different collision energies [37].

We treat the virtual photon as a vector meson and sample two ‘‘partonic participants’’ inside the vector meson to capture its geometry shape fluctuations. Those partonic participants are understood as hot spots composed of valence quarks and their associated gluon cloud. Since the constituent parton distribution functions (PDF) for vector mesons are not well constrained by experiments, we parametrize the vector meson constituent PDF as $xv^{\text{VM}} = N_v x^\alpha (1-x)^\beta$, with $\alpha = \beta = 2$, and N_v is subject to the constraint $\int_0^1 v^{\text{VM}}(x) dx = 1$. This parametrization is

consistent with the shape of the π meson’s valence PDF at low Q^2 [42]. The two partonic participants do not carry all the energy and momentum of the vector meson. The remaining energy and momentum carried by sea quarks and soft gluons are attributed to a soft gluon cloud [37], which is allowed to participate in collisions with the Pb nucleus. We leave the exploration of UPC observables’ sensitivity to the vector meson PDF for future Letter. The transverse positions of the constituent partons are sampled from a 2D Gaussian, $\exp\{-[(x^2 + y^2)/2]Q^2\}$. We use a default $Q^2 = 0.0625 \text{ GeV}^2$ in our model calculations, which translates to a vector meson size of 0.8 fm, close to the values for ρ mesons in Ref. [43]. Below, we will explore the sensitivity of elliptic flow on the photon’s virtuality in $\gamma^* + \text{Pb}$ collisions.

The collective expansion of the produced dense nuclear matter and the evolution of the conserved net baryon current are simulated within a (3 + 1)D viscous hydrodynamic framework, MUSIC [44–47]. As the QGP droplet expands and transitions to the dilute hadronic phase, the fluid dynamic description is switched to a microscopic hadron transport model on a constant energy density surface at $e_{\text{sw}} = 0.20 \text{ GeV}/\text{fm}^3$. The hadronic transport phase is simulated by URQMD [48,49] for the hadron scatterings and resonance decays. The numerical simulations of this hybrid 3D-GLAUBER + MUSIC + URQMD model are carried out using the open-source IEBE-MUSIC framework [50]. All model parameters were fixed in Ref. [37] by calibrating minimum bias $p + p$ collisions. We adjust the specific shear viscosity to $\eta T/(e + P) = 0.08$ and the width of strings in the transverse plane $\sigma_x = 0.4$ fm to describe the p_T -integrated elliptic flow in $p + \text{Pb}$ collisions at 5.02 TeV [51]. The (rapidity-dependent) charged hadron multiplicity distributions studied in Ref. [37] are not changed by these adjustments.

Collectivity in $p + \text{Pb}$ and $\gamma^ + \text{Pb}$ collisions.*—The 3D-GLAUBER + MUSIC + URQMD model can successfully predict the rapidity distributions of particle production in relativistic nuclear collisions of different collision energy and nuclear species [37]. Using this predictive power, we compute the collectivity observables in $\gamma^* + \text{Pb}$ and $p + \text{Pb}$ collisions. We perform numerical simulations for these collisions in their center-of-mass frames. Because of the unequal energies of projectile and target in the laboratory frame, we need to apply a global rapidity shift to all final-state hadrons before comparing the results with experimental measurements. For $p + \text{Pb}$ collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV, the rapidity shift is $\Delta y = 0.465$ in the proton-going direction [52–54]. For $\gamma^* + \text{Pb}$ collisions, the rapidity shift is significantly larger because of the smaller energy of the γ^* . The $\Delta y = y_{\text{beam},\sqrt{s_{\gamma^*N}}} - y_{\text{beam},5.02 \text{ TeV}}$, and is in the Pb-going direction. For example, the $\Delta y = y_{\text{beam},\sqrt{s_{\gamma^*N}}} - y_{\text{beam},5.02 \text{ TeV}} = 1.725$ in the Pb-going direction for $\sqrt{s_{\gamma^*N}} = 894 \text{ GeV}$.




FIG. 1. The charged hadron pseudorapidity distributions $dN_{\text{ch}}/d\eta$ in 0%–90% $p + \text{Pb}$ and $\gamma^* + \text{Pb}$ collisions from the 3D-GLAUBER + MUSIC + URQMD simulations. The theoretical calculations are compared with experimental data from the ATLAS Collaboration in the laboratory frame [25,54].

Figure 1 shows our model results for the charged hadron pseudorapidity distributions $dN_{\text{ch}}/d\eta$ in 0%–90% $p + \text{Pb}$ collisions and in $\gamma^* + \text{Pb}$ collisions for the event class with $N_{\text{ch}} > 10$ and $0.4 < p_T < 5.0$ GeV in the $|\eta| < 2.5$ range. Following the ATLAS analysis [54], centrality classes in $p + \text{Pb}$ collisions are determined using the total transverse energy measured in the Pb-going direction of the forward calorimeter at $-4.9 < \eta < -3.1$. Our model reproduces the shape and magnitude of the $p + \text{Pb}$ $dN_{\text{ch}}/d\eta$ very well.

The predicted $dN_{\text{ch}}/d\eta$ in $\gamma^* + \text{Pb}$ collisions shows a strong asymmetry in the η direction, which clearly demonstrates the strong violation of longitudinal boost invariance. Implementing fluctuations of $\sqrt{s_{\gamma^*N}}$ according to Eq. (1) leads to a good description of the shape of the $dN_{\text{ch}}/d\eta$ distribution in Pb + Pb UPCs measured by the ATLAS Collaboration. Overall, it is remarkable that a final-state-dominated framework can predict the experimental charged hadron rapidity distribution in photonuclear events within 10%. We note that the ATLAS data on $dN_{\text{ch}}/d\eta$ in UPCs have been normalized to the value computed by DPMJET-III $\gamma + \text{Pb}$ at $\eta = 0$ [25]. We divide the ATLAS data of $dN_{\text{ch}}/d\eta$ in UPCs by 1.2 to normalize to our result at $\eta = 0$.

The good description of the charged hadron rapidity distributions provides a solid basis for us to quantitatively study flow observables with the same kinematic cuts as done in the experimental analysis. Figure 2 shows the identified particles' mean transverse momenta $\langle p_T \rangle$ as functions of charged hadron multiplicity in $p + \text{Pb}$ and $\gamma^* + \text{Pb}$ collisions. Compared with the ALICE $p + \text{Pb}$ measurements [52], the 3D-GLAUBER + MUSIC + URQMD framework reproduces the mass hierarchy of the $\langle p_T \rangle$ of pions, kaons, and protons as a result of the system's collective radial expansion. The mean p_T of pions and protons are overestimated by 10%, which can be




FIG. 2. Identified particle mean transverse momenta $\langle p_T \rangle$ as functions of charged hadron multiplicity in $p + \text{Pb}$ (dashed lines) and $\gamma^* + \text{Pb}$ (solid lines) collisions from the 3D-GLAUBER + MUSIC + URQMD framework. The $p + \text{Pb}$ results are compared to the experimental data from the ALICE Collaboration [52].

improved by including bulk viscous effects in the hydrodynamic evolution. Our model predicts that the identified particles' mean p_T in $\gamma^* + \text{Pb}$ collisions are very similar to those in $p + \text{Pb}$ collisions at the same charged hadron multiplicity. This is a consequence of using the same hot spot size in both systems, which leads to a similar amount of radial flow. Our mean p_T result provides a quantitative prediction, and experimental confirmation will be a strong indication that the system produced in $\gamma^* + \text{Pb}$ collisions experiences strong final-state effects.

In Fig. 3, we show the multiplicity dependence of the p_T -integrated anisotropic flow coefficients $v_2\{2\}$ and $v_3\{2\}$ computed with the scalar-product method, which uses two subevents with the kinematic cuts




FIG. 3. Charged hadron anisotropic flow coefficients $v_2\{2\}$ and $v_3\{2\}$ as functions of charged hadron multiplicity N_{ch} in $p + \text{Pb}$ (dashed lines) and $\gamma^* + \text{Pb}$ (solid lines) collisions at LHC energies from the 3D-GLAUBER + MUSIC + URQMD simulations. To compare with the ATLAS data [25,51], we compute the $v_n\{2\}$ coefficients using the appropriate kinematic cuts.

$-2.5 < \eta < -1.0$ and $1.0 < \eta < 2.5$ and $0.4 < p_T < 2$ GeV ($0.5 < p_T < 5.0$ GeV) for $\gamma^* + \text{Pb}$ ($p + \text{Pb}$) collisions. With the specific shear viscosity $\eta T/(e + P) = 0.08$ in hydrodynamic simulations, we fit the ATLAS v_n data for $p + \text{Pb}$ collisions [51]. The 3D-GLAUBER + MUSIC + URQMD framework can nicely reproduce the multiplicity dependence of the experimentally measured $v_n\{2\}$ in $p + \text{Pb}$ collisions. For $N_{\text{ch}} < 20$, the $v_n\{2\}$ drops because of the decreasing lifetime of the hydrodynamic phase. Extrapolating from $p + \text{Pb}$ to $\gamma^* + \text{Pb}$ collisions, our hydrodynamic calculations reproduce the hierarchy observed for the elliptic flow coefficient v_2 in $N_{\text{ch}} \in [20, 60]$ in the ATLAS data using the template fit method.

We have checked that the second-order eccentricity ε_2 in $\gamma^* + \text{Pb}$ initial states is very close to that in $p + \text{Pb}$ systems. However, we find that the longitudinal flow decorrelation is stronger in $\gamma^* + \text{Pb}$ collisions, which results in smaller $v_2\{2\}$ with $|\Delta\eta| > 2$. The reasons for the stronger longitudinal flow decorrelation in $\gamma^* + \text{Pb}$ collisions are (1) the smaller center-of-mass collision energy, which results in a narrower rapidity coverage of particle production, and (2) the larger rapidity shift between the center-of-mass and the lab frames, which increases the decorrelation effects for $|\eta| < 2.5$ in the lab frame. In other words, the initial transverse geometry is less important than the longitudinal structure in these two small systems.

This result underlines the importance of performing full (3 + 1)D simulations when quantitatively studying collectivity in small collision systems, and demonstrates that the elliptic flow hierarchy between $\gamma^* + \text{Pb}$ and $p + \text{Pb}$ collisions is compatible with a picture where final-state effects dominate the generation of momentum anisotropies.

Our model predicts that triangular flow in $\gamma^* + \text{Pb}$ collisions is smaller than that in $p + \text{Pb}$ collisions at the same charged hadron multiplicity, again because of the larger longitudinal decorrelation. Consequently, the ordering of $v_3\{2\}$ between $\gamma^* + \text{Pb}$ and $p + \text{Pb}$ collisions in our model is opposite to the ATLAS data, which show a larger $v_3\{2\}$ in $\gamma^* + \text{Pb}$ collisions. The magnitude of $v_3\{2\}$ in $\gamma^* + \text{Pb}$ collisions may be sensitive to vector meson's detailed substructure fluctuations.

Figure 4 shows our model comparison for the charged hadron p_T -differential elliptic flow $v_2(p_T)$ with the ATLAS measurements in $20 < N_{\text{ch}} < 60$ and $N_{\text{ch}} > 60$ $\gamma^* + \text{Pb}$ and $p + \text{Pb}$ collisions [25,51]. Our $v_2(p_T)$ result for the $N_{\text{ch}} > 60$ events in $p + \text{Pb}$ collisions shows excellent agreement with the ATLAS data, marking a good baseline to study the $v_2(p_T)$ in $\gamma^* + \text{Pb}$ collisions. Comparing this result with the one from the $20 < N_{\text{ch}} < 60$ multiplicity bin of $p + \text{Pb}$ collisions, we see a sizable suppression of $v_2(p_T)$ for $p_T > 1$ GeV in the lower multiplicity bin because of a shorter fireball lifetime.

The $v_2(p_T)$ in $\gamma^* + \text{Pb}$ collisions in the same $20 < N_{\text{ch}} < 60$ multiplicity bin is 10%–15% smaller than the $p + \text{Pb}$ $v_2(p_T)$ across all p_T values because of the larger




FIG. 4. The p_T -differential elliptic flow coefficient $v_2(p_T)$ of charged hadrons in $p + \text{Pb}$ and $\gamma^* + \text{Pb}$ collisions from the 3D-GLAUBER + MUSIC + URQMD simulations are compared to ATLAS data [25,51]. The $v_2(p_T)$ are calculated using the scalar-product method by imposing $|\Delta\eta| > 2$ between the particle of interest and the reference charged hadrons in $-2.5 < \eta < 2.5$ and $0.5 < p_T < 5.0$ GeV ($0.4 < p_T < 2.0$ GeV) for $p + \text{Pb}$ ($\gamma^* + \text{Pb}$) collisions.

longitudinal decorrelation with the reference flow angle in $\gamma^* + \text{Pb}$ collisions. Our model prediction agrees reasonably well with the ATLAS data for $p_T < 2.0$ GeV. The ATLAS UPC $v_2(p_T)$ decreases quickly as p_T increases above 1.6 GeV. This behavior is not seen in our calculations. For p_T above 2 GeV, other physics processes, such as quark recombination, which are not included here, start to be important for anisotropic flow coefficients [16,55].

Finally, we explore the sensitivity of $v_2(p_T)$ to the vector meson transverse size in $\gamma^* + \text{Pb}$ collisions. Because in the experiment the incoming γ^* 's virtuality Q^2 fluctuates from event to event, the projectile vector meson's average size also fluctuates as it is inversely proportional to Q^2 . This adds to the geometric fluctuations resulting from the random positions of the two hot spots (at fixed average size). We estimate the uncertainty on the final $v_2(p_T)$ from such Q^2 fluctuations by running simulations at different values of Q^2 . We perform this exercise at fixed center-of-mass energy, which requires less statistics. As the $v_2\{2\}$ depends only weakly on the collision energy (see Supplemental Material [56]) results for fluctuating energies are expected to be similar. Figure 5 shows that vector mesons with large virtuality result in smaller elliptic flow coefficients because there is less transverse space for the geometry to fluctuate and the average ellipticities are smaller. Increasing the virtuality from 0.04 to 0.25 GeV², the $v_2(p_T)$ in $\gamma^* + \text{Pb}$ decreases monotonically (we remind the reader that the default value is $Q^2 = 0.0625$ GeV²). The overall relative variation is about 30%. Future experiments at an Electron-Ion Collider will provide direct access to the photon's virtuality. Therefore, one will be able to systematically test the predictions from




FIG. 5. The p_T -differential elliptic flow coefficient $v_2(p_T)$ of charged hadrons in $\gamma^* + \text{Pb}$ collisions from the 3D-GLAUBER + MUSIC + URQMD simulations with different photon virtualities are compared to the ATLAS data [25].

the hydrodynamic framework by measuring elliptic anisotropies for different photon virtualities.

Summary.—In this Letter, we have carried out the first dynamical (3 + 1)D simulations that quantitatively study the collectivity in $p + \text{Pb}$ and ultraperipheral $\text{Pb} + \text{Pb}$ collisions at LHC energies within the relativistic hydrodynamic approach. Because these asymmetric collision systems do not have any reasonably wide rapidity window with uniform particle production, it is necessary to go beyond the high-energy Bjorken paradigm and simulate these collisions in full 3D.

The 3D-GLAUBER + MUSIC + URQMD model successfully describes the charged hadron pseudorapidity distributions in 0%–90% $p + \text{Pb}$ and $\gamma^* + \text{Pb}$ collisions in ultraperipheral $\text{Pb} + \text{Pb}$ events. This good agreement provides a baseline for studying momentum anisotropies and their correlations in rapidity.

Although $\gamma^* + \text{Pb}$ events have slightly larger initial eccentricities compared to those in $p + \text{Pb}$ collisions, they also exhibit a stronger longitudinal flow decorrelation, which results in a smaller elliptic momentum anisotropy in $\gamma^* + \text{Pb}$ collisions in a given multiplicity bin. This result highlights the crucial role of nontrivial longitudinal dynamics in describing the hierarchy of the elliptic flow in $p + \text{Pb}$ and $\text{Pb} + \text{Pb}$ UPC events.

The simultaneous description of elliptic flow coefficients in $p + \text{Pb}$ and $\gamma^* + \text{Pb}$ collisions in our model supports the hypothesis that strong final-state interactions can be the dominant source of collectivity in these small systems. Meanwhile, the fact that the model predicts the opposite trend in $v_3\{2\}$ compared to the experimental data suggests that one needs more careful studies of the vector meson's geometry and the interplay with initial-state correlations.

Our study provides a theoretical framework that can bridge the phenomenological studies of collectivity in

relativistic heavy-ion collisions with electron + nucleus collisions at the future Electron-Ion Collider, which will be able to test the picture we have established, namely the dominance of hydrodynamic response to initial spatial geometry in generating azimuthal anisotropies in virtual photon-nucleus collisions, with much more precision.

We thank Jiangyong Jia, Soumya Mohapatra, and Blair Seidlitz for providing the ATLAS data. W. Z. is supported by the National Science Foundation (NSF) under Grant No. ACI-2004571 within the framework of the XSCAPE project of the JETSCAPE collaboration. B. S. and C. S. are supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under DOE Contract No. DE-SC0012704 and Award No. DE-SC0021969, respectively. C. S. acknowledges support from a DOE Office of Science Early Career Award. This work is in part supported within the framework of the Beam Energy Scan Theory (BEST) Topical Collaboration and under Contract No. DE-SC0013460. This research was done using resources provided by the Open Science Grid (OSG) [57,58], which is supported by the National Science Foundation Award No. 2030508.

- [1] U. Heinz and R. Snellings, *Annu. Rev. Nucl. Part. Sci.* **63**, 123 (2013).
- [2] C. Gale, S. Jeon, and B. Schenke, *Int. J. Mod. Phys. A* **28**, 1340011 (2013).
- [3] C. Shen and U. Heinz, *Nucl. Phys. News* **25**, 6 (2015).
- [4] C. Aidala *et al.* (PHENIX Collaboration), *Phys. Rev. Lett.* **120**, 062302 (2018).
- [5] C. Aidala *et al.* (PHENIX Collaboration), *Nat. Phys.* **15**, 214 (2019).
- [6] W. Li, *Mod. Phys. Lett. A* **27**, 1230018 (2012).
- [7] K. Dusling, W. Li, and B. Schenke, *Int. J. Mod. Phys. E* **25**, 1630002 (2016).
- [8] J. L. Nagle and W. A. Zajc, *Annu. Rev. Nucl. Part. Sci.* **68**, 211 (2018).
- [9] J. L. Nagle, A. Adare, S. Beckman, T. Koblesky, J. Orjuela Koop, D. McGlinchey, P. Romatschke, J. Carlson, J. E. Lynn, and M. McCumber, *Phys. Rev. Lett.* **113**, 112301 (2014).
- [10] B. Schenke and R. Venugopalan, *Phys. Rev. Lett.* **113**, 102301 (2014).
- [11] C. Shen, J.-F. Paquet, G. S. Denicol, S. Jeon, and C. Gale, *Phys. Rev. C* **95**, 014906 (2017).
- [12] R. D. Weller and P. Romatschke, *Phys. Lett. B* **774**, 351 (2017).
- [13] H. Mäntysaari, B. Schenke, C. Shen, and P. Tribedy, *Phys. Lett. B* **772**, 681 (2017).
- [14] W. Zhao, Y. Zhou, H. Xu, W. Deng, and H. Song, *Phys. Lett. B* **780**, 495 (2018).
- [15] C. Bierlich and C. O. Rasmussen, *J. High Energy Phys.* **10** (2019) 026.
- [16] W. Zhao, C. M. Ko, Y.-X. Liu, G.-Y. Qin, and H. Song, *Phys. Rev. Lett.* **125**, 072301 (2020).

- [17] K. Dusling and R. Venugopalan, *Phys. Rev. D* **87**, 051502 (R) (2013).
- [18] T. Lappi, B. Schenke, S. Schlichting, and R. Venugopalan, *J. High Energy Phys.* **01** (2016) 061.
- [19] B. Schenke, S. Schlichting, P. Tribedy, and R. Venugopalan, *Phys. Rev. Lett.* **117**, 162301 (2016).
- [20] K. Dusling, M. Mace, and R. Venugopalan, *Phys. Rev. Lett.* **120**, 042002 (2018).
- [21] K. Dusling, M. Mace, and R. Venugopalan, *Phys. Rev. D* **97**, 016014 (2018).
- [22] B. Schenke, C. Shen, and P. Tribedy, *Phys. Lett. B* **803**, 135322 (2020).
- [23] G. Giacalone, B. Schenke, and C. Shen, *Phys. Rev. Lett.* **125**, 192301 (2020).
- [24] B. Schenke, *Rep. Prog. Phys.* **84**, 082301 (2021).
- [25] G. Aad *et al.* (ATLAS Collaboration), *Phys. Rev. C* **104**, 014903 (2021).
- [26] C. A. Bertulani, S. R. Klein, and J. Nystrand, *Annu. Rev. Nucl. Part. Sci.* **55**, 271 (2005).
- [27] A. J. Baltz, *Phys. Rep.* **458**, 1 (2008).
- [28] P. Bozek, A. Bzdak, and G.-L. Ma, *Phys. Lett. B* **748**, 301 (2015).
- [29] W. Ke, J. S. Moreland, J. E. Bernhard, and S. A. Bass, *Phys. Rev. C* **96**, 044912 (2017).
- [30] B. Schenke and S. Schlichting, *Phys. Rev. C* **94**, 044907 (2016).
- [31] C. Shen and S. Alzhrani, *Phys. Rev. C* **102**, 014909 (2020).
- [32] X.-Y. Wu and G.-Y. Qin, [arXiv:2109.03512](https://arxiv.org/abs/2109.03512).
- [33] F. S. Bemfica, M. M. Disconzi, V. Hoang, J. Noronha, and M. Radosz, *Phys. Rev. Lett.* **126**, 222301 (2021).
- [34] C. Shen and L. Yan, *Nucl. Sci. Tech.* **31**, 122 (2020).
- [35] C. Chiu and C. Shen, *Phys. Rev. C* **103**, 064901 (2021).
- [36] C. Plumberg, D. Almaalol, T. Dore, J. Noronha, and J. Noronha-Hostler, *Phys. Rev. C* **105**, L061901 (2022).
- [37] C. Shen and B. Schenke, *Phys. Rev. C* **105**, 064905 (2022).
- [38] J. J. Sakurai, *Ann. Phys. (N.Y.)* **11**, 1 (1960).
- [39] Y. Shi, L. Wang, S.-Y. Wei, B.-W. Xiao, and L. Zheng, *Phys. Rev. D* **103**, 054017 (2021).
- [40] R. Engel, *Z. Phys. C* **66**, 203 (1995).
- [41] C. Shen and B. Schenke, *Phys. Rev. C* **97**, 024907 (2018).
- [42] M. Aicher, A. Schafer, and W. Vogelsang, *Phys. Rev. Lett.* **105**, 252003 (2010).
- [43] J. R. Forshaw, R. Sandapen, and G. Shaw, *Phys. Rev. D* **69**, 094013 (2004).
- [44] B. Schenke, S. Jeon, and C. Gale, *Phys. Rev. Lett.* **106**, 042301 (2011).
- [45] B. Schenke, S. Jeon, and C. Gale, *Phys. Rev. C* **82**, 014903 (2010).
- [46] J.-F. Paquet, C. Shen, G. S. Denicol, M. Luzum, B. Schenke, S. Jeon, and C. Gale, *Phys. Rev. C* **93**, 044906 (2016).
- [47] G. S. Denicol, C. Gale, S. Jeon, A. Monnai, B. Schenke, and C. Shen, *Phys. Rev. C* **98**, 034916 (2018).
- [48] S. A. Bass *et al.*, *Prog. Part. Nucl. Phys.* **41**, 255 (1998).
- [49] M. Bleicher, E. Zabrodin, C. Spieles, S. A. Bass, C. Ernst, S. Soff, L. Bravina, M. Belkacem, H. Weber, H. Stöcker, and W. Greiner, *J. Phys. G* **25**, 1859 (1999).
- [50] The open-source IEBE-MUSIC overarching framework can be downloaded from <https://github.com/chunshen1987/iEBE-MUSIC>.
- [51] M. Aaboud *et al.* (ATLAS Collaboration), *Phys. Rev. C* **96**, 024908 (2017).
- [52] B. B. Abelev *et al.* (ALICE Collaboration), *Phys. Lett. B* **728**, 25 (2014).
- [53] V. Khachatryan *et al.* (CMS Collaboration), *Phys. Lett. B* **742**, 200 (2015).
- [54] G. Aad *et al.* (ATLAS Collaboration), *Eur. Phys. J. C* **76**, 199 (2016).
- [55] W. Zhao, W. Ke, W. Chen, T. Luo, and X.-N. Wang, *Phys. Rev. Lett.* **128**, 022302 (2022).
- [56] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.129.252302> for [brief description].more detail about the center-of-mass energy distribution, and investigates the collective flow coefficients at different center-of-mass energy in $\gamma^* + \text{Pb}$ collisions.
- [57] R. Pordes *et al.*, *J. Phys. Conf. Ser.* **78**, 012057 (2007).
- [58] I. Sfiligoi, D. C. Bradley, B. Holzman, P. Mhashilkar, S. Padhi, and F. Wurthwein, *WRI World Congress* **2**, 428 (2009).