

Coherent Many-Body Oscillations Induced by a Superposition of Broken Symmetry States in the Wake of a Quantum Phase Transition

Jacek Dziarmaga¹, Marek M. Rams¹, and Wojciech H. Zurek²

¹*Jagiellonian University, Institute of Theoretical Physics, Łojasiewicza 11, PL-30348 Kraków, Poland*

²*Theory Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA*

 (Received 23 February 2022; accepted 8 December 2022; published 23 December 2022)

It is now widely accepted that quenches through the critical region of quantum phase transitions result in post-transition states populated with topological defects—analogs of the classical topological defects. However, consequences of the very nonclassical fact that the state after a quench is a *superposition* of distinct, broken-symmetry vacua with different numbers and locations of defects have remained largely unexplored. We identify coherent quantum oscillations induced by such superpositions in observables complementary to the one involved in symmetry breaking. These oscillations satisfy Kibble-Zurek dynamical scaling laws with the quench rate, with an instantaneous oscillation frequency set primarily by the gap of the system. In addition to the obvious fundamental significance of a superposition of different broken symmetry states, quantum coherent oscillations can be used to verify unitarity and test for imperfections of the experimental implementations of quantum simulators.

DOI: [10.1103/PhysRevLett.129.260407](https://doi.org/10.1103/PhysRevLett.129.260407)

Motivation.—Studies of quenches through a symmetry-breaking quantum phase transition at a finite rate have been to a large extent focused on the generation of topological defects. This was clearly the first thing to do, as topological defects are stable and the obvious focus of interest in the classical (i.e., thermodynamic) nonequilibrium phase transitions. By contrast, quantum phase transitions inevitably lead to superpositions of the eigenstates of the post-transition Hamiltonian. Such superpositions (in, e.g., an atom) result in oscillations with the frequency given by the difference between the energies of the two levels involved, and the amplitude set by their initial occupancy. We show that superpositions of the post-transition eigenstates are inevitable in quantum phase transitions and exhibit analogous (many-body) coherent quantum oscillations. We characterize their appearance and properties in models where they can be investigated analytically or numerically, and where they should be accessible to experiments.

The obvious motivation for investigating collective oscillations of many-body systems is because they are there, and because they are a signature of the quantumness of the transition. Moreover, such an oscillatory behavior constitutes a sensitive probe of the imperfections of the experiment, including especially decoherence. We show that the form of the oscillations is simple when the energy levels of the many-body system are degenerate (as then the number of frequencies involved is small). When the degeneracies are lifted by the imperfections of the Hamiltonian (e.g., caused by its implementation), dephasing will result in the loss of coherence. Furthermore, decoherence caused by imperfect isolation of the system will result in nonunitary evolution causing a further gradual

loss of coherence. Therefore, such oscillations can serve as a diagnostic tool to assess how accurate—and especially how quantum—is the implementation of the transition in the emulation experiments: There are now examples of quantum phase transitions that are both solvable and experimentally accessible, creating appealing possibilities to use the exact many-body time-dependent solutions to benchmark experimental implementations. The post-transition oscillations should be relatively easy to prepare and detect in contrast to the more challenging nonlocal “double slit-like” superpositions of topological defects [1].

Kibble-Zurek mechanism.—The Kibble-Zurek mechanism (KZM) has its roots in cosmological symmetry-breaking phase transitions [2]. Kibble considered cooling Universe where causally disconnected regions independently select broken symmetry vacua. This mosaic of broken symmetry domains leads to topologically nontrivial configurations. The extent of such domains is limited by the size of the causal horizon.

This cosmological constraint is not relevant for laboratory experiments. Therefore, a dynamical theory for the continuous phase transitions was proposed and developed [3,4]. KZM employs equilibrium critical exponents to predict the scaling of the defects density as a function of the quench rate. It has been verified in numerous simulations [5–16] and condensed matter experiments [17–41]. Topological defects are central in those studies, as they can persist despite dissipation inevitable in thermodynamic systems.

The quantum version of KZM (QKZM) considers quenches across quantum critical points. It has been developed [42–78] and put to experimental tests [79–89].

Recent experiments target the exactly solvable quantum Ising chain in the transverse field, employing simulators based on Rydberg atoms [87] and superconducting qubits [90]. Scaling of the resulting defect densities appears to be consistent with the QKZM predictions [43–45]. Ongoing experimental developments [91–94] open the possibility to study the quantum dynamics in two-dimensional systems.

Of course, by the time defects are counted, quantum superpositions that should be present in the post-transition state are long gone. Thus, the *quantumness* of phase transition dynamics has not been, as yet, certified in the experiments. Indeed—as approximate scalings observed are not a unique fingerprint of the defect formation mechanism, and it is not clear at what stage the systems used in the experiments decohere and become effectively classical—it would be desirable to directly verify quantumness of the phase transition dynamics. Coherent oscillations we are describing offer that possibility. They can also be used to benchmark quantumness of the hardware used in (e.g., adiabatic) quantum computing.

A smooth ramp crossing the critical point at time t_c can be linearized in its vicinity as

$$\epsilon(t) = \frac{t - t_c}{\tau_Q}, \quad (1)$$

where ϵ measures the distance from the quantum critical point and quench rate is given by τ_Q . The system is prepared in the ground state far from the critical point. The initial evolution adiabatically follows the time-dependent Hamiltonian. This approximate adiabaticity fails at time $-\hat{t}$ before t_c when the reaction rate of the system (set by the gap) becomes comparable to the instantaneous relative ramp rate, namely, $\Delta \propto |\epsilon|^{\nu} \propto |\dot{\epsilon}/\epsilon| = 1/|t|$. This leads to the characteristic timescale

$$\hat{t} \propto \tau_Q^{z\nu/(1+z\nu)}, \quad (2)$$

where z is dynamical critical exponent, and ν is correlation length exponent [3]. In the adiabatic-impulse-adiabatic scenario, the ground state at $-\hat{t} = -\hat{t}/\tau_Q \propto -\tau_Q^{-1/(1+z\nu)}$ fluctuating on a scale set at $-\hat{t}$ survives until $+\hat{t}$, and the correlation length,

$$\hat{\xi} \propto \tau_Q^{\nu/(1+z\nu)}, \quad (3)$$

becomes imprinted for the subsequent adiabatic evolution. This oversimplified scenario correctly predicts the scaling dependence of the characteristic length and timescales on τ_Q . They naturally appear in KZM dynamical scaling hypothesis [95–97]. For an observable \mathcal{O} ,

$$\hat{\xi}^{\Delta_{\mathcal{O}}} \langle \psi(t) | \mathcal{O}_r | \psi(t) \rangle = F_{\mathcal{O}}((t - t_c)/\hat{\xi}^z, r/\hat{\xi}), \quad (4)$$

where $|\psi(t)\rangle$ is the state of the system, $\Delta_{\mathcal{O}}$ is the scaling dimension, $F_{\mathcal{O}}$ is a nonuniversal scaling function, and r is a distance in, e.g., a correlation function. It is expected to hold in the vicinity of the critical point, for t between $\pm\hat{t}$.

In the following, we employ the paradigmatic Ising Hamiltonian in a transverse field,

$$H(t) = -J(t) \sum_{\langle m,n \rangle} \sigma_m^z \sigma_n^z - g(t) \sum_m \sigma_m^x. \quad (5)$$

Here, σ_m^x , σ_m^y , and σ_m^z denote the Pauli matrices on lattice site m , and interactions that are between neighboring sites, $\langle m, n \rangle$. We consider three lattice geometries: (i) an integrable one-dimensional (1D) chain where each site has two neighbors, and two nonintegrable models where each site has four neighbors: (ii) a 1D ladder where sites that are next-nearest neighbors in a chain become adjacent, and (iii) a two-dimensional square lattice geometry (2D). We pictorially represent those lattice geometries as insets in the figures.

Oscillations in one dimension.—We begin with the 1D version [43,45,51,97,98] where we traditionally set $J = 1$ and ramp the transverse field,

$$g(t) = g_c[1 - \epsilon(t)] = g_c - g_c(t - t_c)/\tau_Q, \quad (6)$$

from $t = -\infty$ in the limit of strong field, across the critical point at $g(t_c) = g_c = 1$, to $g(t_s) = 0$ where the transverse field vanishes. The Jordan-Wigner transformation maps the model to a set of independent two-level Landau-Zener systems that can be solved analytically. In particular, the final density of excited quasiparticles or kinks scales like [43,45,99]

$$n \approx \frac{1}{2\pi\sqrt{2\tau_Q}} \propto \hat{\xi}^{-1}, \quad (7)$$

consistent with the critical exponents $z = \nu = 1$. The average defect density (accessed by counting them in the experiments to date) is a very superficial characterization of the final state, which, in fact, should be—prior to the kink count—a quantum superposition of different numbers [51,66] and correlated locations of kinks [77,98].

Breaking with tradition, we do not focus on kinks but rather on the transverse magnetization, σ^x , that does not commute with the kink observables. Its expectation value during and after crossing the critical point is shown in Fig. 1, where we consider linear ramps (6) with several values of the quench time. All the ramps stop at $g = 0$, allowing the system to freely evolve with a purely ferromagnetic Hamiltonian for $t > t_s$. In accordance with the general QKZM scaling hypothesis (4), for slow enough τ_Q the transverse magnetization in the vicinity of the critical point should satisfy




FIG. 1. Coherent oscillations after a quench through a quantum critical point. In panel (a), we show a quench protocol in the 1D transverse-field Ising model, where we allow the system to freely evolve upon reaching zero transverse field at time $t = t_s$. In (b), we track transverse magnetization in the 1D model, where the coherent oscillations for $t > t_s$ are apparent. Their origin can be traced back to the point reached at t_c , see panel (c), where we show data collapse consistent with the dynamical scaling hypothesis in Eq. (8). There is a gradual decrease of oscillation amplitude during the ramp through the ferromagnetic phase, setting the nonzero amplitude observed for $t > t_s$ [note that panel (c) shows only $t < t_s$].

$$\hat{\xi}^{\Delta_x} [\langle \sigma^x(t) \rangle - \langle \sigma^x \rangle_{GS}] = F_{\sigma^x} [(t - t_c) / \hat{t}]. \quad (8)$$

Here $\langle \sigma^x \rangle_{GS}$ is transverse magnetization in the instantaneous ground state for transverse field $g(t)$, and the scaling dimension $\Delta_x = 1$ for a 1D chain. As we can see in Fig. 1(c), the KZM-rescaled plots for different quench timescales τ_Q collapse to a common scaling function. In this integrable case good collapse extends beyond $+\hat{t}$. The function is oscillatory with an instantaneous frequency dominated by twice the quasiparticle gap as two quasiparticles with opposite quasimomenta are the relevant excitation. The amplitude of the oscillations slowly decays with the scaled time, partly due to a dephasing by a

nontrivial quasiparticle dispersion and partly due to the adiabatic evolution of the excited Bogoliubov modes.

The ramps in Fig. 1(a) terminate at $g = 0$ where the transverse magnetization in the ground state is zero: $\langle \sigma^x \rangle_{GS} = 0$. Therefore, $F(\infty) \hat{\xi}^{-1} \propto \tau_Q^{-1/2}$ is the initial transverse magnetization for the subsequent free evolution with $g = 0$, where

$$\langle \sigma_m^x(\tau) \rangle = \langle e^{i\tau H} \sigma_m^x e^{-i\tau H} \rangle_{t_s} = A_0 + A_4 \cos(4\tau + \tilde{\phi}), \quad (9)$$

as each site is uniformly coupled to 2 neighbors. Here, we introduce

$$\tau = t - t_s > 0,$$

as the duration of free evolution with $g = 0$. As we can see there is a constant term plus oscillations with a single frequency. The amplitudes are determined from expectation values in the state at $t = t_s$ at the end of the linear ramp and the beginning of the free evolution [99].

From the exact solution [99], we can extract an asymptotic form for $\tau_Q \gg 1$:

$$\langle \sigma_m^x(\tau) \rangle = 2n + 2Cdn^2 \cos(4\tau + \phi) - \pi^2 n^2 \sin(4\tau). \quad (10)$$

Here, $A_0 = 2n$ is set by the density of kinks in Eq. (7), which is conserved for $t > t_s$. The amplitude

$$A_4^{\text{linear}} = n^2 \sqrt{\pi^4 + 4\pi^2 C d \sin \phi + 4C^2 d^2}, \quad (11)$$

where $C \approx 57\sqrt{6\pi}/80$ is a numerical constant, ϕ is a phase accumulated by the KZ-excited quasiparticles [99], and

$$d = [1 + (3 \ln \tau_Q / 4\pi)^2]^{-3/4} < 1 \quad (12)$$

is a factor due to dephasing of the KZ excitations by their nontrivial dispersion. The constant term and the amplitude are plotted in Fig. 2(a) as functions of the quench time τ_Q .

As we can see, the amplitude is not a simple power law in τ_Q . Irregularities originate from interference between the two oscillatory contributions to (10), from the KZ excitation near the critical point, $\propto \cos(4\tau + \phi)$, and from the abrupt termination of the linear ramp at $g = 0$, $\propto \sin(4\tau)$. To focus on KZ oscillations we eliminate the non-KZ oscillations [104] by using, instead of the all-linear ramp in Eq. (6), a smoother version,

$$\tilde{\epsilon}(t) = \frac{t - t_c}{\tau_Q} - \frac{4}{27} \left(\frac{t - t_c}{\tau_Q} \right)^3, \quad (13)$$

replacing $\epsilon(t)$ in Eq. (6). This protocol starts in the ground state at $g(t_c - \frac{3}{2}\tau_Q) = 2$, and terminates at $g(t_s) = 0$, for $t_s = t_c + \frac{3}{2}\tau_Q$, with a zero time derivative, $\dot{g}(t_s) = 0$. This leads to pure post-KZ oscillation amplitude,




FIG. 2. Scaling of transverse magnetization oscillations amplitudes. In (a), for a 1D chain, we show constant contribution, A_0 (blue line), and the amplitudes of coherent oscillations, A_4 , as a function of τ_Q . We compare the analytical formulas in Eqs. (10) and (14) (lines) with the corresponding exact numerical results (points). We show the results for two quench protocols of the same linear slope $\propto \tau_Q^{-1}$ at the critical point: the first protocol in Eq. (6) stops abruptly at $g = 0$ (green line indicates the amplitude of oscillations), and the second one in Eq. (13) reaches $g = 0$ smoothly (red). The protocols are shown in the inset. In (b), for the ladder, the constant term B_0 closely follows the measured density of excitations, n . The latter follows an expected scaling for a model in the same universality class as a 1D chain, $n \sim \tau_Q^{-1/2}$. The dominant oscillatory contribution, B_8 , can be fitted with $B_8 \sim \tau_Q^{-0.69}$, though we expect that logarithmic corrections from dephasing contribute to the decay of B_8 . Note that in (a) the red line would be consistent with a power law and exponent -1.18 while the analytical solution shows $n^2 \sim \tau_Q^{-1}$ behavior times a logarithmic correction. In (c), we show the data for the 2D square lattice where the dominant contributions, B_0 and B_8 , closely follow the measured excitation density for available times.

$$A_4^{\text{smooth}} = 2Cd n^2, \quad (14)$$

that scales simply as $n^2 \propto \tau_Q^{-1}$, with a logarithmic correction brought in by the dephasing factor (see Fig. 2). The latter is slightly reduced, replacing $\ln \tau_Q$ with $0.2164 + \ln \tau_Q$ in Eq. (12), as the approach to $g = 0$ makes the smooth ramp longer. However, the reduction is negligible when $0.2164 \ll \ln \tau_Q$, because the extra time needed for the smooth ending of the ramp is spent mostly near $g = 0$, where the quasiparticle dispersion is almost flat, and there is little extra dephasing.

The smooth ramp is not the only way to eliminate non-KZ oscillations. For instance, an imperfect termination of

the linear ramp at a finite $g_f \ll 1$ (instead of 0) results in a gradual suppression of the oscillations with time. The small finite transverse field means that the quasiparticle dispersion is nontrivial although almost flat. The non-KZM excitations that span all quasi-momenta, dephase after time $\propto 1/g_f$. On the other hand, the influence on the KZ-part appears in the dephasing factor, replacing $\ln \tau_Q$ with $\ln \tau_Q - g_f^2 + 2\tau g_f/\tau_Q$ in Eq. (12). The KZ-part that originates from small quasimomenta modes, becomes suppressed when $\tau \gg \tau_Q/g_f$. For large enough τ_Q , it becomes much larger than the dephasing time of the non-KZ part, thus opening a time window when the non-KZ oscillations are suppressed but the KZ ones are not. It highlights the stability of KZ-related oscillations.

Oscillations in nonintegrable systems.—Qualitatively similar results can be obtained for nonintegrable systems though they make us resort to numerical simulations, see Fig. 3. For a 1D ladder, we use uniform matrix product




FIG. 3. Oscillations in nonintegrable systems. In (a), we show scaled transverse magnetization (8) in the function of scaled time in the 1D ladder geometry. The critical point of the model is in the same universality class as a 1D chain, with $\nu = z = \Delta_x = 1$. For large enough τ_Q , the plots collapse in the vicinity of the critical point to a single scaling function F_{σ^x} that exhibits oscillatory behavior. In (b), the corresponding data for the 2D transverse Ising model with the scaling dimension $\Delta_x \simeq 1.41$. The values of τ_Q are limited, from below, to be in the scaling limit at the critical point and, from above, to avoid finite-size effects in a finite lattice [78]. The best collapse of complementary quantities used $\xi = \tau_Q^{0.36}$ [78] for similar range of τ_Q 's. Solid lines indicate thermodynamic limit results of iPEPS [107], which become unstable for times longer than shown. Dashed lines indicate the MPS results measured in the center of a finite 11×11 lattice.

states (uMPS) [105] for a system in thermodynamic limit, and in 2D either the MPS [106] on a 11×11 lattice or the iPEPS in the thermodynamic limit [107]. We employ a protocol that is gradually turning on the Ising terms while turning off the transverse field [108],

$$g(t)/g_c = [1 - \epsilon(t)]/2, \quad J(t) = [1 + \epsilon(t)]/2. \quad (15)$$

We use the linear ramp in Eq. (1) for the ladder and a smooth ramp in Eq. (13) for the square lattice. The models exhibit phase transitions (for $J = 1$) at $g_c \approx 3.04438$ in two dimensions [109] and we identify $g_c \approx 2.4785$ for the ferromagnetic ladder.

After the ramp ends, at $g = 0$, the oscillations continue as

$$\langle \sigma_m^x(\tau) \rangle = B_0 + B_4 \cos(4\tau + \phi_4) + B_8 \cos(8\tau + \phi_8), \quad (16)$$

with a constant term and two frequencies of oscillations, resulting from an uniform Ising coupling of a site to 4 neighboring sites [99]. The amplitudes are shown in Fig. 2.

Figure 3 is testing the scaling hypothesis (8) for the nonintegrable models. With increasing τ_Q the plots tend to an oscillatory scaling function in the vicinity of the critical point even though in two dimensions, due to the growth of entanglement with increasing τ_Q , our simulations are limited to relatively fast transitions (i.e., quench times where the integrable 1D Ising also exhibits discrepancies with the limiting slow quench behavior).

Conclusion.—The postquench state is a superposition of different numbers of kinks (excited bonds). Two manifolds of eigenstates that differ by m excited bonds ($m = 2$ for a chain, and $m = 4$ for a ladder) result in oscillations

$$|\dots\downarrow\downarrow\downarrow\downarrow\downarrow\dots\rangle + \mathcal{A}|\dots\downarrow\downarrow\uparrow\downarrow\downarrow\dots\rangle e^{-2mit}. \quad (17)$$

This is the most obvious quantum signature of the consequences of the quantum phase transition.

For a chain the probability of a single-spin flip is $|\mathcal{A}|^2 \propto n^4$ (with a logarithmic correction) in agreement with [98,110] where antibunching of kinks makes it decay faster than n^2 . For a ladder we fit $|\mathcal{A}|^2 \propto \tau_Q^{-1.1}$. In both cases the amplitude of the oscillations follows as a square root of the probability. In two dimensions, the higher oscillation frequency, 8, similarly comes from isolated spin flips similar to (17), while the lower frequency, 4, and the constant term are due to spin flips adjacent to (coarse) domain walls. In 1D cases the dominant energy eigenvalues have nearly the same separation (a multiple of the gap) so the oscillation occurs with an essentially single (time dependent) frequency. In the 2D case the picture becomes slightly more complicated, but the few frequencies are still controlled by the gap size.

The secular part of the response to the quench follows from the same treatment and is also quantum, but the

oscillatory part is a more compelling signature of the quantumness of the transition. Coherent oscillations are vulnerable to decoherence (see Ref. [1] for related discussion) and to imperfect implementation of the Hamiltonian. Decoherence that einselects broken symmetry states is plausible in many-body systems. It will localize kinks while suppressing oscillations, as do the measurements aimed at testing KZM performed to date. Pointer observable is einselected at least in part by the system-environment coupling [111], so, e.g., “quantum limit of decoherence” that favors energy eigenstates [112], is also possible.

This research was funded by the National Science Centre (NCN), Poland, under project 2021/03/Y/ST2/00184 within the QuantERA II Programme that has received funding from the European Union’s Horizon 2020 research and innovation programme under Grant Agreement No. 101017733 (J.D.), NCN under project 2020/38/E/ST3/00150 (MMR), and Department of Energy under the Los Alamos National Laboratory LDRD Program (W.H.Z.). W.H.Z. was also supported by the U.S. Department of Energy, Office of Science, Basic Energy Sciences, Materials Sciences and Engineering Division, Condensed Matter Theory Program.

-
- [1] J. Dziarmaga, W. H. Zurek, and M. Zwolak, *Nat. Phys.* **8**, 49 (2012).
 - [2] T. W. B. Kibble, *Phys. Rep.* **67**, 183 (1980); *Phys. Today* **60**, No. 9, 47 (2007).
 - [3] W. H. Zurek, *Nature (London)* **317**, 505 (1985); *Acta Phys. Pol. B* **24**, 1301 (1993); *Phys. Rep.* **276**, 177 (1996).
 - [4] A. del Campo and W. H. Zurek, *Int. J. Mod. Phys. A* **29**, 1430018 (2014).
 - [5] P. Laguna and W. H. Zurek, *Phys. Rev. Lett.* **78**, 2519 (1997).
 - [6] A. Yates and W. H. Zurek, *Phys. Rev. Lett.* **80**, 5477 (1998).
 - [7] J. Dziarmaga, P. Laguna, and W. H. Zurek, *Phys. Rev. Lett.* **82**, 4749 (1999).
 - [8] N. D. Antunes, L. M. A. Bettencourt, and W. H. Zurek, *Phys. Rev. Lett.* **82**, 2824 (1999).
 - [9] L. M. A. Bettencourt, N. D. Antunes, and W. H. Zurek, *Phys. Rev. D* **62**, 065005 (2000).
 - [10] W. H. Zurek, L. M. A. Bettencourt, J. Dziarmaga, and N. D. Antunes, *Acta Phys. Pol. B* **31**, 2937 (2000), <https://www.actaphys.uj.edu.pl/R/31/12/2937/pdf>.
 - [11] M. Uhlmann, R. Schützhold, and U. R. Fischer, *Phys. Rev. Lett.* **99**, 120407 (2007); *Phys. Rev. D* **81**, 025017 (2010); *New J. Phys.* **12**, 095020 (2010).
 - [12] E. Witkowska, P. Deuar, M. Gajda, and K. Rzażewski, *Phys. Rev. Lett.* **106**, 135301 (2011).
 - [13] A. Das, J. Sabbatini, and W. H. Zurek, *Sci. Rep.* **2**, 352 (2012).
 - [14] J. Sonner, A. del Campo, and W. H. Zurek, *Nat. Commun.* **6**, 7406 (2015).

- [15] P. M. Chesler, A. M. García-García, and H. Liu, *Phys. Rev. X* **5**, 021015 (2015).
- [16] I.-K. Liu, J. Dziarmaga, S.-C. Gou, F. Dalfovo, and N. P. Proukakis, *Phys. Rev. Res.* **2**, 033183 (2020).
- [17] I. Chuang, R. Durrer, N. Turok, and B. Yurke, *Science* **251**, 1336 (1991).
- [18] M. J. Bowick, L. Chandar, E. A. Schiff, and A. M. Srivastava, *Science* **263**, 943 (1994).
- [19] V. M. H. Ruutu, V. B. Eltsov, A. J. Gill, T. W. B. Kibble, M. Krusius, Y. G. Makhlin, B. Plaçais, G. E. Volovik, and W. Xu, *Nature (London)* **382**, 334 (1996).
- [20] C. Bäuerle, Y. M. Bunkov, S. N. Fisher, H. Godfrin, and G. R. Pickett, *Nature (London)* **382**, 332 (1996).
- [21] R. Carmi, E. Polturak, and G. Koren, *Phys. Rev. Lett.* **84**, 4966 (2000).
- [22] R. Monaco, J. Mygind, and R. J. Rivers, *Phys. Rev. Lett.* **89**, 080603 (2002).
- [23] A. Maniv, E. Polturak, and G. Koren, *Phys. Rev. Lett.* **91**, 197001 (2003).
- [24] L. E. Sadler, J. M. Higbie, S. R. Leslie, M. Vengalattore, and D. M. Stamper-Kurn, *Nature (London)* **443**, 312 (2006).
- [25] C. N. Weiler, T. W. Neely, D. R. Scherer, A. S. Bradley, M. J. Davis, and B. P. Anderson, *Nature (London)* **455**, 948 (2008).
- [26] R. Monaco, J. Mygind, R. J. Rivers, and V. P. Koshelets, *Phys. Rev. B* **80**, 180501(R) (2009).
- [27] D. Golubchik, E. Polturak, and G. Koren, *Phys. Rev. Lett.* **104**, 247002 (2010).
- [28] G. D. Chiara, A. del Campo, G. Morigi, M. B. Plenio, and A. Retzker, *New J. Phys.* **12**, 115003 (2010).
- [29] M. Mielenz, J. Brox, S. Kahra, G. Leschhorn, M. Albert, T. Schaetz, H. Landa, and B. Reznik, *Phys. Rev. Lett.* **110**, 133004 (2013).
- [30] S. Ulm, J. Roßnagel, G. Jacob, C. Degünther, S. T. Dawkins, U. G. Poschinger, R. Nigmatullin, A. Retzker, M. B. Plenio, F. Schmidt-Kaler, and K. Singer, *Nat. Commun.* **4**, 2290 (2013).
- [31] K. Pyka, J. Keller, H. L. Partner, R. Nigmatullin, T. Burgermeister, D. M. Meier, K. Kuhlmann, A. Retzker, M. B. Plenio, W. H. Zurek, A. del Campo, and T. E. Mehlstäubler, *Nat. Commun.* **4**, 2291 (2013).
- [32] S. C. Chae, N. Lee, Y. Horibe, M. Tanimura, S. Mori, B. Gao, S. Carr, and S.-W. Cheong, *Phys. Rev. Lett.* **108**, 167603 (2012).
- [33] S.-Z. Lin, X. Wang, Y. Kamiya, G.-W. Chern, F. Fan, D. Fan, B. Casas, Y. Liu, V. Kiryukhin, W. H. Zurek, C. D. Batista, and S.-W. Cheong, *Nat. Phys.* **10**, 970 (2014).
- [34] S. M. Griffin, M. Lilienblum, K. T. Delaney, Y. Kumagai, M. Fiebig, and N. A. Spaldin, *Phys. Rev. X* **2**, 041022 (2012).
- [35] S. Donadello, S. Serafini, M. Tylutki, L. P. Pitaevskii, F. Dalfovo, G. Lamporesi, and G. Ferrari, *Phys. Rev. Lett.* **113**, 065302 (2014).
- [36] S. Deutschländer, P. Dillmann, G. Maret, and P. Keim, *Proc. Natl. Acad. Sci. U.S.A.* **112**, 6925 (2015).
- [37] L. Chomaz, L. Corman, T. Bienaimé, R. Desbuquois, C. Weitenberg, S. Nascimbène, J. Beugnon, and J. Dalibard, *Nat. Commun.* **6**, 6162 (2015).
- [38] V. Yukalov, A. Novikov, and V. Bagnato, *Phys. Lett. A* **379**, 1366 (2015).
- [39] N. Navon, A. L. Gaunt, R. P. Smith, and Z. Hadzibabic, *Science* **347**, 167 (2015).
- [40] I.-K. Liu, S. Donadello, G. Lamporesi, G. Ferrari, S.-C. Gou, F. Dalfovo, and N. P. Proukakis, *Commun. Phys.* **1**, 24 (2018).
- [41] J. Rysti, J. T. Mäkinen, S. Autti, T. Kamppinen, G. E. Volovik, and V. B. Eltsov, *Phys. Rev. Lett.* **127**, 115702 (2021).
- [42] B. Damski, *Phys. Rev. Lett.* **95**, 035701 (2005).
- [43] W. H. Zurek, U. Dorner, and P. Zoller, *Phys. Rev. Lett.* **95**, 105701 (2005).
- [44] A. Polkovnikov, *Phys. Rev. B* **72**, 161201(R) (2005).
- [45] J. Dziarmaga, *Phys. Rev. Lett.* **95**, 245701 (2005).
- [46] J. Dziarmaga, *Adv. Phys.* **59**, 1063 (2010).
- [47] A. Polkovnikov, K. Sengupta, A. Silva, and M. Vengalattore, *Rev. Mod. Phys.* **83**, 863 (2011).
- [48] R. Schützhold, M. Uhlmann, Y. Xu, and U. R. Fischer, *Phys. Rev. Lett.* **97**, 200601 (2006).
- [49] H. Saito, Y. Kawaguchi, and M. Ueda, *Phys. Rev. A* **76**, 043613 (2007).
- [50] V. Mukherjee, U. Divakaran, A. Dutta, and D. Sen, *Phys. Rev. B* **76**, 174303 (2007).
- [51] F. M. Cucchietti, B. Damski, J. Dziarmaga, and W. H. Zurek, *Phys. Rev. A* **75**, 023603 (2007).
- [52] L. Cincio, J. Dziarmaga, M. M. Rams, and W. H. Zurek, *Phys. Rev. A* **75**, 052321 (2007).
- [53] A. Polkovnikov and V. Gritsev, *Nat. Phys.* **4**, 477 (2008).
- [54] K. Sengupta, D. Sen, and S. Mondal, *Phys. Rev. Lett.* **100**, 077204 (2008).
- [55] D. Sen, K. Sengupta, and S. Mondal, *Phys. Rev. Lett.* **101**, 016806 (2008).
- [56] J. Dziarmaga, J. Meisner, and W. H. Zurek, *Phys. Rev. Lett.* **101**, 115701 (2008).
- [57] B. Damski and W. H. Zurek, *Phys. Rev. Lett.* **104**, 160404 (2010).
- [58] C. De Grandi, V. Gritsev, and A. Polkovnikov, *Phys. Rev. B* **81**, 012303 (2010).
- [59] F. Pollmann, S. Mukerjee, A. G. Green, and J. E. Moore, *Phys. Rev. E* **81**, 020101(R) (2010).
- [60] B. Damski, H. T. Quan, and W. H. Zurek, *Phys. Rev. A* **83**, 062104 (2011).
- [61] W. H. Zurek, *J. Phys. Condens. Matter* **25**, 404209 (2013).
- [62] S. Sharma, S. Suzuki, and A. Dutta, *Phys. Rev. B* **92**, 104306 (2015).
- [63] A. Dutta and A. Dutta, *Phys. Rev. B* **96**, 125113 (2017).
- [64] D. Jaschke, K. Maeda, J. D. Whalen, M. L. Wall, and L. D. Carr, *New J. Phys.* **19**, 033032 (2017).
- [65] M. Białończyk and B. Damski, *J. Stat. Mech.* (2018) 073105.
- [66] A. del Campo, *Phys. Rev. Lett.* **121**, 200601 (2018).
- [67] R. Puebla, O. Marty, and M. B. Plenio, *Phys. Rev. A* **100**, 032115 (2019).
- [68] A. Sinha, M. M. Rams, and J. Dziarmaga, *Phys. Rev. B* **99**, 094203 (2019).
- [69] M. M. Rams, J. Dziarmaga, and W. H. Zurek, *Phys. Rev. Lett.* **123**, 130603 (2019).
- [70] S. Mathey and S. Diehl, *Phys. Rev. Res.* **2**, 013150 (2020).

- [71] M. Białończyk and B. Damski, *J. Stat. Mech.* (2020) 013108.
- [72] D. Sadhukhan, A. Sinha, A. Francuz, J. Stefaniak, M. M. Rams, J. Dziarmaga, and W. H. Zurek, *Phys. Rev. B* **101**, 144429 (2020).
- [73] B. S. Revathy and U. Divakaran, *J. Stat. Mech.* (2020) 023108.
- [74] D. Rossini and E. Vicari, *Phys. Rev. Res.* **2**, 023211 (2020).
- [75] K. Hódsági and M. Kormos, *SciPost Phys.* **9**, 55 (2020).
- [76] M. Białończyk and B. Damski, *Phys. Rev. B* **102**, 134302 (2020).
- [77] K. Roychowdhury, R. Moessner, and A. Das, *Phys. Rev. B* **104**, 014406 (2021).
- [78] M. Schmitt, M. M. Rams, J. Dziarmaga, M. Heyl, and W. H. Zurek, *Sci. Adv.* **8**, eabl6850 (2022).
- [79] L. E. Sadler, J. M. Higbie, S. R. Leslie, M. Vengalattore, and D. M. Stamper-Kurn, *Nature (London)* **443**, 312 (2006).
- [80] M. Anquez, B. A. Robbins, H. M. Bharath, M. Boguslawski, T. M. Hoang, and M. S. Chapman, *Phys. Rev. Lett.* **116**, 155301 (2016).
- [81] K. Baumann, R. Mottl, F. Brennecke, and T. Esslinger, *Phys. Rev. Lett.* **107**, 140402 (2011).
- [82] L. W. Clark, L. Feng, and C. Chin, *Science* **354**, 606 (2016).
- [83] D. Chen, M. White, C. Borries, and B. DeMarco, *Phys. Rev. Lett.* **106**, 235304 (2011).
- [84] S. Braun, M. Friesdorf, S. S. Hodgman, M. Schreiber, J. P. Ronzheimer, A. Riera, M. del Rey, I. Bloch, J. Eisert, and U. Schneider, *Proc. Natl. Acad. Sci. U.S.A.* **112**, 3641 (2015).
- [85] B. Gardas, J. Dziarmaga, W. H. Zurek, and M. Zwolak, *Sci. Rep.* **8**, 4539 (2018).
- [86] C. Meldgin, U. Ray, P. Russ, D. Chen, D. M. Ceperley, and B. DeMarco, *Nat. Phys.* **12**, 646 (2016).
- [87] A. Keesling, A. Omran, H. Levine, H. Bernien, H. Pichler, S. Choi, R. Samajdar, S. Schwartz, P. Silvi, S. Sachdev, P. Zoller, M. Endres, M. Greiner, V. Vuletić, and M. D. Lukin, *Nature (London)* **568**, 207 (2019).
- [88] Y. Bando, Y. Susa, H. Oshiyama, N. Shibata, M. Ohzeki, F. J. Gómez-Ruiz, D. A. Lidar, S. Suzuki, A. del Campo, and H. Nishimori, *Phys. Rev. Res.* **2**, 033369 (2020).
- [89] P. Weinberg, M. Tylutki, J. M. Rönkkö, J. Westerholm, J. A. Åström, P. Manninen, P. Törmä, and A. W. Sandvik, *Phys. Rev. Lett.* **124**, 090502 (2020).
- [90] A. D. King, S. Suzuki, J. Raymond *et al.*, *Nat. Phys.* **18**, 1324 (2022).
- [91] S. Ebadi, T. T. Wang, H. Levine, A. Keesling, G. Semeghini, A. Omran, D. Bluvstein, R. Samajdar, H. Pichler, W. W. Ho, S. Choi, S. Sachdev, M. Greiner, V. Vuletic, and M. D. Lukin, *Nature (London)* **595**, 227 (2021).
- [92] P. Scholl, M. Schuler, H. J. Williams, A. A. Eberharter, D. Barredo, K.-N. Schymik, V. Lienhard, L.-P. Henry, T. C. Lang, T. Lahaye, A. M. Läuchli, and A. Browaeys, *Nature (London)* **595**, 233 (2021).
- [93] G. Semeghini, H. Levine, A. Keesling, S. Ebadi, T. T. Wang, D. Bluvstein, R. Verresen, H. Pichler, M. Kalinowski, R. Samajdar, A. Omran, S. Sachdev, A. Vishwanath, M. Greiner, V. Vuletic, and M. D. Lukin, *Science* **374**, 1242 (2021).
- [94] K. J. Satzinger *et al.*, *Science* **374**, 1237 (2021).
- [95] M. Kolodrubetz, B. K. Clark, and D. A. Huse, *Phys. Rev. Lett.* **109**, 015701 (2012).
- [96] A. Chandran, A. Erez, S. S. Gubser, and S. L. Sondhi, *Phys. Rev. B* **86**, 064304 (2012).
- [97] A. Francuz, J. Dziarmaga, B. Gardas, and W. H. Zurek, *Phys. Rev. B* **93**, 075134 (2016).
- [98] R. J. Nowak and J. Dziarmaga, *Phys. Rev. B* **104**, 075448 (2021).
- [99] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.129.260407> for details of the calculations, which includes Refs. [100–103].
- [100] B. Damski and W. H. Zurek, *Phys. Rev. A* **73**, 063405 (2006).
- [101] P. Czarnik, J. Dziarmaga, and P. Corboz, *Phys. Rev. B* **99**, 035115 (2019).
- [102] J. Dziarmaga, *Phys. Rev. B* **104**, 094411 (2021).
- [103] J. Dziarmaga, *Phys. Rev. B* **105**, 054203 (2022).
- [104] To be more precise, making them higher order in powers of n .
- [105] L. Vanderstraeten, J. Haegeman, and F. Verstraete, *SciPost Phys. Lect. Notes* **7** (2019). [10.21468/SciPost-PhysLectNotes.7](https://arxiv.org/abs/10.21468/SciPost-PhysLectNotes.7)
- [106] J. Haegeman, C. Lubich, I. Oseledets, B. Vandereycken, and F. Verstraete, *Phys. Rev. B* **94**, 165116 (2016).
- [107] J. Dziarmaga, *Phys. Rev. B* **106**, 014304 (2022).
- [108] We considered a similar protocol for a 2D system in Ref. [78]. Comparing to that work, here, we rescale the Hamiltonian (and all timescales) by a factor of 2, to have $J = 1$ when $g = 0$.
- [109] H. W. J. Blöte and Y. Deng, *Phys. Rev. E* **66**, 066110 (2002).
- [110] J. Dziarmaga and M. M. Rams, *Phys. Rev. B* **106**, 014309 (2022).
- [111] W. H. Zurek, *Rev. Mod. Phys.* **75**, 715 (2003).
- [112] J. P. Paz and W. H. Zurek, *Phys. Rev. Lett.* **82**, 5181 (1999).