

Emergent Magnetomultipoles and Nonlinear Responses of a Magnetic Hopfion

Yizhou Liu¹, Hikaru Watanabe,¹ and Naoto Nagaosa^{1,2}

¹*RIKEN Center for Emergent Matter Science (CEMS), Wako, Saitama 351-0198, Japan*

²*Department of Applied Physics, University of Tokyo, Tokyo 113-8656, Japan*

 (Received 20 June 2022; revised 13 October 2022; accepted 6 December 2022; published 22 December 2022)

The three-dimensional emergent magnetic field \mathbf{B}^e of a magnetic hopfion gives rise to *emergent* magnetomultipoles in a similar manner to the multipoles of classical electromagnetic field. Here, we show that the nonlinear responses of a hopfion are characterized by its emergent magnetic toroidal moment $T_z^e = \frac{1}{2} \int (\mathbf{r} \times \mathbf{B}^e)_z dV$ and emergent magnetic octupole component $\Gamma^e = \int [(x^2 + y^2)B_z^e - xzB_x^e - yzB_y^e] dV$. The hopfion exhibits nonreciprocal dynamics (nonlinear hopfion Hall effect) under an ac driving current applied along (perpendicular to) the direction of T_z^e . The sign of nonreciprocity and nonlinear Hall angle is determined by the polarity and chirality of hopfion. The nonlinear electrical transport induced by a magnetic hopfion is also discussed. This Letter reveals the vital roles of emergent magnetomultipoles in nonlinear hopfion dynamics and could stimulate further investigations on the dynamical responses of topological spin textures induced by emergent electromagnetic multipoles.

DOI: [10.1103/PhysRevLett.129.267201](https://doi.org/10.1103/PhysRevLett.129.267201)

Topological spin textures (TSTs) like magnetic skyrmion have sparked intense attention due to their novel physical properties and attractive potential for applications [1,2]. Two fundamental quantities characterizing the spin dynamics and the transport phenomena associated with a spin texture are the vector and scalar spin chirality, which are related to the broken spatial-inversion \mathcal{I} and time-reversal symmetry \mathcal{T} , respectively. The scalar spin chirality acts as an effective real-space Berry curvature (the emergent magnetic field \mathbf{B}^e) on conduction electrons and its spatial integral reflects the topological skyrmion number. To date, tremendous efforts have been made to study scalar spin chirality related effects, leading to the discovery of many fascinating phenomena such as topological Hall effect [3–5], skyrmion Hall effect [6–8], emergent electromagnetic inductance [9,10], etc. Therefore, identifying such fundamental quantities relevant to the spin texture's dynamics and transport properties is of vital importance.

The multipole, which originated from classical electromagnetic field, has been a well-established fundamental concept in many branches of physics [11–13]. In condensed matter systems, the multipoles defined within different contents can serve as alluring sources of generating nonlinear and/or nonreciprocal responses [14–19]. The multipole concept can also be applied to the emergent magnetic field \mathbf{B}^e [1,9] associated with TSTs in which *emergent* magnetomultipoles may be further identified. However, unlike their counterparts, such as the momentum-space Berry curvature dipole and multipole [20–23], the role of emergent magnetomultipoles has not been well explored so far.

Recently, interest in realizing three-dimensional (3D) TSTs in magnetic materials has emerged with a focus on

their rich 3D topologies and veiled physical properties [24–28]. One example is the magnetic hopfion, a class of TST with diverse 3D structures [29–35] characterized by the Hopf charge Q_H [36–38]. Even the simplest form of hopfion (i.e., $Q_H = 1$) hosts sophisticated spin structure and 3D emergent magnetic field, leading to interesting dynamical properties such as the entangled current-driven dynamics [39,40] and field-driven resonance modes [41–44]. Hence, the 3D structure of \mathbf{B}^e of the hopfion provides a versatile platform to explore the role of emergent magnetomultipoles.

In this Letter, we show that the emergent magnetomultipoles of a $Q_H = 1$ magnetic hopfion characterizes the nonlinear responses of both the spin dynamics and electron transports. The toruslike hopfion possesses swirlinglike \mathbf{B}^e and it gives rise to an emergent toroidal moment \mathbf{T}^e like the classical toroidal moment of a torus solenoid [13,45,46]. It is found that the hopfion shows nonlinear nonreciprocal translation and rotation with respect to the direction of \mathbf{T}^e under an ac electrical current applied along \mathbf{T}^e . When the ac current is applied perpendicular to \mathbf{T}^e , a nonlinear hopfion Hall effect is identified where the hopfion also translates and rotates with respect to \mathbf{T}^e . These nonlinear dynamics of the hopfion are governed by the emergent toroidal moment T_z^e and the emergent magnetic octupole component Γ^e . On the other hand, we also demonstrate that the hopfion gives rise to nonreciprocal magnetoresistance and nonlinear Hall effect in its transport properties of conduction electrons. From these aspects, the emergent magnetomultipole serves as a new fundamental quantity that characterizes the nonlinear responses of spin textures.

The typical spin configuration of a hopfion $\mathbf{S}_0(\mathbf{r})$ with $Q_H = 1$ is shown in Fig. 1. The Hopf charge is defined as




FIG. 1. Schematic view of a $Q_H = 1$ magnetic hopfion, its emergent magnetic field (\mathbf{B}^e), and the corresponding emergent toroidal moment (\mathbf{T}^e). The half torus represents the equispin surface with $S_z = 0$. The rectangle plane shows the cross-sectional view of the hopfion onto the yz plane, which includes a skyrmion and antiskyrmion pair. In the color scheme, the black color in the core of the torus indicates $S_z = -1$ and the white color in the background indicates $S_z = 1$. The color wheel stands for in-plane spin directions.

$Q_H = \int \mathbf{A}^e(\mathbf{r}) \cdot \mathbf{B}^e(\mathbf{r}) dV$, where $B_i^e(\mathbf{r}) = \frac{1}{2} \epsilon_{ijk} \mathbf{S} \cdot (\partial_j \mathbf{S} \times \partial_k \mathbf{S})$ is the emergent magnetic field and \mathbf{A}^e is the corresponding vector potential ($\mathbf{B}^e = \nabla \times \mathbf{A}^e$). This hopfion structure is axially symmetric around the z axis. The associated emergent magnetic field exhibits a swirlinglike structure as indicated by the red looped arrow in Fig. 1. Although the symmetry of the hopfion results in a vanishing spatial-averaged emergent magnetic field $\langle \mathbf{B}^e(\mathbf{r}) \rangle = 0$, it defines a finite emergent toroidal moment $\mathbf{T}^e = \frac{1}{2} \int \mathbf{r} \times \mathbf{B}^e(\mathbf{r}) dV = T_z^e \hat{z}$ along the hopfion's symmetry axis (z axis) [45,46]. Note that \mathbf{T}^e appears as the integral of the Berry connection over the spin texture [see Eqs. (S13) and (S14) in the Supplemental Material [47]]. As a result, nonreciprocal responses related to \mathbf{T}^e are expected for the hopfion.

A previous study suggests that the current-driven dynamics of hopfion are effectively described by three dynamical modes: translation, rotation, and dilation [39]. For clarity, here we mainly focus on the dynamics of the hopfion related to the z axis, which is most relevant to the nonreciprocal responses. At position $\mathbf{r} = (x, y, z)$ and time t , the translation and rotation modes of the hopfion are expressed as $\mathbf{S}(\mathbf{r}, t) = \mathbf{S}_0[\hat{O}(\mathbf{r} - \mathbf{Z})]$, where $\mathbf{Z} = Z\hat{z}$ characterizes the translation of the hopfion along the z axis, and \hat{O} is the rotation operator. At infinitesimal rotation, for only the rotation around z axis, we have $\hat{O} \approx 1 - i\theta_z L_z$, where $\hat{L}_i = -i\epsilon_{ijk} r_j \partial_k$ is the angular momentum operator and θ_z is the rotation angle of hopfion around z axis. The dilation of the hopfion can be expressed as $\mathbf{S}(\mathbf{r}, t) = \mathbf{S}_0(\lambda \mathbf{r})$ and the static equilibrium dilation $\lambda_0 = 1$ at the initial state.

Based on these dynamical modes, a glance at the dynamics of the hopfion along its symmetry axis can be deduced from the spin's Berry phase term of the Lagrangian [6,50,51] $L_{BP} = \int (1 - \cos\theta) \dot{\phi} dV$, where θ and ϕ are the polar and azimuthal angle of the localized

spin \mathbf{S} with unit length. The variation of the spin's Berry phase term $\delta L_{BP} = \int \mathbf{S} \cdot \delta \mathbf{S} \times \dot{\mathbf{S}} dV$ can be expressed in terms of the slowly varying dynamical modes of the hopfion after integrating out the spin configuration,

$$\delta L_{BP}^z = (-T_z^e \dot{Z} + \Gamma^e \dot{\theta}_z) \delta \lambda, \quad (1)$$

where T_z^e is the emergent toroidal moment and $\Gamma^e = \int [(x^2 + y^2) B_z^e - xz B_x^e - yz B_y^e] dV$ is an emergent magnetic octupole component of the hopfion [52,53]. In Eq. (1), the translation and rotation are coupled to the dilation via T_z^e and Γ^e rather than the emergent magnetic field, whose spatial average vanishes for the hopfion. This distinguishes the dynamics of the hopfion from other previously studied TSTs in which the dynamical modes (e.g., longitudinal and transverse motion) are coupled via the gyrovector (total emergent magnetic flux of the spin texture) [1,6,54–58]. In principle, the velocity coupled with the toroidal moment can generate nonreciprocal effects [18,59], and thus the nonreciprocal dynamics of the hopfion is also expected from the coupled dynamical modes.

To carefully check the nonreciprocal dynamics of the hopfion, spin dynamics simulations were performed. A magnetic hopfion stabilized in a frustrated magnet is employed as the initial state. The corresponding model Hamiltonian is $\mathcal{H} = -\sum_{\langle i,j \rangle} J_{ij} \mathbf{S}_i \cdot \mathbf{S}_j$, where the summation of the exchange interaction is extended up to fourth nearest neighbor on a cubic mesh [47]. The presence of a hopfion spontaneously breaks the inversion symmetry of this system.

The spin dynamics are calculated by solving the Landau-Lifshitz-Gilbert (LLG) equation with the spin transfer torque (STT) terms [60–62]:

$$\begin{aligned} \frac{d\mathbf{S}}{dt} = & -\gamma_0 \mathbf{S} \times \mathbf{H}_{\text{eff}} + \frac{\alpha}{S} \mathbf{S} \times \frac{d\mathbf{S}}{dt} + \frac{pa^3}{2eS} (\mathbf{j} \cdot \nabla) \mathbf{S} \\ & + \frac{pa^3 \beta}{2eS^2} (\mathbf{j} \cdot \nabla) \mathbf{S} \times \mathbf{S}. \end{aligned} \quad (2)$$

Here γ_0 is the gyromagnetic ratio, α is the damping constant, p is the spin polarization, a is the lattice constant, and \mathbf{H}_{eff} is the effective magnetic field. S is the spin length, which is fixed to be 1 for simplicity. The effects of STT are described by the last two terms in Eq. (2), where \mathbf{j} is the current density and β quantifies the nonadiabaticity of the STT.

Typical results of the hopfion dynamics under an ac driving current $j_z = j \sin(\omega t)$ applied along the z axis are summarized in Fig. 2. As shown in the snapshots [Fig. 2(a)] within a single driving period, for the first (second) half of the period with positive (negative) applied current, the hopfion moves along the current direction with a clockwise (counterclockwise) rotation and an expansion (shrinkage). Interestingly, after each cycle, the hopfion gains a net translation and rotation. Snapshots after multiple cycles




FIG. 2. Nonreciprocal dynamics of a magnetic hopfion driven by a sinusoidal-type ac current applied along the z axis. (a) Snapshots of the hopfion dynamics within a single driving cycle. (b) Snapshots of the hopfion dynamics showing clear net translation and rotation after 10 and 20 driving cycles. (c) and (d) show the translation and rotation of the ac hopfion dynamics. Inset of (d) also shows the change of the hopfion's size over time.

with more pronounced net translation and rotation are shown in Fig. 2(b) for clarity (also see movie for the nonreciprocal dynamics in the Supplemental Material [47]). The steady-state ac dynamics plotted in Figs. 2(c) and 2(d) also show that a net translation and rotation are associated with the oscillating dynamics and they increase with time in a linear manner. Therefore, the hopfion possesses nonreciprocal translation and rotation under an ac driving current. On the other hand, the size of hopfion d oscillates around an equilibrium value [Fig. 2(d) inset] so that the steady-state ac dynamics can be sustained. The current density dependences of the hopfion's net velocity v and net angular velocity $\dot{\theta}$ are plotted in Figs. 3(a) and 3(c). The quadratic dependence on the current density suggests a nonlinear nonreciprocal dynamics of the hopfion.

To better understand this nonlinear nonreciprocal dynamics, we derive the corresponding equation of motion via the generalized Thiele's approach based on the dynamical modes of the hopfion [39,56,63]. Other than the translation, rotation, and dilation of the hopfion, it is necessary to take into account two additional effects in order to capture the nonreciprocal dynamics. First, there is always an energy change associated with the dilation of the hopfion. Although this energy change can be ignored in describing the linear dc dynamics, it is quite essential here




FIG. 3. Current density and $(\beta/\alpha - 1)$ dependence of the hopfion's net velocity [(a) and (b)] and net angular velocity [(c) and (d)]. Inset in (a) and (c) is the corresponding driving frequency dependence. Blue and red dots are simulation results. Black lines represent the j^2 fitting in (a) and (c), the $(1/\omega^2)$ fitting in the inset of (a) and (c), and the $(\beta/\alpha - 1)^2$ fitting in (b) and (d), respectively. The ac current is applied along the z axis.

for generating the nonlinear nonreciprocal dynamics. The formation of the hopfion requires a minimum number of spins and it assigns an energy limit bounded with the shrinking of hopfion. Hence, the energy of the hopfion is asymmetric with respect to the dilation (expansion and shrinkage) and can be effectively described by an anharmonic potential $E = \frac{1}{2}m(\lambda - 1)^2 + \frac{1}{6}\chi(\lambda - 1)^3$ (for more details, see Supplemental Material [47]). In addition, the dilation also leads to an effective field $\mathbf{H}_\lambda = (\zeta/\mu_s)\mathbf{S}_0 \times [(\lambda - 1)\mathbf{r} \cdot \partial_r]\mathbf{S}_0$ acted on the hopfion, where μ_s is the magnetic moment and ζ is a phenomenological parameter similar to m and γ . These two factors associated with the dilation of the hopfion are key ingredients for understanding the nonreciprocal dynamics of the hopfion beyond the linear response regime.

Based on these dynamical modes, the equations of motion for the hopfion are obtained as

$$-T_z^e \dot{\lambda} + \alpha K_{RR} \dot{Z} + \alpha K_{R\Theta} \dot{\theta}_z = \beta K_{RR} \xi j_z - \frac{\gamma_0 \zeta}{\mu_s} T_z^e (\lambda - 1), \quad (3)$$

$$\Gamma^e \dot{\lambda} + \alpha K_{R\Theta} \dot{Z} + \alpha K_{\Theta\Theta} \dot{\theta}_z = \beta K_{R\Theta} \xi j_z + \frac{\gamma_0 \zeta}{\mu_s} \Gamma^e (\lambda - 1), \quad (4)$$

$$T_z^e \dot{Z} - \Gamma^e \dot{\theta}_z + \alpha K_{\lambda\lambda} \dot{\lambda} = \xi T_z^e j_z - \frac{\gamma_0}{\mu_s} m (\lambda - 1) - \frac{\gamma_0}{2\mu_s} \chi (\lambda - 1)^2, \quad (5)$$

with $K_{RR} = \int (\partial_z \mathbf{S})^2 dV$, $K_{R\Theta} = \int \partial_z \mathbf{S} \cdot (x \partial_y - y \partial_x) \mathbf{S} dV$, $K_{\Theta\Theta} = \int [(x \partial_y - y \partial_x) \mathbf{S}]^2 dV$, and $K_{\lambda\lambda} = \int (\mathbf{r} \cdot \partial_r \mathbf{S})^2 dV$. By setting the current density $j_z = j \sin(\omega t)$, these equations can be solved in the ac limit hierarchically with respect to the order of j .

We then solve the dynamical modes as $\dot{Z} \approx \dot{Z}_1(j) + \dot{Z}_2(j^2)$, $\dot{\theta}_z \approx \dot{\theta}_{z1}(j) + \dot{\theta}_{z2}(j^2)$, and $\lambda \approx \lambda_0 + \lambda_1(j) + \lambda_2(j^2)$ up to the second order of j (for details of the solution, see Supplemental Material [47]). While the first order solutions (with j linear terms) only contain oscillating dynamics, their second order solutions (with j^2 terms) include non-oscillating terms, i.e., the terms related to the nonreciprocal dynamics. By further imposing the symmetry of the hopfion, the following simplified dependences can be obtained [47]

$$\dot{Z}_2^{\text{dc}} \sim \frac{1}{\omega^2} T_z^e \left(\frac{\beta}{\alpha} - 1 \right)^2 j^2, \quad (6)$$

$$\dot{\theta}_{z2}^{\text{dc}} \sim -\frac{1}{\omega^2} \Gamma^e \left(\frac{\beta}{\alpha} - 1 \right)^2 j^2. \quad (7)$$

\dot{Z}_2^{dc} and $\dot{\theta}_{z2}^{\text{dc}}$ represent the nonlinear nonreciprocal translation and rotation of the hopfion. They have the same dependence on the current density j , driving frequency ω , and $[(\beta/\alpha) - 1]$. These dependences are also consistent with the simulation results as shown in Fig. 3. The anomaly in the frequency dependence might be due to some resonances or other internal modes of the hopfion, which are difficult to capture using the current formalism.

Next, we discuss the nonlinear hopfion responses under an ac current applied perpendicular to the direction of T_z^e (e.g., along the x axis). A previous study suggests the dc hopfion dynamics under an in-plane current include a rotation of the hopfion plane, which is equivalent to a rotation of T_z^e [39]. Therefore, although the current and T_z^e are initially perpendicular to each other, as time increases, the current component along T_z^e develops and nonlinear dynamics of the hopfion are still expected.

The simulation results for an ac current applied along the x axis are shown in Fig. 4. While the hopfion oscillates along the x axis, the net translation and rotation are still bounded with the z axis. Hence, the hopfion moves perpendicular to the applied ac current direction, which is a nonlinear hopfion Hall effect. Moreover, the current density dependence, frequency dependence, and (β/α) dependence of the hopfion's net velocity and angular velocity have also been checked, which all agree well with the analytical results (see Supplemental Materials [47]).

The equation of motion [Eq. (6) and Eq. (7)] indicates that T_z^e and Γ^e characterize the nonlinear dynamics of hopfion. The nonreciprocities of \dot{Z}_2^{dc} and $\dot{\theta}_{z2}^{\text{dc}}$ are determined by T_z^e and Γ^e , respectively. Therefore, it is possible to control the nonlinear dynamics of hopfion by manipulating its spin configuration (e.g., switch the chirality or polarity). Although both the translation and rotation of the hopfion show nonlinear responses, the corresponding symmetries are not the same. T_z^e breaks both \mathcal{T} and \mathcal{I} . As a result, the velocity of the hopfion reverses its sign upon \mathcal{T} and \mathcal{I} operation but is invariant under \mathcal{IT} operation [Fig. 5(a)].




FIG. 4. Nonlinear hopfion Hall effect under an ac current applied along the x axis. (a) The translation of ac hopfion dynamics along different axes. (b) The ac rotation dynamics of the hopfion around the z axis. Inset of (b) shows the schematic of the nonlinear hopfion Hall effect.

On the other hand, the rotation of the hopfion is determined by Γ^e , a parity-even emergent magnetomultipole. So the net rotation of the hopfion reverses its sign upon \mathcal{T} or \mathcal{IT} only [Fig. 5(b)].

Finally, to facilitate a more complete understanding, we study the nonlinear electron transport effects induced by a hopfion, which is effectively the counterpart of the STT effect. The second-order current density in response to the external electric field \mathbf{E} is defined as

$$j_{2,a} = \chi_{abc} E_b E_c \quad (8)$$

where χ_{abc} is the second-order conductivity tensor. By considering a double exchange model (in which the itinerant electrons are coupled to the localized spins via the exchange coupling) [64,65], χ_{abc} can be obtained by solving the semiclassical Boltzmann equation [66–68] (for details of the derivation, see Ref. [47]). For the hopfion spin texture studied here ($\mathbf{T}^e = T_z^e \hat{\mathbf{z}}$), three typical second-order conductivity components $\chi_{zzz}/2 = \chi_{zxx} = \chi_{zyy} = \chi \propto T_z^e$ are obtained (detailed solutions can be found in [47]). So the emergent toroidal moment of the hopfion induces two types of nonlinear responses. χ_{zxx} and χ_{zyy} characterize a nonlinear Hall effect and χ_{zzz} characterizes a nonreciprocal transport, both along the direction of T_z^e . These nonlinear transport effects are consistent with the nonlinear spin dynamics of the hopfion.

Our results reveal the vital role of the emergent magnetomultipoles in determining the nonlinear spin dynamics and




FIG. 5. Nonreciprocal translation (a) and rotation (b) for different symmetry operations on the hopfion configuration.

nonlinear electrical transports related to a magnetic hopfion. We anticipate the hopfions in other physical systems may also show nonlinear responses in a similar manner as the vector field itself breaks at least the inversion symmetry [69–72]. As an example, we demonstrate the nonlinear nonreciprocal dynamics of a hopfion stabilized in a typical chiral magnet with conical background (see Supplemental Materials [47]). In addition to hopfion, the emergent magnetomultipoles can also be applied to other TSTs in both two- and three-dimensional systems and can be engineered by tailoring different kinds of TSTs. This Letter also provides an alternative working principle for the potential implementations of a hopfion in 3D spintronic devices [25,28].

We thank W. Koshibae for fruitful discussions. N. N. was supported by JST CREST Grants No. JMPJCR1874 and No. JPMJCR16F1, and JSPS KAKENHI Grant No. 18H03676. H. W. acknowledges support from JSPS KAKENHI No. JP21J00453. Y. L. acknowledges financial support from the RIKEN Special Postdoctoral Researcher (SPDR) Program and JST CREST Grant No. JPMJCR20T1.

-
- [1] N. Nagaosa and Y. Tokura, *Nat. Nanotechnol.* **8**, 899 (2013).
 [2] A. Fert, N. Reyren, and V. Cros, *Nat. Rev. Mater.* **2**, 17031 (2017).
 [3] B. Binz and A. Vishwanath, *Physica (Amsterdam)* **403B**, 1336 (2008).
 [4] A. Neubauer, C. Pfleiderer, B. Binz, A. Rosch, R. Ritz, P. G. Niklowitz, and P. Böni, *Phys. Rev. Lett.* **102**, 186602 (2009).
 [5] N. Nagaosa, J. Sinova, S. Onoda, A. H. MacDonald, and N. P. Ong, *Rev. Mod. Phys.* **82**, 1539 (2010).
 [6] J. Zang, M. Mostovoy, J. H. Han, and N. Nagaosa, *Phys. Rev. Lett.* **107**, 136804 (2011).
 [7] W. Jiang, X. Zhang, G. Yu, W. Zhang, X. Wang, M. Benjamin Jungfleisch, J. E. Pearson, X. Cheng, O. Heinonen, K. L. Wang, Y. Zhou, A. Hoffmann, and S. G. E. te Velthuis, *Nat. Phys.* **13**, 162 (2017).
 [8] K. Litzius, I. Lemesch, B. Krüger, P. Bassirian, L. Caretta, K. Richter, F. Büttner, K. Sato, O. A. Tretiakov, J. Förster, R. M. Reeve, M. Weigand, I. Bykova, H. Stoll, G. Schütz, G. S. D. Beach, and M. Kläui, *Nat. Phys.* **13**, 170 (2017).
 [9] N. Nagaosa, *Proc. Jpn. Acad. Ser. B* **95**, 278 (2019).
 [10] T. Yokouchi, F. Kagawa, M. Hirschberger, Y. Otani, N. Nagaosa, and Y. Tokura, *Nature (London)* **586**, 232 (2020).
 [11] L. D. Landau and E. M. Lifshitz, *The Classical Theory of Fields: 4th Edition*, Graduate Texts in Contemporary Physics (Butterworth-Heinemann, London, 1975).
 [12] K. S. Thorne, *Rev. Mod. Phys.* **52**, 299 (1980).
 [13] V. M. Dubovik and V. V. Tugushev, *Phys. Rep.* **187**, 145 (1990).
 [14] N. A. Spaldin, M. Fiebig, and M. Mostovoy, *J. Phys. Condens. Matter* **20**, 434203 (2008).
 [15] Y. Tokura and N. Nagaosa, *Nat. Commun.* **9**, 3740 (2018).
 [16] T.-h. Arima, J.-H. Jung, M. Matsubara, M. Kubota, J.-P. He, Y. Kaneko, and Y. Tokura, *J. Phys. Soc. Jpn.* **74**, 1419 (2005).
 [17] J. Matsuno, T. Lottermoser, T. Arima, M. Kawasaki, and Y. Tokura, *Phys. Rev. B* **75**, 180403(R) (2007).
 [18] S.-W. Cheong, D. Talbayev, V. Kiryukhin, and A. Saxena, *npj Quantum Mater.* **3**, 19 (2018).
 [19] S. Bhowal and N. A. Spaldin, *Phys. Rev. Lett.* **128**, 227204 (2022).
 [20] I. Sodemann and L. Fu, *Phys. Rev. Lett.* **115**, 216806 (2015).
 [21] Q. Ma, S.-Y. Xu, H. Shen, D. MacNeill, V. Fatemi, T.-R. Chang, A. M. Mier Valdivia, S. Wu, Z. Du, C.-H. Hsu, S. Fang, Q. D. Gibson, K. Watanabe, T. Taniguchi, R. J. Cava, E. Kaxiras, H.-Z. Lu, H. Lin, L. Fu, N. Gedik, and P. Jarillo-Herrero, *Nature (London)* **565**, 337 (2019).
 [22] C.-P. Zhang, X.-J. Gao, Y.-M. Xie, H. C. Po, and K. T. Law, *arXiv:2012.15628*.
 [23] Z. Z. Du, H.-Z. Lu, and X. C. Xie, *Nat. Rev. Phys.* **3**, 744 (2021).
 [24] L. Faddeev and A. J. Niemi, *Nature (London)* **387**, 58 (1997).
 [25] A. Fernández-Pacheco, R. Streubel, O. Fruchart, R. Hertel, P. Fischer, and R. P. Cowburn, *Nat. Commun.* **8**, 15756 (2017).
 [26] C. Donnelly, M. Guizar-Sicairos, V. Scagnoli, S. Gliga, M. Holler, J. Raabe, and L. J. Heyderman, *Nature (London)* **547**, 328 (2017).
 [27] C. Donnelly, K. L. Metlov, V. Scagnoli, M. Guizar-Sicairos, M. Holler, N. S. Bingham, J. Raabe, L. J. Heyderman, N. R. Cooper, and S. Gliga, *Nat. Phys.* **17**, 316 (2021).
 [28] S.-H. Yang, R. Naaman, Y. Paltiel, and S. S. P. Parkin, *Nat. Rev. Phys.* **3**, 328 (2021).
 [29] P. Sutcliffe, *Phys. Rev. Lett.* **118**, 247203 (2017).
 [30] Y. Liu, R. K. Lake, and J. Zang, *Phys. Rev. B* **98**, 174437 (2018).
 [31] P. Sutcliffe, *J. Phys. A* **51**, 375401 (2018).
 [32] J.-S. B. Tai and I. I. Smalyukh, *Phys. Rev. Lett.* **121**, 187201 (2018).
 [33] F. N. Rybakov, N. S. Kiselev, A. B. Borisov, L. Döring, C. Melcher, and S. Blügel, *arXiv:1904.00250*.
 [34] R. Voinescu, J.-S. B. Tai, and I. I. Smalyukh, *Phys. Rev. Lett.* **125**, 057201 (2020).
 [35] N. Kent, N. Reynolds, D. Raftrey, I. T. G. Campbell, S. Virasawmy, S. Dhuey, R. V. Chopdekar, A. Hierro-Rodríguez, A. Sorrentino, E. Pereiro, S. Ferrer, F. Hellman, P. Sutcliffe, and P. Fischer, *Nat. Commun.* **12**, 1562 (2021).
 [36] J. H. C. Whitehead, *Proc. Natl. Acad. Sci. U.S.A.* **33**, 117 (1947).
 [37] F. Wilczek and A. Zee, *Phys. Rev. Lett.* **51**, 2250 (1983).
 [38] N. Manton and P. Sutcliffe, *Topological Solitons* (Cambridge University Press, Cambridge, England, 2004).
 [39] Y. Liu, W. Hou, X. Han, and J. Zang, *Phys. Rev. Lett.* **124**, 127204 (2020).
 [40] X. S. Wang, A. Qaiumzadeh, and A. Brataas, *Phys. Rev. Lett.* **123**, 147203 (2019).
 [41] D. Raftrey and P. Fischer, *Phys. Rev. Lett.* **127**, 257201 (2021).

- [42] L. Bo, L. Ji, C. Hu, R. Zhao, Y. Li, J. Zhang, and X. Zhang, *Appl. Phys. Lett.* **119**, 212408 (2021).
- [43] Z. Khodzhaev and E. Turgut, *J. Phys. Condens. Matter* **34**, 225805 (2022).
- [44] S. Li, J. Xia, L. Shen, X. Zhang, M. Ezawa, and Y. Zhou, *Phys. Rev. B* **105**, 174407 (2022).
- [45] B. Göbel, C. A. Akosa, G. Tatara, and I. Mertig, *Phys. Rev. Res.* **2**, 013315 (2020).
- [46] S. S. Pershoguba, D. Andreoli, and J. Zang, *Phys. Rev. B* **104**, 075102 (2021).
- [47] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.129.267201> for details of the derivations, numerical simulations, nonreciprocal dynamics of hopfion in chiral magnets, and movie of the nonreciprocal dynamics, which includes Refs. [48] and [49].
- [48] G. P. Müller, M. Hoffmann, C. Dißelkamp, D. Schürhoff, S. Mavros, M. Sallermann, N. S. Kiselev, H. Jónsson, and S. Blügel, *Phys. Rev. B* **99**, 224414 (2019).
- [49] A. Vansteenkiste, J. Leliaert, M. Dvornik, M. Helsen, F. Garcia-Sanchez, and B. Van Waeyenberge, *AIP Adv.* **4**, 107133 (2014).
- [50] A. Auerbach, *Interacting Electrons and Quantum Magnetism*, Graduate Texts in Contemporary Physics (Springer-Verlag, New York, 1994).
- [51] G. Tatara, H. Kohno, and J. Shibata, *Phys. Rep.* **468**, 213 (2008).
- [52] H. Watanabe and Y. Yanase, *Phys. Rev. B* **98**, 245129 (2018).
- [53] S. Hayami, M. Yatsushiro, Y. Yanagi, and H. Kusunose, *Phys. Rev. B* **98**, 165110 (2018).
- [54] N. Papanicolaou and T. N. Tomaras, *Nucl. Phys.* **B360**, 425 (1991).
- [55] K. Y. Guslienko, *J. Nanosci. Nanotechnol.* **8**, 2745 (2008).
- [56] O. A. Tretiakov, D. Clarke, G.-W. Chern, Y. B. Bazaliy, and O. Tchernyshyov, *Phys. Rev. Lett.* **100**, 127204 (2008).
- [57] K. Everschor-Sitte and M. Sitte, *J. Appl. Phys.* **115**, 172602 (2014).
- [58] C. Psaroudaki and D. Loss, *Phys. Rev. Lett.* **120**, 237203 (2018).
- [59] S.-W. Cheong and X. Xu, *npj Quantum Mater.* **7**, 40 (2022).
- [60] J. C. Slonczewski, *J. Magn. Magn. Mater.* **159**, L1 (1996).
- [61] L. Berger, *Phys. Rev. B* **54**, 9353 (1996).
- [62] S. Zhang and Z. Li, *Phys. Rev. Lett.* **93**, 127204 (2004).
- [63] A. A. Thiele, *Phys. Rev. Lett.* **30**, 230 (1973).
- [64] J. Ye, Y. B. Kim, A. J. Millis, B. I. Shraiman, P. Majumdar, and Z. Tešanović, *Phys. Rev. Lett.* **83**, 3737 (1999).
- [65] P. Bruno, V. K. Dugaev, and M. Taillefumier, *Phys. Rev. Lett.* **93**, 096806 (2004).
- [66] H. Isobe, S.-Y. Xu, and L. Fu, *Sci. Adv.* **6**, eaay2497 (2020).
- [67] H. Ishizuka and N. Nagaosa, *Nat. Commun.* **11**, 2986 (2020).
- [68] H. Isobe and N. Nagaosa, *J. Phys. Soc. Jpn.* **91**, 115001 (2022).
- [69] G. Volovik and V. Mineev, *Sov. Phys. JETP* **46**, 401 (1977), <http://jetp.ras.ru/cgi-bin/e/index/e/46/2/p401?a=list>.
- [70] E. Babaev, *Phys. Rev. Lett.* **88**, 177002 (2002).
- [71] Y. Kawaguchi, M. Nitta, and M. Ueda, *Phys. Rev. Lett.* **100**, 180403 (2008).
- [72] P. J. Ackerman and I. I. Smalyukh, *Nat. Mater.* **16**, 426 (2017).