

Photon Ring Astrometry for Superradiant Clouds

Yifan Chen^{1,*}, Xiao Xue,^{2,3,†} Richard Brito^{4,‡} and Vitor Cardoso^{1,4,§}

¹*Niels Bohr International Academy, Niels Bohr Institute, Blegdamsvej 17, 2100 Copenhagen, Denmark*

²*II. Institute of Theoretical Physics, Universität Hamburg, 22761 Hamburg, Germany*

³*Deutsches Elektronen-Synchrotron DESY, Notkestr. 85, 22607 Hamburg, Germany*

⁴*CENTRA, Departamento de Física, Instituto Superior Técnico—IST, Universidade de Lisboa—UL, Avenida Rovisco Pais 1, 1049 Lisboa, Portugal*



(Received 16 November 2022; revised 25 January 2023; accepted 16 February 2023; published 14 March 2023)

Gravitational atoms produced from the superradiant extraction of rotational energy of spinning black holes can reach energy densities significantly higher than that of dark matter, turning black holes into powerful potential detectors for ultralight bosons. These structures are formed by coherently oscillating bosons, which induce oscillating metric perturbations deflecting photon geodesics passing through their interior. The deviation of nearby geodesics can be further amplified near critical bound photon orbits. We discuss the prospect of detecting this deflection using photon ring autocorrelations with the Event Horizon Telescope and its next-generation upgrade, which can probe a large unexplored region of the cloud mass parameter space when compared with previous constraints.

DOI: 10.1103/PhysRevLett.130.111401

Introduction.—Ultralight bosons, such as axions, dark photons, or massive tensors, are well-motivated particles beyond the standard model. They are predictions of fundamental theories with extra dimensions [1–4], and are excellent dark matter candidates [5–9] with masses above $\sim 10^{-22}$ eV. The lightness of such fields enhances wavelike properties on astrophysical scales making their phenomenology extremely rich [10].

An important example of this rich phenomenology occurs when the boson Compton wavelength is comparable to the gravitational radius of a rotating black hole (BH). Then, a dense bound state can be formed via superradiant extraction of the BH rotational energy [11–13]. Through this process, a “boson cloud” can form outside the horizon, giving rise to a system also known as a gravitational atom [3,14–16]. Assuming that the BH has no external supplement of angular momentum, and considering a single unstable bound state, up to $\sim \mathcal{O}(10)\%$ of the BH mass can be transferred to the cloud [17–19]. The local energy density of this cloud can be orders of magnitude higher than the one of virialized dark matter. Thus, the environment outside rotating BHs can be used as a powerful detector for ultralight bosons [13]; this can be achieved via BH mass and spin measurements [17,20–29], gravitational-wave emission [20–23,26,30–42], detection of binary BH systems [43–49], axion-induced birefringence [50–53], shadow evolution [54–57], and lensing due to the extended energy distribution [58–61]. Some of these signatures benefit from the unprecedented spatial resolution achieved by the Event Horizon Telescope (EHT) [62–70] and are expected to improve substantially with its next generation upgrade (ngEHT) [71–74].

In this Letter, we introduce a new method to search for boson clouds. We propose to use the trapping properties of BHs, which are tightly connected with the existence of unstable, bound null geodesics. The existence of such geodesics is of central importance for the interpretation of BH images [73,75–88]. For near-critical orbits, i.e., null geodesics arbitrarily close to the bound photon orbits, the motion is unbound, and photons can propagate to asymptotic observers while probing the near-horizon region. Photons from the near-critical motion circle the BH a number of times and can be used as an astrometry tool to detect oscillating lensing effects induced by real ultralight bosons.

We work in units where $\hbar = c = 1$ and adopt the metric convention $(-, +, +, +)$. Greek indices take values $(0, 1, 2, 3)$, while spatial indices are denoted by latin letters, i.e., $x^\mu = (x^0, x^i)$. For clarity, we use subscripts or superscripts V, T when referring to quantities related to a massive vector or tensor field, respectively.

Oscillatory deflections from boson clouds.—Consider a boson field of mass μ , a BH of mass M_{BH} , and a dimensionless angular momentum a_J pointing along the z axis. When characterizing the BH-boson state, it is useful to employ a Boyer-Lindquist coordinate system. Ultralight bosons can bind with the BH to form a hydrogenlike gravitational atom with discrete quantum numbers [13,14,89] and a gravitational fine-structure constant $\alpha \equiv G_N M_{\text{BH}} \mu$, where G_N is Newton’s gravitational constant. The eigenstates of the boson are characterized by an energy $\omega \sim \mu$ [13,14]. When the angular velocity Ω of the BH is larger than the angular phase velocity of the bosons, i.e., $\omega/m < \Omega$, where m is the azimuthal number of the

boson, then a bound superradiant state is possible, which grows via extraction of rotational energy from the BH [12,13].

Their energy-momentum tensor $T_{\mu\nu}$ induces both static and oscillating metric perturbations in the metric. For vector fields, only the traceless part of T_{ij}^V oscillates. Therefore, neglecting subdominant spatial derivatives, one finds that the oscillating part of the metric perturbation is given by $H_{ij}^V \approx -T_{ij}^{\text{osc}}/(2\mu^2 m_{\text{pl}}^2)$, where $m_{\text{pl}} \equiv 1/\sqrt{8\pi G_N}$ is the reduced Planck mass. On the other hand, massive tensors behave as a localized effective strain $H_{\mu\nu}^T$ proportional to its field value and a dimensionless coupling constant α_h characterizing the strength of the interaction between the massive tensor and photons [90,91]. We introduce dimensionless parameters $\epsilon_V \equiv (\Psi_0^V/m_{\text{pl}})^2/2$ and $\epsilon_T \equiv \alpha_h \Psi_0^T/m_{\text{pl}}$ to characterize the perturbative strain amplitude generated from a vector and a tensor cloud, respectively, where Ψ_0^V and Ψ_0^T are their corresponding maximal field value. See the Supplemental Material [98] for details.

Considering metric perturbations $H_{\mu\nu} \equiv \epsilon h_{\mu\nu}$ expanded around a Kerr background $g_{\mu\nu}$, we have $g_{\mu\nu} \approx g_{\mu\nu}^K + \epsilon h_{\mu\nu}$, where $\epsilon \ll 1$ controls the amplitude of the perturbations. The photon geodesics in this perturbed metric can be similarly expanded, $x^\mu \approx x_{(0)}^\mu + \epsilon x_{(1)}^\mu$, where $x_{(1)}^\mu$ satisfies

$$\frac{d^2 x_{(1)}^\mu}{d\lambda^2} = -2\Gamma_{\alpha\beta}^{K\mu} \frac{dx_{(0)}^\alpha}{d\lambda} \frac{dx_{(1)}^\beta}{d\lambda} - \Gamma_{\alpha\beta}^{(1)\mu} \frac{dx_{(0)}^\alpha}{d\lambda} \frac{dx_{(0)}^\beta}{d\lambda}. \quad (1)$$

Here, $x_{(0)}^\mu$ and $\Gamma_{\alpha\beta}^{K\mu}$ are the unperturbed null geodesics and the connection of the Kerr background, respectively, and λ is an affine parameter. $\Gamma_{\alpha\beta}^{(1)\mu}$ is the perturbed connection generated from bosonic clouds.

We use a backward ray tracing code on a Kerr background, KGEO [108], to compute $x_{(0)}^\mu$ from a faraway camera toward the BH employing the integral method [109–112]. We consider an faraway observer at an inclination angle $\theta_o = 17^\circ$ from the black hole with dimensionless spin $a_J = 0.94$, consistent with the EHT observations of M87* [62]. We calculate the geodesics starting at different pixels on the observer plane and keep only those that hit the BH equatorial plane at least two times. The initial ($\lambda = 0$) components of photon geodesics are fixed to be $r(\lambda = 0) = 10^3 r_g$, $t(\lambda = 0) = 0$, without loss of generality, where $r_g \equiv G_N M_{\text{BH}}$ is the gravitational radius. The plasma medium effects, such as absorption and scattering effects, are neglected since the accretion flow is considered to be optically thin.

An example of a near-critical unperturbed geodesic $x_{(0)}^\mu$ is shown in the top panel of Fig. 1 using a white line that, due to strong lensing effects, crosses the gray equatorial plane of the Kerr BH several times. We also show samples of the perturbed spatial photon geodesics $x_{(0)}^\mu + \epsilon_T x_{(1)}^\mu$ for




FIG. 1. Examples of photon geodesics connecting a faraway observer from a Kerr BH surrounded by a bosonic cloud, that execute multiple orbits intersecting the equatorial plane (gray plane in top panel). Top panel: white line shows the unperturbed geodesic, while different colors show perturbed geodesics $x_{(0)}^\mu + \epsilon_T x_{(1)}^\mu$ at different oscillation phases of bosons. Bottom panel: deviation of the geodesics in Kerr-Schild coordinates in terms of the affine parameter λ . The dashed vertical line corresponds to the point where the unperturbed orbit $x_{(0)}^\mu$ first crosses the equatorial plane.

eight evenly spaced initial phases to account for the full oscillation behavior of the cloud. As expected, deviations start to be apparent after entering the photon ring orbit.

The deviation $x_{(1)}^\mu$ is shown in the bottom panel of Fig. 1 for a massive tensor and vector cloud in the ground state with $\alpha = 0.2$. To avoid (unphysical) singularities at the poles, we show $x_{(1)}^\mu$ using Cartesian Kerr-Schild coordinates (t, x, y, z) , in units of the gravitational radius r_g . The evolution of $x_{(1)}^\mu$ as a function of the affine parameter can be divided in two stages. The first stage (bottom-left panel) is dominated by perturbative deviations from cloud-induced metric fluctuations far away from the BH. In the second stage, when the orbit is close to the photon ring, deviations are exponentially amplified, as seen on the bottom right panel of Fig. 1.

The second term on the right-hand side of Eq. (1) dominates during the first stage, which is especially important for photons propagating at the typical radial scale of the cloud, i.e., r_g/α^2 , where our Newtonian approximation for the cloud profile, as discussed in the Supplemental Material [98], is reasonable. The oscillation of the boson fields leave imprints on $x_{(1)}^\mu$, with the

frequency of the oscillations for vector clouds being twice the one of the tensor cloud case. The exponential growth of $x_{(1)}^\mu$ starts when $x_{(0)}^\mu$ enters a nearly bound orbit where the first term in Eq. (1) becomes important. This growth is caused by the instability of the photon ring orbit [75–82,87,111]. More precisely, a face-on observer would see a deviation increase by a factor of ~ 20 between two sequential crossings of the BH equatorial plane.

Astrometry with photon ring autocorrelations.—The next question concerns the observability of such an effect. One sensitive and realistic probe is to use photon ring autocorrelations as proposed in Ref. [88], which is an especially useful observable for BHs observed nearly face on, such as M87* and SgrA*.

The strong gravity region outside Kerr BHs is responsible for the existence of bound null geodesics and for near-critical trajectories where photons propagate around the BH multiple times before reaching a narrow ring region on the observer plane. These strongly lensed photons form the photon ring with a locally enhanced intensity [85–87]. One way to observe the strongly lensed photons, without the need of significant improvements of the current spatial resolution, is to exploit the time variability of the emission from the accretion flow, for example. Indeed, intensity fluctuations due to the flow's turbulent nature were observed by the EHT [70,113]. For each emission point, there are multiple geodesics connecting it to the observer's image plane that differ by the number k of half orbits around the BH. In the image plane, we can use polar coordinates (ρ, φ) , where ρ is the radial distance from the BH and φ is the polar angle. Photons with common origin but different k reach the photon ring separated in time t and angle φ , rendering the two-point intensity fluctuation correlation [88]

$$\mathcal{C}(T, \Phi) \equiv \langle \Delta I(t, \varphi) \Delta I(t + T, \varphi + \Phi) \rangle \quad (2)$$

to have peaks at $T \approx N\tau_0$, $\Phi \approx N\delta_0$ for a background Kerr BH, where ΔI is the intensity fluctuation after an integration in the radial direction on the photon ring, $N \equiv k - k'$ is the difference between a photon pair executing k and k' half orbits, and $\{\tau_0, \delta_0\}$ are the critical parameters characterizing the time delay and azimuthal lapse of the bound photon orbits, respectively [111]. This is a universal prediction dependent only on the space-time near the photon shell, especially for optically and geometrically thin emission flows [114].

In the presence of boson clouds, small oscillations around the unperturbed geodesics grow exponentially close to the photon ring orbit, leading to a periodic shift in the peak positions, $T \approx N\tau_0 + \Delta T^N$ and $\Phi \approx N\delta_0 + \Delta\Phi^N$. For photons emitted from a geometrically thin disk at the equatorial plane, the deviations can be calculated using

$$\begin{aligned} \Delta T^N/\epsilon &= x_{(1)}^t(\lambda_N) - x_{(1)}^t(\lambda_0), \\ \Delta\Phi^N/\epsilon &= x_{(1)}^\phi(\lambda_N) - x_{(1)}^\phi(\lambda_0), \end{aligned} \quad (3)$$

where λ_N represents the affine parameter at which the perturbed geodesics crosses the equatorial plane, i.e., $z_{(0)} + \epsilon z_{(1)} = 0$, for the $N+1$ times. Since $x_{(1)}^\mu$ grows exponentially after λ_0 , one can safely neglect $x_{(1)}^\mu(\lambda_0)$ in Eq. (3). Notice that one can get an equivalent gauge-invariant description of the geodesics deflection and shifts on the image plane using the deviation of the conserved quantities in the Kerr space-time instead.

The fundamental resolution of the autocorrelation is limited by the intrinsic correlation length of the source, which can be inferred from general relativistic magnetohydrodynamic simulations [88]. Taking $\ell_\phi \approx 4.3^\circ$ and $\ell_t \approx 3r_g$ as correlation lengths for ϕ and t , respectively [88], $\Delta T^N/\ell_t$ is typically smaller than $\Delta\Phi^N/\ell_\phi$. We thus focus on $\Delta\Phi^N$, fitting it to be $\Delta\Phi^N = \Phi_0^N \cos(\omega t + \delta)$ for $N = 1$ and 2, where Φ_0^N is the amplitude and δ is the relative oscillation phase. The phase δ is always well fit by the sum of 2φ and a small deviation, representing an $m = 2$ mode of metric perturbation and a small inclination angle.

The spatially varying Φ_0^N , as shown in the Supplemental Material [98], would require weighting the intensity fluctuation spectrum as a function of the photon ring region, which is beyond the scope of this study. We will instead estimate the detectability of the signal using a specific point on the critical curve with the largest amplitude $\varphi \approx 0.7\pi$, which is also the point with the most significant intensity contributed by the lensed photon [57]. We show Φ_0^N at this point as a function of α for the ground-state tensor and vector cloud in Fig. 2. As expected, Φ_0^{N2} is typically larger due to the exponential sequential growth. For completeness, we also consider the higher vector field mode (2122) (this notation is explained in the Supplemental Material [98]), which has a smaller maximum value for Φ_0^N when compared to the ground-state mode due to the $\sin^2\theta$ suppression for geodesics reaching a nearly face-on observer.




FIG. 2. Oscillation amplitude Φ_0 of the azimuthal lapse at $\varphi \approx 0.7\pi$ on the critical curve at different values of α with $a_J = 0.94$, $\theta_o = 17^\circ$, and $\epsilon = 10^{-3}$.

Prospective constraints.—In a realistic setup, the finite spatial resolution of EHT or ngEHT leads to a Gaussian smearing of the correlation peak in the (T, Φ) plane. For ngEHT, assuming an $\sim 10\mu$ as spatial resolution [120,121], the 1σ width of the Gaussian packet in the Φ direction is around 10° for both M87* and SgrA*. On the other hand, in the presence of a boson cloud, there will be periodic oscillations of the central position of the Gaussian packet, causing a broadening of its width that becomes especially evident when the oscillation amplitude Φ_0^N is comparable or larger than the “original” width without oscillations. Thus, if a correlation peak predicted assuming a vacuum Kerr BH is observed, one can impose the conservative limit that $2\Phi_0^N < 10^\circ$. Notice that such a conservative estimate does not exploit the full oscillatory features. Once a detection of a correlation peak has been made, one can also check whether an oscillation in the center of the Gaussian packet provides a better description than a stationary peak, which can potentially give a sensitivity beyond the one limited by the Gaussian smearing. With less than 1 order-of-magnitude-longer observation times or a slightly improved spatial resolution, one could, in principle, resolve the intrinsic correlation of the plasma with $\ell_\phi \approx 4.3^\circ$. We thus use this limit as an optimistic criterion for our prospective constraints.

The oscillation amplitude Φ_0^N shown in Fig. 2 is proportional to ϵ_T or ϵ_V , which in turn is related to the total mass in the cloud as discussed in the Supplemental Material [98]. The results are summarized in Fig. 3, where the dashed lines show the optimistic limit based on the intrinsic correlation length, while the bands show conservative constraints that could be obtained immediately after a detection of the corresponding autocorrelation peak. Here we take $\alpha_h = 1$ for massive tensors and limit the α range for the vector mode (1011) to be below 0.5, which is the limit imposed by assuming that the ground-state cloud formed due to superradiance.

For comparison, we also show constraints taken from Ref. [61] based on joining EHT observations of M87*’s photon ring size with measurements of the BH mass-to-distance ratio obtained from stellar (or gas) dynamics. Those constraints correspond to the red region labeled “star motion + ring size.” Furthermore, there is a theoretical bound on the maximum mass a cloud can reach from extracting BH rotation energy through superradiance [19], assuming that there is no angular momentum supplement to the BH, which could be breached due to a potential accretion process [17,122].

The photon ring autocorrelations with the $N = 1$ subring could already go far beyond previous limits for a tensor cloud, while a detection of the $N = 2$ subring can probe both vector and tensor clouds sensitively. In the Supplemental Material, we show similar results for different BH spins [98]. In general, we find qualitatively similar results for smaller spins, but with smaller oscillations due to




FIG. 3. Prospects for constraints on the mass of a vector cloud or a tensor cloud with $\alpha_h = 1$ and a BH dimensionless spin $a_J = 0.94$. The red region comes from constraints from stellar motion jointly with ring size measurements using the current EHT with 10% error bars [61], and the green region is the theoretical bound on the maximum superradiant extraction for the considered modes [19]. The constraint bands show limits ranging from a conservative criterium based on ngEHT’s spatial resolution to an optimistic criterion based on the intrinsic azimuthal correlation length of the accretion flow.

the fact that the photon ring radius gets larger for smaller spins. However, even for $a_J = 0.5$, the $N = 2$ subring can always constrain interesting regions of the parameter space.

Discussion.—Metric perturbations induced by superradiant clouds lead to oscillatory deflections of photon geodesics. Because of strong gravitational effects, searching for this effect benefits from two aspects compared to previous astrometry searches for ultralight bosonic dark matter [90,95,96,123–129]: (i) Superradiant clouds can reach large field values outside a BH, and (ii) any deviation of the vacuum geodesics grows exponentially close to the photon ring orbit. Importantly, this method relies solely on observations of the photon ring. With longer integration times, the EHT could already detect the $N = 1$ subring [88], while the expected improvements in baseline coverage, spatial resolution, dynamic ranges, and multifrequency observations of the ngEHT could help shorten the integration time.

Another observable that we did not consider is the time delay that can be seen in Fig. 1, which is also the dominant observation channel for a scalar cloud. The prospects to detect such a time delay are less interesting due to the limited time resolution of the photon ring autocorrelation. However, in the presence of an emission source with shorter time correlation lengths, such as a hot spot [130–135] or a nearby pulsar [136–140], echoes from those sources could be used as astrometry with a dramatically enhanced time resolution similar to pulsar timing array searches.

Our discussion can also be applied to other profiles of ultralight bosons, such as a soliton core dark matter, whose typical field value is $\Psi_{\text{DM}} \approx 1.4 \times 10^{11} \text{ GeV}$ ($\epsilon_V \sim 10^{-15}$ and $\epsilon_T \sim 10^{-7} \alpha_h$) and radius $r_c \simeq 40 \text{ pc}$ for $\mu = 10^{-21} \text{ eV}$ [141]. Using a flat background approximation, the propagation of light in the soliton core leads to a spatial deviation $\sim \epsilon r_c \simeq 10 \alpha_h r_g$ before entering the photon ring orbit of SgrA*. Considering the amplification process in the photon ring, we estimate to be able to constrain α_h to be less than 10^{-4} using the $N = 1$ subring and an order-of-magnitude better for $N = 2$. These bounds are comparable with those obtained with pulsar timing arrays or planetary motion [90].

Finally, we should note that the oscillating cloud will also cause emission of gravitational waves (GWs). Assuming M87* and α that maximizes the constraints in Fig. 3, the half-life [142] of the cloud due to GW emission will typically be of order $\sim 10^4$ years [22,26], which could potentially be significantly longer if the fields self-interact [30,143–146]. This also opens the potential to joint detections with LISA, which could detect GWs emitted by boson clouds for $\mu \gtrsim 10^{-19} \text{ eV}$ [13].

This work is supported by the Villum Investigator program supported by the VILLUM Foundation (Grant No. VIL37766) and the DNRF Chair program (Grant No. DNRF162) by the Danish National Research Foundation, and under the European Union’s H2020 ERC Advanced Grant “Black holes: gravitational engines of discovery” Grant Agreement No. Gravitas-101052587, and by Fundação para a Ciência e Tecnologia I. P., Portugal under Project No. 2022.01324.PTDC. X. X. is supported by Deutsche Forschungsgemeinschaft under Germany’s Excellence Strategy EXC2121 “Quantum Universe” Grant No. 390833306. The views and opinions expressed are those of the author only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them. R. B. acknowledges financial support provided by FCT, under the Scientific Employment Stimulus—Individual Call—Grant No. 2020.00470.CEECIND.

*yifan.chen@nbi.ku.dk

[†]xiao.xue@desy.de

[‡]richard.brito@tecnico.ulisboa.pt

[§]vitor.cardoso@nbi.ku.dk

- [1] P. Svrcek and E. Witten, *J. High Energy Phys.* **06** (2006) 051.
- [2] S. A. Abel, M. D. Goodsell, J. Jaeckel, V. V. Khoze, and A. Ringwald, *J. High Energy Phys.* **07** (2008) 124.
- [3] A. Arvanitaki, S. Dimopoulos, S. Dubovsky, N. Kaloper, and J. March-Russell, *Phys. Rev. D* **81**, 123530 (2010).
- [4] M. Goodsell, J. Jaeckel, J. Redondo, and A. Ringwald, *J. High Energy Phys.* **11** (2009) 027.
- [5] J. Preskill, M. B. Wise, and F. Wilczek, *Phys. Lett.* **120B**, 127 (1983).
- [6] L. F. Abbott and P. Sikivie, *Phys. Lett.* **120B**, 133 (1983).
- [7] M. Dine and W. Fischler, *Phys. Lett.* **120B**, 137 (1983).
- [8] A. E. Nelson and J. Scholtz, *Phys. Rev. D* **84**, 103501 (2011).
- [9] W. Hu, R. Barkana, and A. Gruzinov, *Phys. Rev. Lett.* **85**, 1158 (2000).
- [10] L. Hui, *Annu. Rev. Astron. Astrophys.* **59**, 247 (2021).
- [11] R. Penrose and R. M. Floyd, *Nature (London)* **229**, 177 (1971).
- [12] Y. B. Zel’Dovich, *JETP Lett.* **14**, 180 (1971), http://jetpletters.ru/ps/1604/article_24607.shtml.
- [13] R. Brito, V. Cardoso, and P. Pani, *Lect. Notes Phys.* **906**, 1 (2015).
- [14] S. L. Detweiler, *Phys. Rev. D* **22**, 2323 (1980).
- [15] V. Cardoso and S. Yoshida, *J. High Energy Phys.* **07** (2005) 009.
- [16] S. R. Dolan, *Phys. Rev. D* **76**, 084001 (2007).
- [17] R. Brito, V. Cardoso, and P. Pani, *Classical Quantum Gravity* **32**, 134001 (2015).
- [18] W. E. East and F. Pretorius, *Phys. Rev. Lett.* **119**, 041101 (2017).
- [19] C. A. R. Herdeiro, E. Radu, and N. M. Santos, *Phys. Lett. B* **824**, 136835 (2022).
- [20] A. Arvanitaki and S. Dubovsky, *Phys. Rev. D* **83**, 044026 (2011).
- [21] A. Arvanitaki, M. Baryakhtar, and X. Huang, *Phys. Rev. D* **91**, 084011 (2015).
- [22] M. Baryakhtar, R. Lasenby, and M. Teo, *Phys. Rev. D* **96**, 035019 (2017).
- [23] R. Brito, S. Ghosh, E. Barausse, E. Berti, V. Cardoso, I. Dvorkin, A. Klein, and P. Pani, *Phys. Rev. D* **96**, 064050 (2017).
- [24] V. Cardoso, O. J. C. Dias, G. S. Hartnett, M. Middleton, P. Pani, and J. E. Santos, *J. Cosmol. Astropart. Phys.* **03** (2018) 043.
- [25] H. Davoudiasl and P. B. Denton, *Phys. Rev. Lett.* **123**, 021102 (2019).
- [26] R. Brito, S. Grillo, and P. Pani, *Phys. Rev. Lett.* **124**, 211101 (2020).
- [27] M. J. Stott, [arXiv:2009.07206](https://arxiv.org/abs/2009.07206).
- [28] C. Ünal, F. Pacucci, and A. Loeb, *J. Cosmol. Astropart. Phys.* **05** (2021) 007,
- [29] A. K. Saha, P. Parashari, T. N. Maity, A. Dubey, S. Bouri, and R. Laha, [arXiv:2208.03530](https://arxiv.org/abs/2208.03530).

- [30] H. Yoshino and H. Kodama, *Prog. Theor. Phys.* **128**, 153 (2012).
- [31] H. Yoshino and H. Kodama, *Prog. Theor. Exp. Phys.* **2014**, 043E02 (2014).
- [32] H. Yoshino and H. Kodama, *Classical Quantum Gravity* **32**, 214001 (2015).
- [33] R. Brito, S. Ghosh, E. Barausse, E. Berti, V. Cardoso, I. Dvorkin, A. Klein, and P. Pani, *Phys. Rev. Lett.* **119**, 131101 (2017).
- [34] M. Isi, L. Sun, R. Brito, and A. Melatos, *Phys. Rev. D* **99**, 084042 (2019); **102**, 049901(E) (2020).
- [35] N. Siemonsen and W. E. East, *Phys. Rev. D* **101**, 024019 (2020).
- [36] L. Sun, R. Brito, and M. Isi, *Phys. Rev. D* **101**, 063020 (2020); **102**, 089902(E) (2020).
- [37] C. Palomba *et al.*, *Phys. Rev. Lett.* **123**, 171101 (2019).
- [38] S. J. Zhu, M. Baryakhtar, M. A. Papa, D. Tsuna, N. Kawanaka, and H.-B. Eggenstein, *Phys. Rev. D* **102**, 063020 (2020).
- [39] L. Tsukada, R. Brito, W. E. East, and N. Siemonsen, *Phys. Rev. D* **103**, 083005 (2021).
- [40] C. Yuan, R. Brito, and V. Cardoso, *Phys. Rev. D* **104**, 044011 (2021).
- [41] R. Abbott *et al.* (KAGRA, VIRGO, and LIGO Scientific Collaborations), *Phys. Rev. D* **105**, 102001 (2022).
- [42] C. Yuan, Y. Jiang, and Q.-G. Huang, *Phys. Rev. D* **106**, 023020 (2022).
- [43] D. Baumann, H. S. Chia, and R. A. Porto, *Phys. Rev. D* **99**, 044001 (2019).
- [44] O. A. Hannuksela, K. W. K. Wong, R. Brito, E. Berti, and T. G. F. Li, *Nat. Astron.* **3**, 447 (2019).
- [45] D. Baumann, H. S. Chia, R. A. Porto, and J. Stout, *Phys. Rev. D* **101**, 083019 (2020).
- [46] V. De Luca and P. Pani, *J. Cosmol. Astropart. Phys.* **08** (2021) 032.
- [47] D. Baumann, G. Bertone, J. Stout, and G. M. Tomaselli, *Phys. Rev. D* **105**, 115036 (2022).
- [48] D. Baumann, G. Bertone, J. Stout, and G. M. Tomaselli, *Phys. Rev. Lett.* **128**, 221102 (2022).
- [49] P. S. Cole, G. Bertone, A. Coogan, D. Gaggero, T. Karydas, B. J. Kavanagh, T. F. M. Spieksma, and G. M. Tomaselli, [arXiv:2211.01362](https://arxiv.org/abs/2211.01362).
- [50] Y. Chen, J. Shu, X. Xue, Q. Yuan, and Y. Zhao, *Phys. Rev. Lett.* **124**, 061102 (2020).
- [51] G.-W. Yuan, Z.-Q. Xia, C. Tang, Y. Zhao, Y.-F. Cai, Y. Chen, J. Shu, and Q. Yuan, *J. Cosmol. Astropart. Phys.* **03** (2021) 018.
- [52] Y. Chen, Y. Liu, R.-S. Lu, Y. Mizuno, J. Shu, X. Xue, Q. Yuan, and Y. Zhao, *Nat. Astron.* **6**, 592 (2022).
- [53] Y. Chen, C. Li, Y. Mizuno, J. Shu, X. Xue, Q. Yuan, Y. Zhao, and Z. Zhou, *J. Cosmol. Astropart. Phys.* **09** (2022) 073.
- [54] R. Roy and U. A. Yajnik, *Phys. Lett. B* **803**, 135284 (2020).
- [55] G. Creci, S. Vandoren, and H. Witek, *Phys. Rev. D* **101**, 124051 (2020).
- [56] R. Roy, S. Vagnozzi, and L. Visinelli, *Phys. Rev. D* **105**, 083002 (2022).
- [57] Y. Chen, R. Roy, S. Vagnozzi, and L. Visinelli, *Phys. Rev. D* **106**, 043021 (2022).
- [58] P. V. P. Cunha and C. A. R. Herdeiro, *Gen. Relativ. Gravit.* **50**, 42 (2018).
- [59] P. V. P. Cunha, C. A. R. Herdeiro, and E. Radu, *Universe* **5**, 220 (2019).
- [60] A. Amorim *et al.* (GRAVITY Collaboration), *Mon. Not. R. Astron. Soc.* **489**, 4606 (2019).
- [61] I. Sengo, P. V. P. Cunha, C. A. R. Herdeiro, and E. Radu, *J. Cosmol. Astropart. Phys.* **01** (2023) 047.
- [62] K. Akiyama *et al.* (Event Horizon Telescope Collaboration), *Astrophys. J. Lett.* **875**, L1 (2019).
- [63] K. Akiyama *et al.* (Event Horizon Telescope Collaboration), *Astrophys. J. Lett.* **875**, L4 (2019).
- [64] K. Akiyama *et al.* (Event Horizon Telescope Collaboration), *Astrophys. J. Lett.* **875**, L6 (2019).
- [65] K. Akiyama *et al.* (Event Horizon Telescope Collaboration), *Astrophys. J. Lett.* **875**, L5 (2019).
- [66] K. Akiyama *et al.* (Event Horizon Telescope Collaboration), *Astrophys. J. Lett.* **910**, L12 (2021).
- [67] K. Akiyama *et al.* (Event Horizon Telescope Collaboration), *Astrophys. J. Lett.* **910**, L13 (2021).
- [68] K. Akiyama *et al.* (Event Horizon Telescope Collaboration), *Astrophys. J. Lett.* **930**, L12 (2022).
- [69] K. Akiyama *et al.* (Event Horizon Telescope Collaboration), *Astrophys. J. Lett.* **930**, L14 (2022).
- [70] K. Akiyama *et al.* (Event Horizon Telescope Collaboration), *Astrophys. J. Lett.* **930**, L15 (2022).
- [71] A. W. Raymond, D. Palumbo, S. N. Paine, L. Blackburn, R. Córdova Rosado, S. S. Doebleman, J. R. Farah, M. D. Johnson, F. Roelofs, R. P. J. Tilanus *et al.*, *Astrophys. J. Suppl. Ser.* **253**, 5 (2021).
- [72] The next generation Event Horizon Telescope, <https://www.ngeht.org/>.
- [73] P. Tiede, M. D. Johnson, D. W. Pesce, D. C. M. Palumbo, D. O. Chang, and P. Galison, *Galaxies* **10**, 111 (2022).
- [74] A. Chael, S. Issaoun, D. w. Pesce, M. D. Johnson, A. Ricarte, C. M. Fromm, and Y. Mizuno, [arXiv:2210.12226](https://arxiv.org/abs/2210.12226).
- [75] M. A. Podurets, *Sov. Astron.* **8**, 868 (1965), <https://adsabs.harvard.edu/full/1965SvA.....8..868P>.
- [76] W. L. Ames and K. S. Thorne, *Astrophys. J.* **151**, 659 (1968).
- [77] J.-P. Luminet, *Astron. Astrophys.* **75**, 228 (1979), <https://articles.adsabs.harvard.edu/full/1979A%26A....75..228L>.
- [78] G. Campbell and R. Matzner, *J. Math. Phys. (N.Y.)* **14**, 1 (1973).
- [79] C. T. Cunningham and J. M. Bardeen, *Astrophys. J. Lett.* **173**, L137 (1972).
- [80] H. Falcke, F. Melia, and E. Agol, *Astrophys. J. Lett.* **528**, L13 (2000).
- [81] S. Chandrasekhar, *The Mathematical Theory of Black Holes* (Oxford University Press, New York, 1983).
- [82] V. Cardoso, A. S. Miranda, E. Berti, H. Witek, and V. T. Zanchin, *Phys. Rev. D* **79**, 064016 (2009).
- [83] V. Cardoso and R. Vicente, *Phys. Rev. D* **100**, 084001 (2019).
- [84] V. Cardoso, F. Duque, and A. Foschi, *Phys. Rev. D* **103**, 104044 (2021).
- [85] T. Johannsen and D. Psaltis, *Astrophys. J.* **718**, 446 (2010).

- [86] S. E. Gralla, D. E. Holz, and R. M. Wald, *Phys. Rev. D* **100**, 024018 (2019).
- [87] M. D. Johnson *et al.*, *Sci. Adv.* **6**, eaaz1310 (2020).
- [88] S. Hadar, M. D. Johnson, A. Lupsasca, and G. N. Wong, *Phys. Rev. D* **103**, 104038 (2021).
- [89] D. Baumann, H. S. Chia, J. Stout, and L. ter Haar, *J. Cosmol. Astropart. Phys.* **12** (2019) 006.
- [90] J. M. Armaleo, D. López Nacir, and F. R. Urban, *J. Cosmol. Astropart. Phys.* **09** (2020) 031.
- [91] Previous constraints assume either couplings to fermions [90,92–94] or massive tensors as the dominant dark matter component [90,95–97].
- [92] M. Hohmann, *Phys. Rev. D* **95**, 124049 (2017).
- [93] C. Talmadge, J. P. Berthias, R. W. Hellings, and E. M. Standish, *Phys. Rev. Lett.* **61**, 1159 (1988).
- [94] M. Sereno and P. Jetzer, *Mon. Not. R. Astron. Soc.* **371**, 626 (2006).
- [95] S. Sun, X.-Y. Yang, and Y.-L. Zhang, *Phys. Rev. D* **106**, 066006 (2022).
- [96] C. Unal, F. R. Urban, and E. D. Kovetz, arXiv:2209.02741.
- [97] J. M. Armaleo, D. López Nacir, and F. R. Urban, *J. Cosmol. Astropart. Phys.* **01** (2020) 053.
- [98] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.130.111401> for a detailed discussion of gravitational atoms, photon ring autocorrelation, oscillation amplitude map on the photon ring, and prospects for lower black hole spins, which includes Refs. [99–107].
- [99] R. Brito, V. Cardoso, and P. Pani, *Phys. Rev. D* **88**, 023514 (2013).
- [100] K. Aoki, K.-i. Maeda, Y. Misonoh, and H. Okawa, *Phys. Rev. D* **97**, 044005 (2018).
- [101] M. Jain and M. A. Amin, *Phys. Rev. D* **105**, 056019 (2022).
- [102] M. Moscibrodzka and C. F. Gammie, *Mon. Not. R. Astron. Soc.* **475**, 43 (2018).
- [103] S. C. Noble, P. K. Leung, C. F. Gammie, and L. G. Book, *Classical Quantum Gravity* **24**, S259 (2007).
- [104] F. Tamburini, B. Thidé, and M. Della Valle, *Mon. Not. R. Astron. Soc.* **492**, L22 (2020).
- [105] J. Feng and Q. Wu, *Mon. Not. R. Astron. Soc.* **470**, 612 (2017).
- [106] K. Akiyama *et al.* (Event Horizon Telescope Collaboration), *Astrophys. J. Lett.* **930**, L16 (2022).
- [107] A. Cruz-Osorio, C. M. Fromm, Y. Mizuno, A. Nathania, Z. Younsi, O. Porth, J. Davelaar, H. Falcke, M. Kramer, and L. Rezzolla, *Nat. Astron.* **6**, 103 (2022).
- [108] A. Chael, KGEO, 2022, <https://github.com/achael/kggeo>.
- [109] B. Carter, *Phys. Rev.* **174**, 1559 (1968).
- [110] J. M. Bardeen, Black Holes, in *Proceedings of the Les Houches Summer School of Theoretical Physics*, edited by C. DeWitt and B. S. DeWitt (Gordon & Breach, New York, 1973), pp. 215–240.
- [111] S. E. Gralla and A. Lupsasca, *Phys. Rev. D* **101**, 044031 (2020).
- [112] S. E. Gralla and A. Lupsasca, *Phys. Rev. D* **101**, 044032 (2020).
- [113] M. Wielgus *et al.* (Event Horizon Telescope Collaboration), *Astrophys. J. Lett.* **930**, L19 (2022).
- [114] The EHT observation of M87* [66,67] and SgrA* [68–70,115] favors a magnetically arrested disk model, which is geometrically thin in the inner region [116–119].
- [115] M. Wielgus, M. Moscibrodzka, J. Vos, Z. Gelles, I. Martí-Vidal, J. Farah, N. Marchili, C. Goddi, and H. Messias, *Astron. Astrophys.* **665**, L6 (2022).
- [116] I. V. Igumenshchev, R. Narayan, and M. A. Abramowicz, *Astrophys. J.* **592**, 1042 (2003).
- [117] R. Narayan, I. V. Igumenshchev, and M. A. Abramowicz, *Publ. Astron. Soc. Jpn.* **55**, L69 (2003).
- [118] J. C. McKinney, A. Tchekhovskoy, and R. D. Blandford, *Mon. Not. R. Astron. Soc.* **423**, 3083 (2012).
- [119] A. Tchekhovskoy, *Launching of Active Galactic Nuclei Jets*, edited by I. Contopoulos *et al.* (Springer, 2015), Vol. 414, p. 45.
- [120] A. Chael, M. D. Johnson, and A. Lupsasca, *Astrophys. J.* **918**, 6 (2021).
- [121] R. Lico *et al.*, arXiv:2301.05699.
- [122] L. Hui, Y. T. A. Law, L. Santoni, G. Sun, G. M. Tomaselli, and E. Trincherini, arXiv:2208.06408.
- [123] A. Khmelnitsky and V. Rubakov, *J. Cosmol. Astropart. Phys.* **02** (2014) 019.
- [124] N. K. Porayko, X. Zhu, Y. Levin, L. Hui, G. Hobbs *et al.*, *Phys. Rev. D* **98**, 102002 (2018).
- [125] M. Bošković, F. Duque, M. C. Ferreira, F. S. Miguel, and V. Cardoso, *Phys. Rev. D* **98**, 024037 (2018).
- [126] H.-K. Guo, Y. Ma, J. Shu, X. Xue, Q. Yuan, and Y. Zhao, *J. Cosmol. Astropart. Phys.* **05** (2019) 015.
- [127] K. Nomura, A. Ito, and J. Soda, *Eur. Phys. J. C* **80**, 419 (2020).
- [128] X. Xue, Z. Q. Xia, X. Zhu, Y. Zhao, J. Shu *et al.* (PPTA Collaboration), *Phys. Rev. Res.* **4**, L012022 (2022).
- [129] G.-W. Yuan, Z.-Q. Shen, Y.-L. S. Tsai, Q. Yuan, and Y.-Z. Fan, *Phys. Rev. D* **106**, 103024 (2022).
- [130] K. Moriyama and S. Mineshige, *Publ. Astron. Soc. Jpn.* **67**, 106 (2015).
- [131] K. Moriyama, S. Mineshige, M. Honma, and K. Akiyama, *Astrophys. J.* **887**, 227 (2019)..
- [132] G. N. Wong, *Astrophys. J.* **909**, 217 (2021).
- [133] P. M. Chesler, L. Blackburn, S. S. Doebleman, M. D. Johnson, J. M. Moran, R. Narayan, and M. Wielgus, *Classical Quantum Gravity* **38**, 125006 (2021).
- [134] A. Gußmann, *J. High Energy Phys.* **08** (2021) 160.
- [135] A. Andrianov, S. Chernov, I. Girin, S. Likhachev, A. Lyakhovets, and Y. Shchekinov, *Phys. Rev. D* **105**, 063015 (2022).
- [136] P. Kocherlakota, S. Biswas, P. S. Joshi, S. Bhattacharyya, C. Chakraborty, and A. Ray, *Mon. Not. R. Astron. Soc.* **490**, 3262 (2019).
- [137] T. Kimpson, K. Wu, and S. Zane, *Mon. Not. R. Astron. Soc.* **497**, 5421 (2020).
- [138] T. Kimpson, K. Wu, and S. Zane, *Astron. Astrophys.* **644**, A167 (2020).
- [139] K. Wu, *Universe* **8**, 78 (2022).
- [140] B. Ben-Salem and E. Hackmann, *Mon. Not. R. Astron. Soc.* **516**, 1768 (2022).
- [141] H.-Y. Schive, T. Chiueh, and T. Broadhurst, *Nat. Phys.* **10**, 496 (2014).

-
- [142] Notice that the decay due to GW emission goes as $\propto 1/(1+t/\tau_{\text{GW}})$ with τ_{GW} the half-life [17,23].
 - [143] H. Fukuda and K. Nakayama, *J. High Energy Phys.* 01 (2020) 128.
 - [144] M. Baryakhtar, M. Galanis, R. Lasenby, and O. Simon, *Phys. Rev. D* **103**, 095019 (2021).
 - [145] W. E. East, *Phys. Rev. Lett.* **129**, 141103 (2022).
 - [146] H. Omiya, T. Takahashi, T. Tanaka, and H. Yoshino, [arXiv:2211.01949](https://arxiv.org/abs/2211.01949).