

Direct Measurement of the Cosmic-Ray Helium Spectrum from 40 GeV to 250 TeV with the Calorimetric Electron Telescope on the International Space Station

O. Adriani,^{1,2} Y. Akaïke,^{3,4} K. Asano,⁵ Y. Asaoka,⁵ E. Berti,^{2,6} G. Bigongiari,^{7,8} W. R. Binns,⁹ M. Bongi,^{1,2} P. Brogi^{7,8,*},
 A. Bruno,¹⁰ J. H. Buckley,⁹ N. Cannady,^{11,12,13} G. Castellini,⁶ C. Checchia,^{7,8} M. L. Cherry,¹⁴ G. Collazuol,^{15,16}
 G. A. de Nolfo,¹⁰ K. Ebisawa,¹⁷ A. W. Ficklin,¹⁴ H. Fuke,¹⁷ S. Gonzi,^{1,2,6} T. G. Guzik,¹⁴ T. Hams,¹¹ K. Hibino,¹⁸
 M. Ichimura,¹⁹ K. Ioka,²⁰ W. Ishizaki,⁵ M. H. Israel,⁹ K. Kasahara,²¹ J. Kataoka,²² R. Kataoka,²³ Y. Katayose,²⁴ C. Kato,²⁵
 N. Kawanaka,²⁰ Y. Kawakubo,¹⁴ K. Kobayashi,^{3,4,†} K. Kohri,²⁶ H. S. Krawczynski,⁹ J. F. Krizmanic,¹² P. Maestro,^{7,8}
 P. S. Marrocchesi,^{7,8} A. M. Messineo,^{8,27} J. W. Mitchell,¹² S. Miyake,²⁸ A. A. Moiseev,^{12,13,29} M. Mori,³⁰ N. Mori,²
 H. M. Motz,³¹ K. Munakata,²⁵ S. Nakahira,¹⁷ J. Nishimura,¹⁷ S. Okuno,¹⁸ J. F. Ormes,³² S. Ozawa,³³ L. Pacini,^{2,6} P. Papini,²
 B. F. Rauch,⁹ S. B. Ricciarini,^{2,6} K. Sakai,^{11,12,13} T. Sakamoto,³⁴ M. Sasaki,^{12,13,29} Y. Shimizu,¹⁸ A. Shiomi,³⁵ P. Spillantini,¹
 F. Stolzi,^{7,8} S. Sugita,³⁴ A. Sulaj,^{7,8} M. Takita,⁵ T. Tamura,¹⁸ T. Terasawa,⁵ S. Torii,³ Y. Tsunesada,^{36,37} Y. Uchihori,³⁸
 E. Vannuccini,² J. P. Wefel,¹⁴ K. Yamaoka,³⁹ S. Yanagita,⁴⁰ A. Yoshida,³⁴ K. Yoshida,²¹ and W. V. Zober⁹

(CALET Collaboration)

¹*Department of Physics, University of Florence, Via Sansone, 1–50019, Sesto Fiorentino, Italy*

²*INFN Sezione di Firenze, Via Sansone, 1–50019, Sesto Fiorentino, Italy*

³*Waseda Research Institute for Science and Engineering, Waseda University, 17 Kikuicho, Shinjuku, Tokyo 162-0044, Japan*

⁴*JEM Utilization Center, Human Spaceflight Technology Directorate, Japan Aerospace Exploration Agency, 2-1-1 Sengen, Tsukuba, Ibaraki 305-8505, Japan*

⁵*Institute for Cosmic Ray Research, The University of Tokyo, 5-1-5 Kashiwa-no-Ha, Kashiwa, Chiba 277-8582, Japan*

⁶*Institute of Applied Physics (IFAC), National Research Council (CNR), Via Madonna del Piano, 10, 50019, Sesto Fiorentino, Italy*

⁷*Department of Physical Sciences, Earth and Environment, University of Siena, via Roma 56, 53100 Siena, Italy*

⁸*INFN Sezione di Pisa, Polo Fibonacci, Largo B. Pontecorvo, 3–56127 Pisa, Italy*

⁹*Department of Physics and McDonnell Center for the Space Sciences, Washington University, One Brookings Drive, St. Louis, Missouri 63130-4899, USA*

¹⁰*Heliospheric Physics Laboratory, NASA/GSFC, Greenbelt, Maryland 20771, USA*

¹¹*Center for Space Sciences and Technology, University of Maryland, Baltimore County, 1000 Hilltop Circle, Baltimore, Maryland 21250, USA*

¹²*Astroparticle Physics Laboratory, NASA/GSFC, Greenbelt, Maryland 20771, USA*

¹³*Center for Research and Exploration in Space Sciences and Technology, NASA/GSFC, Greenbelt, Maryland 20771, USA*

¹⁴*Department of Physics and Astronomy, Louisiana State University, 202 Nicholson Hall, Baton Rouge, Louisiana 70803, USA*

¹⁵*Department of Physics and Astronomy, University of Padova, Via Marzolo, 8, 35131 Padova, Italy*

¹⁶*INFN Sezione di Padova, Via Marzolo, 8, 35131 Padova, Italy*

¹⁷*Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, 3-1-1 Yoshinodai, Chuo, Sagami-hara, Kanagawa 252-5210, Japan*

¹⁸*Kanagawa University, 3-27-1 Rokkakubashi, Kanagawa, Yokohama, Kanagawa 221-8686, Japan*

¹⁹*Faculty of Science and Technology, Graduate School of Science and Technology, Hirosaki University, 3, Bunkyo, Hirosaki, Aomori 036-8561, Japan*

²⁰*Yukawa Institute for Theoretical Physics, Kyoto University, Kitashirakawa Oiwake-cho, Sakyo-ku, Kyoto, 606-8502, Japan*

²¹*Department of Electronic Information Systems, Shibaura Institute of Technology, 307 Fukasaku, Minuma, Saitama 337-8570, Japan*

²²*School of Advanced Science and Engineering, Waseda University, 3-4-1 Okubo, Shinjuku, Tokyo 169-8555, Japan*

²³*National Institute of Polar Research, 10-3, Midori-cho, Tachikawa, Tokyo 190-8518, Japan*

²⁴*Faculty of Engineering, Division of Intelligent Systems Engineering, Yokohama National University, 79-5 Tokiwadai, Hodogaya, Yokohama 240-8501, Japan*

²⁵*Faculty of Science, Shinshu University, 3-1-1 Asahi, Matsumoto, Nagano 390-8621, Japan*

²⁶*Institute of Particle and Nuclear Studies, High Energy Accelerator Research Organization, 1-1 Oho, Tsukuba, Ibaraki, 305-0801, Japan*

²⁷*University of Pisa, Polo Fibonacci, Largo B. Pontecorvo, 3–56127 Pisa, Italy*

²⁸*Department of Electrical and Electronic Systems Engineering, National Institute of Technology (KOSEN), Ibaraki College, 866 Nakane, Hitachinaka, Ibaraki 312-8508, Japan*

²⁹*Department of Astronomy, University of Maryland, College Park, Maryland 20742, USA*

³⁰*Department of Physical Sciences, College of Science and Engineering, Ritsumeikan University, Shiga 525-8577, Japan*

³¹*Faculty of Science and Engineering, Global Center for Science and Engineering, Waseda University, 3-4-1 Okubo, Shinjuku, Tokyo 169-8555, Japan*

³²*Department of Physics and Astronomy, University of Denver,**Physics Building, Room 211, 2112 East Wesley Avenue, Denver, Colorado 80208-6900, USA*³³*Quantum ICT Advanced Development Center, National Institute of Information and Communications Technology, 4-2-1 Nukui-Kitamachi, Koganei, Tokyo 184-8795, Japan*³⁴*College of Science and Engineering, Department of Physics and Mathematics, Aoyama Gakuin University, 5-10-1 Fuchinobe, Chuo, Sagami-hara, Kanagawa 252-5258, Japan*³⁵*College of Industrial Technology, Nihon University, 1-2-1 Izumi, Narashino, Chiba 275-8575, Japan*³⁶*Graduate School of Science, Osaka Metropolitan University, Sugimoto, Sumiyoshi, Osaka 558-8585, Japan*³⁷*Nambu Yoichiro Institute for Theoretical and Experimental Physics, Osaka Metropolitan University, Sugimoto, Sumiyoshi, Osaka 558-8585, Japan*³⁸*National Institutes for Quantum and Radiation Science and Technology, 4-9-1 Anagawa, Inage, Chiba 263-8555, Japan*³⁹*Nagoya University, Furo, Chikusa, Nagoya 464-8601, Japan*⁴⁰*College of Science, Ibaraki University, 2-1-1 Bunkyo, Mito, Ibaraki 310-8512, Japan*

(Received 19 January 2023; accepted 3 April 2023; published 27 April 2023)

We present the results of a direct measurement of the cosmic-ray helium spectrum with the CALET instrument in operation on the International Space Station since 2015. The observation period covered by this analysis spans from October 13, 2015, to April 30, 2022 (2392 days). The very wide dynamic range of CALET allowed for the collection of helium data over a large energy interval, from ~ 40 GeV to ~ 250 TeV, for the first time with a single instrument in low Earth orbit. The measured spectrum shows evidence of a deviation of the flux from a single power law by more than 8σ with a progressive spectral hardening from a few hundred GeV to a few tens of TeV. This result is consistent with the data reported by space instruments including PAMELA, AMS-02, and DAMPE and balloon instruments including CREAM. At higher energy we report the onset of a softening of the helium spectrum around 30 TeV (total kinetic energy). Though affected by large uncertainties in the highest energy bins, the observation of a flux reduction turns out to be consistent with the most recent results of DAMPE. A double broken power law is found to fit simultaneously both spectral features: the hardening (at lower energy) and the softening (at higher energy). A measurement of the proton to helium flux ratio in the energy range from 60 GeV/n to about 60 TeV/n is also presented, using the CALET proton flux recently updated with higher statistics.

DOI: [10.1103/PhysRevLett.130.171002](https://doi.org/10.1103/PhysRevLett.130.171002)

Introduction.—The observation of spectral features departing from a single power law in the energy spectra of cosmic-ray nuclei can provide additional insight into the general phenomenology of cosmic-ray (CR) acceleration and propagation in the Galaxy. The deviations observed by several experiments [1–15] are not easily accommodated within the conventional models of Galactic cosmic-ray acceleration and propagation. These unexpected features have prompted new theoretical interpretations in terms of revised acceleration and propagation mechanisms, as well as the possible contribution of local sources in the injection spectra of Galactic cosmic rays [16–31]. Therefore, accurate measurements of the high-energy spectra of individual elements and of their flux ratios (most notably secondary to primary) are of particular interest to parametrize the energy dependence of spectral features in terms of spectral index variations and smoothness parameters. Input from the new

instruments launched to low Earth orbit in the last decade can provide additional discrimination power among the proposed theoretical models and improve our understanding of CR origin.

At rigidities below a few TV, measurements are carried out either by magnetic spectrometers [8,9] or calorimeters [4,7,10,32,33]. The latter can reach a region of higher energies where new spectral features have been recently observed [1,2,34].

The Calorimetric Electron Telescope (CALET) [35–38] is a space-based instrument equipped with a thick homogeneous calorimeter, optimized for the measurement of the all-electron spectrum [39,40], yet with excellent capabilities to measure the hadronic component of cosmic rays including proton, light, and heavy nuclei (up to nickel and above) [2,14,41,42] in the energy range up to ~ 1 PeV. In this Letter, we present a direct measurement of the cosmic-ray helium spectrum in kinetic energy E from 40 GeV to 250 TeV with CALET.

CALET instrument.—CALET is an all-calorimetric instrument, consisting of three main subdetectors. A charge detector (CHD) is followed by a 3 radiation-length (X_0) thick imaging calorimeter (IMC) and by a 27 X_0 thick total absorption calorimeter (TASC). The CHD, positioned at the

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

top of the apparatus, consists of a two layered hodoscope of plastic scintillator paddles, arranged along two orthogonal directions. The IMC is a fine grained sampling calorimeter alternating thin layers of Tungsten absorber with x, y layers of scintillating fibers (with 1 mm^2 cross section) read out individually. It reconstructs the early shower profile and the trajectory of the impinging particle with good angular resolution, also providing an independent charge measurement via multiple dE/dx sampling [43]. The TASC is a homogeneous calorimeter with 12 layers of tightly packed lead-tungstate (PbWO_4) logs, providing an energy measurement over a very large dynamic range (more than 6 orders of magnitude) spanning four different gain ranges [44]. A more complete description of the instrument is given in the Supplemental Material of Ref. [39].

The instrument was launched on August 19, 2015, and placed on the JEM EF (Japanese Experiment Module Exposed Facility) on the International Space Station. Scientific observations [38] started on October 13, 2015, and smooth and continuous operations have taken place since then.

Data analysis.—Flight data collected from October 13, 2015, to April 30, 2022, were analyzed (2392 days). The total observation live time is 48 459.7 hours, and the live time fraction to total time is about 84.4%. The data analysis generally follows the same procedures used for the CALET analysis of protons [2,15], C-O [14], Fe [41], and Ni [42].

A highly efficient reconstruction of hadronic tracks is of primary importance for the flux measurement. The combinatorial Kalman filter tracking algorithm (KF) [45], already used in the proton spectrum analysis [15], provides good performances also for helium tracks.

The shower energy of each event is calculated as the TASC energy deposit sum (hereafter E_{TASC}), and is calibrated using penetrating protons and He particles selected in flight by a dedicated trigger mode. A seamless stitching of adjacent gain ranges is performed on flight data and complemented by the confirmation of the instrument linearity over the whole range during preflight ground measurements with a UV pulsed laser, as described in Ref. [44].

Time-dependent variations occurring during the long-term observation period are also corrected for each sensor, using penetrating particles as gain monitor [39].

Detailed Monte Carlo (MC) simulations have been performed, based on the EPICS simulation package [46,47]. In order to assess the relatively large uncertainties in the modeling of hadronic interactions, a series of beam tests were carried out at the CERN-SPS using the CALET beam test model [48–50]. Trigger efficiency and energy response derived from MC simulations were tuned using the beam test results obtained in 2015 with ion beams of 13, 19, and 150 GeV/n. For helium nuclei a shower energy correction of 10.4% (8%) at 13(19) GeV/n was applied, while a 3.2% energy independent correction was applied at

150 GeV/n and above. A log-linear interpolation provided the correction factors for intermediate energies not measured at CERN. No correction is applied to the trigger efficiency since beam test measurements are consistent with the MC simulations.

In the analysis of hadrons, especially in the high-energy region where no beam test calibrations are possible, a comparison between different MC models is mandatory. To this extent, we have run simulations with FLUKA [51–53] and compared them with EPICS.

A preselection of well-reconstructed and well-contained events is applied, prior to charge identification, to minimize the background contamination of the selected helium sample. The following criteria are applied.

Trigger: Only events taken with the onboard high-energy trigger mode are retained. This mode is designed to ensure maximum exposure to electrons above 10 GeV and to other high-energy shower events. Consistency between MC and flight data (FD) for triggered events is obtained by applying an offline trigger filter requiring more severe conditions than the onboard trigger. It removes residual effects due to positional and temporal variations of the detector gain.

Track quality cut: Selected events are required to have a good primary track candidate reconstructed in both views with the KF algorithm. A minimum number of points are required for each track segment, and a χ^2 cut is applied. In this way an angular resolution for He nuclei of about 0.1° and an impact point (IP) resolution of $\sim 400 \mu\text{m}$ on the CHD top layer are achieved.

Geometrical condition: The reconstructed events are required to traverse the whole detector (i.e., from CHD top to TASC bottom, with 2 cm clearance from the edges of the TASC top layer) and be contained inside a fiducial region (acceptance A_1), with a geometric factor (GF) of $0.051 \text{ m}^2 \text{ sr}$ ($\sim 49\%$ of the total GF).

Electron rejection: An electron rejection cut is applied, based on a fractional quantity known as “Molière concentration along the track” and calculated by summing all energy deposits inside one Molière radius around each IMC fiber matched to the track and normalized to the total energy deposit sum in the IMC. By requiring this quantity to be less than 0.75, when the fraction of the TASC energy deposited in the last layer is greater than 0.01, more than 90% of electrons are rejected while retaining a very high efficiency for helium nuclei ($> 99.9\%$ for $E > 50 \text{ GeV}$).

Off-Acceptance rejection cuts: Hadronic interactions and the combinatorial track reconstruction are responsible for the occasional misidentification of one of the secondary tracks as the primary track. This results in a number of events erroneously reconstructed inside the fiducial acceptance A_1 . To reject most of these events, different topological cuts are applied using the TASC information. The fractional energy deposit in each of the first two TASC

layers is required to be less than 0.3 to reject laterally incident tracks. The residual between the impact points of a track onto the first two layers of the TASC and the center of gravity of the corresponding energy deposits is required (consistency cut) not to exceed the size of two lead-tungstate logs (± 2 cm). Taking advantage of the TASC granularity, the shower axis is reconstructed with the method of moments (see Ref. [54] for details), and is required to cross the TASC-X1 layer. This cut rejects, with very high efficiency, lateral events erroneously reconstructed inside the fiducial region. A small correction (of a few %) is applied to the cut efficiency to take into account small discrepancies between FD and MC.

The identification of cosmic-ray nuclei via a measurement of their charge is carried out with two independent subsystems that are routinely used to cross calibrate each other: the CHD and the IMC. Tracking allows one to select the CHD paddles crossed by the primary particle and, after application of position and time-dependent calibrations and corrections [44], the information from the two CHD layers is combined into a single charge estimator. The IMC, being equipped with individually read out scintillating fibers, provides multiple dE/dx measurements up to a maximum of 16 samples. The impact point of the impinging particle is reconstructed at first [43], and only the dE/dx ionization clusters from the layers upstream of the IP are used. The charge is evaluated as the truncated mean of the valid samples rejecting 30% of the highest ones. The nonlinear response due to the saturation of the scintillation light in the fibers is corrected for, both in IMC and CHD, by fitting the light yield according to a quenching model described in Refs. [55,56].

To mitigate the effects of the increase of the back-scattered background with energy, both charge measurements are calibrated to the nominal peak positions. This calibration is applied separately to FD and MC simulations by EPICS and FLUKA. To ensure a perfect match between FD and MC, the MC data are finely tuned with FD (separately for EPICS and FLUKA), fitting the proton and helium charge distributions in several energy slices with an asymmetric Landau distribution convoluted with a Gaussian. The full width at half maximum (FWHM) and peak position of the charge distribution are extracted for each energy slice and used, on an event by event basis, to finely tune the MC distributions and to perform an energy dependent charge cut, resulting in an almost flat charge selection efficiency ($\sim 65\%$). More details are given in the Supplemental Material [57].

Background contamination is estimated from MC simulations of protons, helium, and from FD, as a function of the observed energy. The MC simulations are used to evaluate the relative contributions and the FD to assess the proton and helium relative abundances. Charge contamination from protons misidentified as helium is the dominant component. Other not negligible contributions come from

off-acceptance helium and protons misreconstructed inside the acceptance $A1$. Depending on the energy, the estimated overall contamination ranges from a few percent to $\sim 20\%$ at the highest energies where the proton background becomes dominant. The estimated background is then subtracted bin by bin from the dN/dE distribution of helium candidates.

In order to take into account the relatively limited energy resolution (the observed energy fraction is around 35% and the energy resolution is 30–40%), energy unfolding is necessary to correct for significant bin-to-bin migration effects and to infer the primary particle energy. In this analysis, we applied an iterative unfolding method based on the Bayes theorem [59] implemented in the RooUnfold package [60] in ROOT [61], using the response matrix derived from the MC. Convergence is obtained within two iterations, given the relatively accurate prior distribution obtained from the previous observations of AMS-02 [3] and CREAM-I [7]. The energy bin width is chosen to be commensurate with the resolution of the TASC.

The energy spectrum is obtained from the unfolded energy distribution as follows:

$$\Phi(E) = \frac{N(E)}{\Delta E \times \varepsilon(E) \times S\Omega \times T} \quad (1)$$

$$N(E) = U[N_{\text{obs}}(E_{\text{TASC}}) - N_{\text{bg}}(E_{\text{TASC}})] \quad (2)$$

where ΔE denotes the energy bin width; E is the particle kinetic energy, calculated as the geometric mean of the lower and upper bounds of the bin; $N(E)$ is the bin content in the unfolded distribution; $\varepsilon(E)$ the overall selection efficiency (Fig. S2 of the Supplemental Material [57]); T is the live time; $S\Omega$ the “fiducial” geometrical acceptance; U the unfolding procedure; $N_{\text{obs}}(E_{\text{TASC}})$ the bin content of the observed energy distribution (including background); and $N_{\text{bg}}(E_{\text{TASC}})$ the background events in the same bin.

Systematic uncertainties.—The systematic uncertainties can be categorized into energy independent and energy dependent ones. The former includes systematic effects in the normalization and were studied in Ref. [39]. This uncertainty is estimated around 4.1% as the quadratic sum of the uncertainties on live time (3.4%), radiation environment (1.8%), and long-term stability (1.4%).

The energy dependent uncertainties include the following contributions.

Trigger: The absolute calibration of the trigger efficiency was performed at the beam test. The main source of uncertainty comes from the accuracy of the calibration. A possible systematic bias in the trigger efficiency due to normalization was included in the uncertainty, by scanning the offline trigger threshold applied to the TASC-X1 signal between 100 and 150 MIP signals.

Shower energy correction: The absolute calibration of the energy response in the low-energy region was carried out using the beam test data. Both the accuracy of the calibration and the uncertainty in the model used to fit the test beam data are taken into account in the systematics.

Track reconstruction and acceptance: The effects of tracking on the flux were evaluated by studying its dependence on the goodness-of-tracking cuts. To investigate the uncertainty of the acceptance, restricted acceptance regions have been studied, and the resultant fluxes were compared.

Background subtraction: Background subtraction is only slightly dependent on the simulated spectral shape. Different reweighting functions (including E^α with $-2.9 \leq \alpha \leq -2.5$) were adopted for the MC spectrum, and the relative differences with respect to the reference case were included in the systematic uncertainty for each energy bin.

Unfolding: The uncertainties from the unfolding procedure were evaluated by applying different response matrices computed by varying the spectral index (between -2.9 and -2.5) of the MC generation spectrum, or the number of iterations of the Bayesian method.

Charge identification (ID) and off-acceptance rejection cuts: The flux stability against the selection cut efficiencies was studied around the reference value, and the differences with respect to the reference case were accounted as systematic error. The thresholds of each cut were varied separately in an appropriate range (± 1 FWHM for the charge ID cut) around the reference value, and the differences versus the reference case were accounted for as systematic error.

MC model: A second Monte Carlo (FLUKA) is used to evaluate the smearing matrix and the relevant selection efficiencies. For each bin, a systematic error is obtained by a comparison of FLUKA with EPICS results.

Considering all of the above contributions, the total systematic uncertainty remains below 10% up to ~ 60 TeV. Above it increases moderately, remaining commensurate with the statistical error as summarized in Fig. S5 of the Supplemental Material [57] where the total uncertainty is shown with all the relevant contributions listed above.

Two independent helium analyses were carried out by separate groups inside the CALET collaboration, using different event selections and background rejection procedures. The results of the two analyses are consistent with each other within the errors.

Results.—The energy spectrum of CR helium, as measured by CALET in an interval of kinetic energy per particle from ~ 40 GeV to ~ 250 TeV, is shown in Fig. 1 where the statistical and systematic uncertainties are bounded within a gray band. The measured helium flux and the statistical and systematic errors are tabulated in




FIG. 1. Cosmic-ray helium spectrum measured by CALET (red markers), compared with previous direct observations [1,3,7]. The error bars represent only the statistical error; the gray band represents the quadratic sum of statistical and systematic error. The light violet colored band shows the systematic uncertainty of Ref. [1].

Table I of the Supplemental Material [57]. The CALET spectrum is compared with previous observations from space-based [1,3] and balloon-borne [7,10] experiments. Our spectrum is in good agreement with the very accurate measurements by AMS-02 in the lower energy region below a few TeV, as well as with the measurements from calorimetric instruments in the higher energy region, in particular with the recent measurement of DAMPE [1].

In Fig. 2, a fit of CALET data with a “double broken power law” (DBPL) [Eq. (3)] is shown in the energy range from 60 GeV to 250 TeV:

$$\Phi(E) = C \left(\frac{E}{\text{GeV}} \right)^\gamma \left[1 + \left(\frac{E}{E_0} \right)^{\frac{\Delta\gamma}{\gamma}} \right]^{-\frac{\Delta\gamma}{\gamma}} \left[1 + \left(\frac{E}{E_1} \right)^{\frac{\Delta\gamma_1}{\gamma_1}} \right]^{-\frac{\Delta\gamma_1}{\gamma_1}} \quad (3)$$




FIG. 2. Fit of CALET data with a DBPL function [Eq. (3)]. The result is consistent with other recent measurements [1] within the errors. Both statistical and systematic uncertainties are taken into account [57].




FIG. 3. Energy dependence of the spectral index calculated within a sliding energy window for CALET data. The spectral index is determined for each point by fitting the data using ± 2 bins. The gray band indicates the uncertainty range including systematics.

A progressive hardening from a few hundred GeV to a few tens TeV is observed. The fit returns a power law index of $\gamma = -2.703^{+0.005}_{-0.006}(\text{stat})^{+0.032}_{-0.009}(\text{syst})$, $\Delta\gamma = 0.25^{+0.02}_{-0.01}(\text{stat})^{+0.02}_{-0.03}(\text{syst})$, first break energy $E_0 = 1319^{+113}_{-93}(\text{stat})^{+267}_{-124}(\text{syst})$ GeV, and smoothness parameter $S = 2.7^{+0.6}_{-0.5}(\text{stat})^{+3.0}_{-0.9}(\text{syst})$. The onset of a flux softening above a few tens of TeV is also observed, with a second spectral index variation $\Delta\gamma_1 = -0.22^{+0.07}_{-0.10}(\text{stat})^{+0.03}_{-0.04}(\text{syst})$ and second break energy $E_1 = 33.2^{+9.8}_{-6.2}(\text{stat})^{+1.8}_{-2.3}(\text{syst})$ TeV. Given the relatively large uncertainties of the data in the highest energy bins, the second smoothness parameter S_1 cannot be effectively constrained and is kept fixed at a value of $S_1 = 30$.

The index change $\Delta\gamma$ is proven to be different from zero by more than 8σ , taking into account both statistical and systematic error [57]. The fit parameters are generally consistent, within the errors, with the recent results of




FIG. 4. The CALET proton [2] and helium fluxes are shown as functions of kinetic energy per nucleon, together with previous measurements from other experiments [1,3,7,9,34,63,64].

DAMPE [1], although $\Delta\gamma_1$ seems to indicate a less pronounced softening in our data.

The spectral hardening and softening can be easily seen in Fig. 3 where the spectral index is shown as a function of kinetic energy. For each point the spectral index is fitted within a sliding energy window of ± 2 bins. The black marker in the plot represents the index γ with its statistical error, while the gray band represents the quadratic sum of statistical and systematic uncertainties.

Differences between the proton and helium spectra can contribute important constraints on acceleration models (e.g., Ref. [16]). To ease the comparison in Fig. 4, we show the CALET proton spectrum published in Ref. [2] and the helium spectrum from this analysis, in kinetic energy per nucleon. The ^3He contribution to the flux is taken into account assuming the same $^3\text{He}/^4\text{He}$ ratio as measured by AMS-02 [62] and extrapolating it to higher energies with use of a single power-law fit.

Using the CALET proton flux of Ref. [2], we present the p/He flux ratio in Fig. 5 as measured by CALET with high statistical precision in a wide energy range from 60 GeV/n to ~ 60 TeV/n. Both the statistical and systematic errors are shown; details on the systematic uncertainty can be found in the Supplemental Material [57]. Measurements from other experiments [10,65] are included in the same plot. Our result is found to be in agreement with previous measurements from magnetic spectrometers [3,8] up to their maximum detectable rigidity (~ 2 TV), as shown in Fig. S8 of the Supplemental Material [57]. The measured p/He ratio is tabulated in Table II and III of the Supplemental Material [57], as a function of kinetic energy per nucleon and rigidity respectively.




FIG. 5. Energy spectrum of p/He ratio as measured by CALET; the red bars represent statistical error only; the gray band represents the quadratic sum of statistical and systematic errors. Results of previous measurements from CREAM [10] and PAMELA (calorimeter analysis) [65,66] are shown as reference.

Conclusion.—CALET has measured the cosmic-ray helium spectrum covering, for the first time with a single instrument on the International Space Station, the large energy range from 40 GeV to 250 TeV. Our spectrum is not consistent with a single power law (at $> 8\sigma$ level), and its shape confirms the presence of a hardening above a few hundred GeV (where a Smoothly Broken Power Law function fits the spectrum well) and the onset of a flux softening above a few tens TeV. A DBPL fits both spectral features with parameters that are found to be consistent, within the errors, with the most recent results of DAMPE [1]. Using the CALET proton flux [2], we also measured the p/He ratio in the interval from 60 GeV/n to ~ 60 TeV/n. Owing to the partial cancellation of systematic errors in the ratio, this measurement can provide important information on the respective acceleration and propagation mechanisms.

We gratefully acknowledge JAXA's contributions to the development of CALET and to the operations aboard the JEM EF on the International Space Station. We also wish to express our sincere gratitude to ASI (Agenzia Spaziale Italiana) and NASA for their support of the CALET project. This work was supported in part by JSPS Grant-in-Aid for Scientific Research (S) Grant No. 19H05608, and by the MEXT-Supported Program for the Strategic Research Foundation at Private Universities (2011–2015)(Grant No. S1101021) at Waseda University. The CALET effort in Italy is supported by ASI under Agreement No. 2013-018-R.0 and its amendments. The CALET effort in the United States is supported by NASA through Grants No. 80NSSC20K0397, No. 80NSSC20K0399, and No. NNH18ZDA001N-APRA18-0004, and under Award No. 80GSFC21M0002.

*paolo.brogi@unisi.it

†kenkou@aoni.waseda.jp

- [1] F. Alemanno *et al.* (DAMPE Collaboration), *Phys. Rev. Lett.* **126**, 201102 (2021).
- [2] O. Adriani *et al.* (CALET Collaboration), *Phys. Rev. Lett.* **129**, 101102 (2022).
- [3] M. Aguilar *et al.* (AMS Collaboration), *Phys. Rev. Lett.* **115**, 211101 (2015).
- [4] A. Panov *et al.*, *Bull. Russ. Acad. Sci. Phys.* **71**, 494 (2007).
- [5] H. Ahn *et al.*, *Astrophys. J.* **707**, 593 (2009).
- [6] H. Ahn *et al.*, *Astrophys. J. Lett.* **714**, L89 (2010).
- [7] Y. Yoon *et al.*, *Astrophys. J.* **728**, 122 (2011).
- [8] O. Adriani *et al.*, *Science* **332**, 69 (2011).
- [9] M. Aguilar *et al.* (AMS Collaboration), *Phys. Rev. Lett.* **114**, 171103 (2015).
- [10] Y. Yoon *et al.*, *Astrophys. J.* **839**, 5 (2017).
- [11] M. Aguilar *et al.* (AMS Collaboration), *Phys. Rev. Lett.* **119**, 251101 (2017).
- [12] M. Aguilar *et al.* (AMS Collaboration), *Phys. Rev. Lett.* **120**, 021101 (2018).
- [13] M. Aguilar *et al.* (AMS Collaboration), *Phys. Rev. Lett.* **121**, 051103 (2018).
- [14] O. Adriani *et al.* (CALET Collaboration), *Phys. Rev. Lett.* **125**, 251102 (2020).
- [15] O. Adriani *et al.* (CALET Collaboration), *Phys. Rev. Lett.* **122**, 181102 (2019).
- [16] M. A. Malkov, P. H. Diamond, and R. Z. Sagdeev, *Phys. Rev. Lett.* **108**, 081104 (2012).
- [17] A. Erlykin and A. Wolfendale, *Astropart. Phys.* **35**, 449 (2012).
- [18] S. Thoudam and J. Hörandel, *Mon. Not. R. Astron. Soc.* **421**, 1209 (2012).
- [19] G. Bernard, T. Delahaye, Y.-Y. Keum, W. Liu, P. Salati, and R. Taillet, *Astron. Astrophys.* **555**, A48 (2013).
- [20] P. Blasi, E. Amato, and P. D. Serpico, *Phys. Rev. Lett.* **109**, 061101 (2012).
- [21] R. Aloisio and P. Blasi, *J. Cosmol. Astropart. Phys.* **07** (2013) 001.
- [22] S. Thoudam and J. Hörandel, *Astron. Astrophys.* **567**, A33 (2014).
- [23] Y. Ohira and K. Ioka, *Astrophys. J. Lett.* **729**, L13 (2011).
- [24] Y. Ohira, N. Kawanaka, and K. Ioka, *Phys. Rev. D* **93**, 083001 (2016).
- [25] P. Biermann, J. K. Becker, J. Dreyer, A. Meli, E.-S. Seo, and T. Stanev, *Astrophys. J.* **725**, 184 (2010).
- [26] V. Ptuskin, V. Zirakashvili, and E. Seo, *Astrophys. J.* **763**, 47 (2013).
- [27] V. Zatsepin and N. Sokolskaya, *Astron. Astrophys.* **458**, 1 (2006).
- [28] N. Tomassetti, *Astrophys. J. Lett.* **752**, L13 (2012).
- [29] A. Vladimirov, G. Jóhannesson, I. Moskalenko, and T. Porter, *Astrophys. J.* **752**, 68 (2012).
- [30] N. Tomassetti, *Phys. Rev. D* **92**, 063001 (2015).
- [31] C. Evoli, P. Blasi, G. Morlino, and R. Aloisio, *Phys. Rev. Lett.* **121**, 021102 (2018).
- [32] E. Atkin *et al.*, *J. Cosmol. Astropart. Phys.* **07** (2017) 020.
- [33] E. Atkin *et al.*, *JETP Lett.* **108**, 5 (2018).
- [34] Q. An *et al.* (DAMPE Collaboration), *Sci. Adv.* **5**, eaax3793 (2019).
- [35] P. S. Marrocchesi (CALET Collaboration), *Proc. Sci., ICRC2021* (2021) 010.
- [36] S. Torii (CALET Collaboration), *Proc. Sci., ICRC2019* (2019) 142.
- [37] S. Torii *et al.* (CALET Collaboration), *Proc. Sci., ICRC2017* (2017) 1092.
- [38] Y. Asaoka, Y. Ozawa, S. Torii *et al.* (CALET Collaboration), *Astropart. Phys.* **100**, 29 (2018).
- [39] O. Adriani *et al.* (CALET Collaboration), *Phys. Rev. Lett.* **119**, 181101 (2017).
- [40] O. Adriani *et al.* (CALET Collaboration), *Phys. Rev. Lett.* **120**, 261102 (2018).
- [41] O. Adriani *et al.* (CALET Collaboration), *Phys. Rev. Lett.* **126**, 241101 (2021).
- [42] O. Adriani *et al.* (CALET Collaboration), *Phys. Rev. Lett.* **128**, 131103 (2022).
- [43] P. Brogi *et al.* (CALET Collaboration), *Proc. Sci., ICRC2015* (2015) 595.
- [44] Y. Asaoka, Y. Akaïke, Y. Komiya, R. Miyata, S. Torii *et al.* (CALET Collaboration), *Astropart. Phys.* **91**, 1 (2017).

- [45] P. Maestro, N. Mori *et al.* (CALET Collaboration), *Proc. Sci., ICRC2017* (**2017**) 208.
- [46] K. Kasahara, in *Proceedings of 24th International Cosmic Ray Conference (Rome, Italy)* (1995), Vol. 1, p. 399, <https://adsabs.harvard.edu/full/1995ICRC....1..399K>.
- [47] EPICS and COSMOS versions are 9.20 and 8.00, respectively.
- [48] Y. Akaïke *et al.* (CALET Collaboration), in *Proceedings of 33rd International Cosmic Ray Conference (ICRC2013)* 726 (2013), <https://inspirehep.net/literature/1412775>.
- [49] T. Niita, S. Torii, Y. Akaïke, Y. Asaoka, K. Kasahara, S. Ozawa, and T. Tamura, *Adv. Space Res.* **55**, 2500 (2015).
- [50] Y. Akaïke *et al.* (CALET Collaboration), *Proc. Sci., ICRC2015* (**2015**) 613.
- [51] T. Böhlen, *Nucl. Data Sheets* **120**, 211 (2014).
- [52] A. Ferrari, P. Sala, A. Fassó, and J. Ranft, in Reports No. INFN/TC_05/11, No. SLAC-R-773, No. CERN-2005-10, 2005, [10.5170/CERN-2005-010](https://arxiv.org/abs/10.5170/CERN-2005-010).
- [53] The version of FLUKA is Fluka2011.2c.4.
- [54] J. J. Gomez *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **262**(2–3), 284 (1987).
- [55] R. Voltz, J. Lopes da Silva, G. Laustriat, and A. Coche, *JChPh* **45**, 3306 (1966).
- [56] G. Tarle *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. B* **6**, 504 (1985).
- [57] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.130.171002>, for supporting figures and the tabulated fluxes as well as the description of data analysis procedure and the detailed assessment of systematic uncertainties, which includes Ref. [58].
- [58] P. Brogi *et al.* (CALET Collaboration), *Proc. Sci., ICRC2021* (**2021**) 101.
- [59] G. D’Agostini, *Nucl. Instrum. Methods Phys. Res., Sect. A* **362**, 487 (1995).
- [60] T. Adye, in [arXiv:1105.1160v1](https://arxiv.org/abs/1105.1160v1).
- [61] R. Brun and F. Rademakers, *Nucl. Instrum. Methods Phys. Res., Sect. A* **389**, 81 (1997).
- [62] M. Aguilar *et al.* (AMS Collaboration), *Phys. Rev. Lett.* **123**, 181102 (2019).
- [63] V. Grebenyuk *et al.* (NUCLEON Collaboration), *Adv. Space Res.* **64**, 2546 (2019).
- [64] A. Panov *et al.* (ATIC Collaboration), *Bull. Russian Acad. Sci.* **73**, 564 (2009).
- [65] O. Adriani *et al.*, *Adv. Space Res.* **51**, 2, 219(2013).
- [66] D. Maurin *et al.*, *Universe* **6**, 8, 102 (2020).