


Observation of a Resonant Structure near the $D_s^+ D_s^-$ Threshold in the $B^+ \rightarrow D_s^+ D_s^- K^+$ Decay

R. Aaij *et al.**
(LHCb Collaboration)

 (Received 28 October 2022; revised 6 February 2023; accepted 28 February 2023; published 14 August 2023)

An amplitude analysis of the $B^+ \rightarrow D_s^+ D_s^- K^+$ decay is carried out to study for the first time its intermediate resonant contributions, using proton-proton collision data collected with the LHCb detector at center-of-mass energies of 7, 8, and 13 TeV. A near-threshold peaking structure, referred to as $X(3960)$, is observed in the $D_s^+ D_s^-$ invariant-mass spectrum with significance greater than 12 standard deviations. The mass, width, and the quantum numbers of the structure are measured to be $3956 \pm 5 \pm 10$ MeV, $43 \pm 13 \pm 8$ MeV, and $J^{PC} = 0^{++}$, respectively, where the first uncertainties are statistical and the second systematic. The properties of the new structure are consistent with recent theoretical predictions for a state composed of $c\bar{c}s\bar{s}$ quarks. Evidence for an additional structure is found around 4140 MeV in the $D_s^+ D_s^-$ invariant mass, which might be caused either by a new resonance with the 0^{++} assignment or by a $J/\psi\phi \leftrightarrow D_s^+ D_s^-$ coupled-channel effect.

DOI: [10.1103/PhysRevLett.131.071901](https://doi.org/10.1103/PhysRevLett.131.071901)

Exotic hadrons (hadrons that are not composed either of a quark-antiquark pair or of three quarks or three antiquarks are collectively called exotic hadrons) play a crucial role in studies of quantum chromodynamics (QCD), and provide a unique window to understand the nature of the strong interaction. Dozens of charged states with hidden charm or beauty, which imply exotic nature, such as $Z_c(4430)^+$ [1,2], $Z_b(10610)^+$ [3], $Z_c(3900)^+$ [4–6], $Z_c(4020)^+$ [7,8], $P_c(4450)^+$ [9,10], $Z_{cs}(3985)^+$ [11], $Z_{cs}(4000)^+$ [12], have been recently discovered by various experiments. (The inclusion of charge-conjugate processes is always implied and natural units with $\hbar = c = 1$ are used throughout the Letter.) Over the last two years, the LHCb Collaboration reported three new open-charm tetraquark states, $X_{0,1}(2900)^0$ [13,14] and $T_{cc}(3875)^+$ [15,16], composed of $cs\bar{u}\bar{d}$ and $cc\bar{u}\bar{d}$ quarks, respectively. Interestingly, most of these states have masses close to thresholds of hadron pairs, which may indicate that they are hadronic molecules loosely bound by deuteronlike meson-exchange forces [17–20]. There are a number of other possible explanations, including that these particles are compact multiquark states [21–23], hadroquarkonia in which a $c\bar{c}$ core is bound to light quarks and/or gluons via chromoelectric dipole interactions [24,25], or cusps produced by near-threshold

kinematics involving open-charm hadrons, or other dominant processes [26,27].

The $\chi_{c0}(3930)$ state was observed by the LHCb Collaboration in the $D^+ D^-$ invariant-mass spectrum [14]. The mass and width of this state are consistent with those of the $X(3915)$ resonance observed in the $\omega J/\psi$ invariant-mass spectrum [28–31]. Moreover, the $X(3915)$ has preferred spin (J), parity (P), and charge-parity (C) quantum numbers of $J^{PC} = 0^{++}$ [31,32], so the two states are treated as a single hadron in the following discussions unless otherwise specified. However, the $\chi_{c0}(3930)$ state is not considered to be consistent with being a candidate for either the $\chi_{c0}(2P)$ or $\chi_{c0}(3P)$ state [33–37]. Lebed *et al.* [38] propose that it is the lightest $c\bar{c}s\bar{s}$ state. Calculations based on QCD sum rules [39] favor the $\chi_{c0}(3930)$ state as a $0^{++} [cq][\bar{c}\bar{q}]$ (where $q = u, d$) or $[cs][\bar{c}\bar{s}]$ tetraquark. Recent lattice QCD results also indicate that this state is dominated by the $c\bar{c}s\bar{s}$ constituents [40]. The $D_s^+ D_s^-$ molecular interpretation is also possible, as suggested by the quark delocalization color-screening model [41] and other phenomenological studies [42,43]. All these developments point to a potential resonant structure in the vicinity of the threshold in the $D_s^+ D_s^-$ invariant-mass spectrum.

Previously, only the Belle experiment studied the $D_s^+ D_s^-$ invariant-mass spectrum in processes involving initial-state radiation, where only 1^{--} charmonium(like) states can contribute [44]. The $B^+ \rightarrow D_s^+ D_s^- K^+$ process, given its large branching fraction measured in the accompanying paper [45], provides a good opportunity to study resonances in the $D_s^+ D_s^-$ final states, both scalars and those of higher spin, such as the 0^{++} charmonium(like) states $\chi_{c0}(4500)$ and $\chi_{c0}(4700)$ possibly having an intrinsic $c\bar{c}s\bar{s}$

*Full author list given at the end of the Letter.

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

component [12], the well-known 1^{--} charmonium states, such as $\psi(4040)$, $\psi(4160)$, $\psi(4260)$, $\psi(4415)$, and $\psi(4660)$ [32,46,47].

In this Letter, an amplitude analysis of about 360 reconstructed $B^+ \rightarrow D_s^+ D_s^- K^+$ signal decays is presented, leading to the first observation of a near-threshold peaking structure in the $D_s^+ D_s^-$ system, denoted by $X(3960)$. The analysis is based on proton-proton (pp) collision data collected by the LHCb experiment at center-of-mass energies of 7, 8, and 13 TeV between 2011 and 2018, corresponding to an integrated luminosity of 9 fb^{-1} . The D_s^+ candidates are reconstructed via the $D_s^+ \rightarrow K^- K^+ \pi^+$ decay. The details of the detector, data and simulation, selection criteria, background composition, and B^+ invariant-mass fit can be found in the accompanying article [45].

To improve the resolution on the masses of the two-body combinations that are used in the amplitude analysis, the four momentum of each final-state particle is determined from a kinematic fit [48] where the B^+ mass is constrained to its known value [32]. Figure 1 shows the resulting Dalitz-plot distribution for the $B^+ \rightarrow D_s^+ D_s^- K^+$ signal decays, where the non- B^+ background is subtracted by the $sPlot$ technique [49] with the reconstructed B^+ mass as the discriminating variable. The most evident feature is the band near the $D_s^+ D_s^-$ threshold. To validate that this peaking structure is not due to the combinatorial background, the $D_s^+ D_s^-$ invariant-mass distribution of candidates in the B^+ mass region from 5360 to 5600 MeV is investigated and no peak is observed.

Employing an unbinned maximum-likelihood method, an amplitude fit with the $sFit$ technique [50] is performed to investigate the intermediate states and determine the quantum numbers J^{PC} of any new particle. Two known 1^{--} charmonium states, $\psi(4260)$ and $\psi(4660)$ [32,46,47], and two new 0^{++} X states are needed to fit the structures in the $D_s^+ D_s^-$ spectrum. One of these scalars, $X(3960)$, describes the $D_s^+ D_s^-$ threshold enhancement and the other,

designated $X_0(4140)$, is necessary to model the dip around 4140 MeV, as shown in Fig. 2. The subscript 0 is used to distinguish the latter from the 1^{++} $X(4140)$ state seen in the $J/\psi\phi$ final state [32]. Additionally, an S -wave three-body phase-space function [32] is employed to model the nonresonant (NR) $B^+ \rightarrow D_s^+ D_s^- K^+$ component. Since no significant contribution of any state is observed in either the $D_s^- K^+$ or $D_s^+ K^+$ systems, these five contributions constitute the baseline model.

The helicity formalism [51] is used to construct the amplitude model of the $B^+ \rightarrow D_s^+ D_s^- K^+$ decay, with a similar approach applied to previous LHCb analyses of B^+ and B_s^0 decays to three pseudoscalar particles [14,52–54]. The resonant structure near the $D_s^+ D_s^-$ mass threshold is parametrized by a Flatté-like function [19,32,55] depending on the invariant mass m

$$R(m|M_0, g_j) = \frac{1}{M_0^2 - m^2 - iM_0 \sum_j g_j \rho_j(m)}, \quad (1)$$

where M_0 is the mass of the resonance, g_j denotes the coupling of this resonance to the j th channel, $\rho_j(m)$ is the phase-space factor [32] for the j th two-body decay. When the value of m is below the threshold of the channel j , i.e., $q_j^2 < 0$, an analytic continuation is applied for $q_j = i\sqrt{-q_j^2}$ [55,56]. The total width of the resonance is calculated as $\Gamma_0 = \sum_j g_j \rho_j(M_0)$. In the baseline model, only the $D_s^+ D_s^-$ channel ($j = 1$) is included in the Flatté-like parametrization.

Other resonances are modeled by a relativistic Breit-Wigner function $\text{BW}(m|M_0, \Gamma_0)$ with a mass-dependent width [32]. The radius of each resonance entering the Blatt-Weisskopf barrier factor [57–59] is set to 3 GeV^{-1} , corresponding to about 0.6 fm.

The total probability density function is the squared modulus of the total decay amplitude multiplied by the efficiency, normalized to ensure that the integral over the Dalitz plot is unity. The fit fraction \mathcal{F}_i expresses the fraction of the total rate due to the component i , and the interference fraction \mathcal{I}_{ij} describes the interference between components i and j . They are defined in Eqs. (18) and (19) of Ref. [53], such that $\sum_i \mathcal{F}_i + \sum_{i < j} \mathcal{I}_{ij} = 1$.

As shown in Fig. 2, the two-body mass distributions are well modeled by the baseline amplitude fit. The corresponding numerical results are summarized in Table I, including the mass, width, fit fraction, and significance (\mathcal{S}) of each component. The significance of a given component is evaluated by assuming that the change of twice the negative log-likelihood ($-2 \ln \mathcal{L}$) between the baseline fit and the fit without that component obeys a χ^2 distribution, where the number of degrees of freedom (d.o.f) is given by the change in the number of free parameters. All the components included in the baseline model have a statistical significance greater than three standard deviations (σ),




FIG. 1. Dalitz-plot distribution for the $B^+ \rightarrow D_s^+ D_s^- K^+$ decay after background subtraction.




FIG. 2. Background-subtracted invariant-mass distributions (top left) $m(D_s^+ D_s^-)$, (top right) $m(D_s^+ K^+)$ and (bottom) $m(D_s^- K^+)$ for the $B^+ \rightarrow D_s^+ D_s^- K^+$ signal. The projections of the fit with the baseline amplitude model are also shown.

where the $X(3960)$ and $X_0(4140)$ states are found to be 14.6σ and 4.1σ significant, respectively. The obtained significances for the $X(3960)$ and $X_0(4140)$ resonances are also validated using pseudoexperiments.

The J^{PC} assignment for the system of a pair of oppositely charged pseudoscalar mesons must be in the series 0^{++} , 1^{--} , 2^{++} , etc. States with higher intrinsic spin are not expected to contribute significantly in the current dataset. To determine the $X(3960)$ quantum numbers, fits with the baseline model are performed under alternative J^{PC} hypotheses, 1^{--} , 2^{++} , instead of 0^{++} . The significance to reject a J^{PC} hypothesis is computed as $\sqrt{\Delta(-2 \ln \mathcal{L})}$, where $\Delta(-2 \ln \mathcal{L}) = -[2 \ln \mathcal{L}(0^{++}) - 2 \ln \mathcal{L}(J^{PC})]$, and

indicates the likelihood difference between the fits for the preferred 0^{++} assignment and for each alternative J^{PC} hypothesis. To ensure that for different J^{PC} hypotheses this resonance corresponds to the same particle, the mass and the width are limited to be within a $\pm 3\sigma$ range of the baseline fit results. The 0^{++} assignment is preferred over 1^{--} and 2^{++} hypotheses by 9.3σ and 12.3σ , respectively. Similarly, replacing the baseline 0^{++} assignment by 1^{--} or 2^{++} for the $X_0(4140)$ state deteriorates the fit quality. The 0^{++} assignment is favored over 1^{--} (2^{++}) hypothesis at a 3.5σ (4.2σ) level. Within the baseline model this 0^{++} state produces the dip around 4140 MeV via destructive interference with the 0^{++} NR and $X(3960)$ components, with

TABLE I. Summary of the main results obtained with the baseline model, where the first uncertainty is statistical and the second systematic. The last column shows the signal significance with (without) the systematic uncertainty included.

| Component | J^{PC} | M_0 (MeV) | Γ_0 (MeV) | \mathcal{F} (%) | $S(\sigma)$ |
|--------------|----------|---------------------|-------------------|--------------------------|-------------|
| $X(3960)$ | 0^{++} | $3956 \pm 5 \pm 10$ | $43 \pm 13 \pm 8$ | $25.4 \pm 7.7 \pm 5.0$ | 12.6 (14.6) |
| $X_0(4140)$ | 0^{++} | $4133 \pm 6 \pm 6$ | $67 \pm 17 \pm 7$ | $16.7 \pm 4.7 \pm 3.9$ | 3.8 (4.1) |
| $\psi(4260)$ | 1^{--} | 4230 [60] | 55 [60] | $3.6 \pm 0.4 \pm 3.2$ | 3.2 (3.6) |
| $\psi(4660)$ | 1^{--} | 4633 [32] | 64 [32] | $2.2 \pm 0.2 \pm 0.8$ | 3.0 (3.2) |
| NR | 0^{++} | ... | ... | $46.1 \pm 13.2 \pm 11.3$ | 3.1 (3.4) |

the interference fractions of, respectively, $(-22.4 \pm 6.4)\%$ and $(-5.2 \pm 3.9)\%$, where the uncertainties are statistical only.

Systematic uncertainties on the measured resonance properties are evaluated, and are summarized in Table 1 in Supplemental Material [61]. Corrections, derived from calibration samples, are applied to account for possible discrepancies between data and simulation in the hardware trigger and particle-identification responses. The uncertainty due to the limited size of the simulation samples is evaluated using the bootstrap method [62]. Additional resonances, not included in the baseline model (states in the $D_s^+ D_s^-$ system: $0^{++} \chi_{c0}(4500)$ and $\chi_{c0}(4700)$ [12], $1^{--} \psi(4040)$, $\psi(4160)$, and $\psi(4415)$ [32], and $2^{++} \chi_{c2}(3930)$ [14]; and in the $D_s^- K^+$ system: $0^+ \bar{D}_0^*(2300)^0$ [32], $1^- \bar{D}_1^*(2600)^0$ [32,63] and $\bar{D}_1^*(2760)^0$ [64], and $2^+ \bar{D}_2^*(2460)^0$ [32]) are utilized to estimate the uncertainty due to insufficient consideration of possible amplitude components. None of these states significantly improve the baseline model. The $c \bar{c} s \bar{s}$ candidates $\chi_{c0}(4500)$ and $\chi_{c0}(4700)$ have statistical significances of 0.8σ and 1.3σ , respectively, and their fit fractions are $(0.6 \pm 1.0)\%$ ($<3.5\%$ at 90% confidence level) and $(2.4 \pm 1.8)\%$ ($<6.7\%$ at 90% confidence level), where the uncertainties are statistical. The Blatt-Weisskopf hadron size is varied between 1.5 and 4.5 GeV^{-1} . The fixed masses and widths of two baseline ψ states are varied by their corresponding uncertainties. The Flatté-like parametrization for the $X(3960)$ state is replaced by a constant-width relativistic Breit-Wigner function. The uncertainty due to the possible bias of the *sFit* method is evaluated using pseudoexperiments. The total systematic uncertainties on mass, width, and fit fraction are obtained by adding all contributions in quadrature, assuming that each source is independent. Regarding the total significance for each component in the baseline model, the smallest significance among these systematic tests is selected.

The measured mass and width of the $X(3960)$ state are consistent with those of the $\chi_{c0}(3930)$ meson [14] within 3σ . Assuming that the $X(3960)$ in the $D_s^+ D_s^-$ system and the $\chi_{c0}(3930)$ in the $D^+ D^-$ system are the same state, the baseline model is extended by adding a second channel ($D^+ D^-$) in the Flatté-like parametrization. The corresponding fit projections and numerical results can be found in Supplemental Material [61]. The likelihood is essentially unchanged while the number of d.o.f. is increased by one compared to the baseline fit. The coupling strength of the $X(3960)$ state to $D_s^+ D_s^-$ ($D^+ D^-$) is found to be 0.33 ± 1.18 (0.15 ± 0.33) GeV . The masses and fit fractions of all components are consistent with those in the baseline one-channel Flatté-like model.

In the case that the $X(3960)$ and $\chi_{c0}(3930)$ states are the same particle, the partial width ratio of such an X resonance decaying to $D_s^+ D_s^-$ and $D^+ D^-$ final states is calculated as

$$\frac{\Gamma(X \rightarrow D^+ D^-)}{\Gamma(X \rightarrow D_s^+ D_s^-)} = \frac{\mathcal{B}^{(1)} \mathcal{F}_X^{(1)}}{\mathcal{B}^{(2)} \mathcal{F}_X^{(2)}} = 0.29 \pm 0.09 \pm 0.10 \pm 0.08, \quad (2)$$

where the superscripts (1) and (2) indicate the $B^+ \rightarrow D^+ D^- K^+$ and $B^+ \rightarrow D_s^+ D_s^- K^+$ channels, respectively, $\mathcal{F}_X^{(1)} = (3.70 \pm 0.92)\%$ is the fit fraction of the $\chi_{c0}(3930)$ state in the $B^+ \rightarrow D^+ D^- K^+$ decay [14], $\mathcal{F}_X^{(2)}$ is the fit fraction of the $X(3960)$ resonance presented in this Letter, and the branching fraction ratio $\mathcal{B}^{(1)}/\mathcal{B}^{(2)}$ is taken from the accompanying paper [45]. The first uncertainty is statistical, the second systematic, and the third is due to uncertainties in the measured branching fractions, $\mathcal{B}(D^+ \rightarrow K^- \pi^+ \pi^+)$ and $\mathcal{B}(D_s^+ \rightarrow K^- K^+ \pi^+)$ [32], and the uncertainty on $\mathcal{F}_X^{(1)}$ [14]. This ratio is compatible with that of the couplings mentioned above.

It is well known that the creation of an $s\bar{s}$ quark pair from the vacuum is suppressed relative to $u\bar{u}$ or $d\bar{d}$ pairs. Moreover, the $X \rightarrow D_s^+ D_s^-$ decay, occurring near the threshold, has a rather smaller phase-space factor than that of $X \rightarrow D^+ D^-$. These two features indicate that $\Gamma(X \rightarrow D^+ D^-)$ should be considerably larger than $\Gamma(X \rightarrow D_s^+ D_s^-)$ if X does not have any intrinsic $s\bar{s}$ content. However, the value measured in Eq. (2) contradicts this expectation. This implies that the $X(3960)$ and $\chi_{c0}(3930)$ are either not the same resonance, or they are the same nonconventional charmoniumlike state, for instance, a candidate containing the dominant $c\bar{c}s\bar{s}$ constituents predicted in recent theoretical models [38–43,65]. Further studies are needed to gain insights into the nature of the $D_s^+ D_s^-$ threshold enhancement, in particular the measurement of the relative branching fraction for the $D_{(s)} \bar{D}_{(s)}$ and $\omega J/\psi$ channels, produced in a different environment, such as from two-photon fusion processes by the Belle II experiment.

There is no obvious candidate within conventional charmonium multiplets for $X(3960)$ or $\chi_{c0}(3930)$ assignment. First of all, the mass of the $\chi_{c0}(3930)$ state is far from predictions for the $\chi_{c0}(3P)$, which lies within the range 4131–4292 MeV [33,35]. For the $\chi_{c0}(2P)$ state, most potential models predict a mass in the range 3842–3868 MeV [34–36], except the Godfrey-Isgur model which gives 3916 MeV [33]. Second, the $\chi_{c0}(3930)$ state interpreted as $\chi_{c0}(2P)$ would give too small a mass splitting with respect to the $\chi_{c2}(3930)$ state [14] identified as $\chi_{c2}(2P)$ [33–36]. In addition, interpreting the $\chi_{c0}(3930)$ state as the $\chi_{c0}(2P)$ charmonium would result in inconsistent decay widths, as the Okubo-Zweig-Iizuka (OZI) [66,67] suppressed channel $\chi_{c0}(3930) \rightarrow \omega J/\psi$ has a decay width larger than theoretical expectations, whereas the S -wave OZI-allowed $\chi_{c0}(3930) \rightarrow D\bar{D}$ mode has smaller decay width than the expectations [36,37]. As a consequence, neither the $X(3960)$ nor the $\chi_{c0}(3930)$ is likely to be a pure $\chi_{c0}(2P)$ or $\chi_{c0}(3P)$ charmonium state.

To test the possibility that the dip in the $D_s^+ D_s^-$ invariant mass around 4140 MeV can be produced by the opening of the nearby $J/\psi\phi$ threshold, without introducing an additional resonance, we employ a simple K -matrix model that contains the single resonance $X(3960)$ and two coupled channels, $D_s^+ D_s^-$ and $J/\psi\phi$. The K matrix reads

$$\begin{pmatrix} \mathcal{K}_{D_s^+ D_s^- \rightarrow D_s^+ D_s^-} & \mathcal{K}_{D_s^+ D_s^- \rightarrow J/\psi\phi} \\ \mathcal{K}_{J/\psi\phi \rightarrow D_s^+ D_s^-} & \mathcal{K}_{J/\psi\phi \rightarrow J/\psi\phi} \end{pmatrix} \equiv \begin{pmatrix} \mathcal{K}_{11} & \mathcal{K}_{12} \\ \mathcal{K}_{21} & \mathcal{K}_{22} \end{pmatrix}, \quad (3)$$

where $\mathcal{K}_{12} = \mathcal{K}_{21}$, and the subscripts 1 and 2 represent $D_s^+ D_s^-$ and $J/\psi\phi$ final states, respectively. One possible choice for the 2×2 K -matrix parametrization [32] is

$$\mathcal{K}_{ba}(m) = \sum_R \frac{g_b^R g_a^R}{M_R^2 - m^2} + f_{ba}, \quad (4)$$

where M_R refers to the bare mass of the resonance R , m is the $D_s^+ D_s^-$ invariant mass, g_a^R denotes the bare coupling of the resonance R to the channel a , and the f_{ba} is a real matrix parametrizing the nonpole part of the K matrix. As the $X(3960)$ mass is about 160 MeV lower than the $J/\psi\phi$ threshold and its width is less than 50 MeV, the coupling of the $X(3960)$ state to $J/\psi\phi$ should be negligible, giving $g_2^R = 0$. This results in the $X(3960)$ resonance entering the \mathcal{K}_{11} element only. The production amplitude is expressed in the P -vector formalism [32,68,69], which gives

$$\mathcal{P}_b(m) = \sum_R \frac{\beta_R g_b^R}{M_R^2 - m^2} + \beta_b, \quad (5)$$

where β_R and β_b are complex free parameters due to rescattering effects or missing channels [60]. The amplitude \mathcal{M} is

$$\mathcal{M}_a = \sum_b (I - i\rho\mathcal{K})_{ab}^{-1} \mathcal{P}_b, \quad (6)$$

where $\rho = \text{diag}\{\rho_{11}, \rho_{22}\}$ is the diagonal matrix composed of phase-space factors, I represents the identity matrix, and $a = 1$ for the $D_s^+ D_s^-$ channel under consideration.

The fit demonstrates that the dip around the $J/\psi\phi$ threshold can also be modeled by the $J/\psi\phi \rightarrow D_s^+ D_s^-$ rescattering, and results in a $-2 \ln \mathcal{L}$ that is worse by 6.0, while the number of d.o.f. is increased by one, compared to the baseline fit. The fit projections and numerical results can be found in Supplemental Material [61]. Since the fit quality of the K -matrix parametrization is close to that of the baseline model, a strong conclusion cannot be drawn whether the dip is due to destructive interference with the $X_0(4140)$ resonance or caused by the $J/\psi\phi \rightarrow D_s^+ D_s^-$ rescattering.

In addition, it is found that the fits with the two-channel Flatté-like and K -matrix parametrizations are unstable, due to having too many free parameters for the limited data

sample size. Consequently, the statistical uncertainties for some parameters are large. Therefore, neither of these parametrizations are taken as the baseline model.

In conclusion, the first amplitude analysis of the $B^+ \rightarrow D_s^+ D_s^- K^+$ decay is performed using pp collision data with an integrated luminosity of 9 fb^{-1} collected by the LHCb experiment between 2011 and 2018. A peaking structure near the $D_s^+ D_s^-$ mass threshold, denoted as $X(3960)$, is observed with a significance larger than 12σ . Its quantum numbers are determined to be $J^{PC} = 0^{++}$, favored over 1^{--} or 2^{++} with a significance greater than 9σ . As argued above, the $X(3960)$ and $\chi_{c0}(3930)$ states are unlikely to be the same pure conventional charmonium state. The $X(3960)$ resonance presented in this Letter could be a candidate for an exotic state predominantly consisting of $c\bar{c}s\bar{s}$ constituents, as suggested in recent theoretical literature [38–43,65]. If predominant $c\bar{c}s\bar{s}$ content is confirmed, this state should be labeled $T_{\psi\phi}^f(3960)$ in the new naming scheme for exotic hadrons [70]. In addition, a dip around 4140 MeV can be described either by a $0^{++} X_0(4140)$ resonance having a significance of 3.5σ , or the coupled-channel effect of the $J/\psi\phi \leftrightarrow D_s^+ D_s^-$ reaction. The data from the forthcoming Run 3 of the LHCb experiment and from the Belle II experiment will be critical to clarify the nature of these phenomena.

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ, and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG, and MPG (Germany); INFN (Italy); NWO (Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MICINN (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (U.K.); DOE NP and NSF (U.S.). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (Netherlands), PIC (Spain), GridPP (U.K.), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), Polish WLCG (Poland), and NERSC (USA). We are indebted to the communities behind the multiple open-source software packages on which we depend. Individual groups or members have received support from ARC and ARDC (Australia); Minciencias (Colombia); AvH Foundation (Germany); EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union); A*MIDEX, ANR, IPhU and Labex P2IO, and Région Auvergne-Rhône-Alpes (France); Key Research Program of Frontier Sciences of CAS, CAS PIFI, CAS CCEPP, Fundamental Research Funds for the Central Universities, and Sci. & Tech. Program of Guangzhou (China); GVA, XuntaGal, GENCAT, and Prog. Atracción Talento, CM (Spain); SRC (Sweden); the Leverhulme Trust, the Royal Society, and UKRI (U.K.).

- [1] S. K. Choi *et al.* (Belle Collaboration), Observation of a Resonance-Like Structure in the $\pi^\pm\psi'$ Mass Distribution in Exclusive $B \rightarrow K\pi^\pm\psi'$ Decays, *Phys. Rev. Lett.* **100**, 142001 (2008).
- [2] K. Chilikin *et al.* (Belle Collaboration), Experimental constraints on the spin and parity of the $Z(4430)^+$, *Phys. Rev. D* **88**, 074026 (2013).
- [3] A. Bondar *et al.* (Belle Collaboration), Observation of Two Charged Bottomonium-Like Resonances in $\Upsilon(5S)$ Decays, *Phys. Rev. Lett.* **108**, 122001 (2012).
- [4] M. Ablikim *et al.* (BESIII Collaboration), Observation of a Charged Charmoniumlike Structure in $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ at $\sqrt{s} = 4.26$ GeV, *Phys. Rev. Lett.* **110**, 252001 (2013).
- [5] Z. Q. Liu *et al.* (Belle Collaboration), Study of $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ and Observation of a Charged Charmoniumlike State at Belle, *Phys. Rev. Lett.* **110**, 252002 (2013); **111**, 019901(E) (2013).
- [6] T. Xiao, S. Dobbs, A. Tomaradze, and K. K. Seth, Observation of the charged hadron $Z_c^\pm(3900)$ and evidence for the neutral $Z_c^0(3900)$ in $e^+e^- \rightarrow \pi\pi J/\psi$ at $\sqrt{s} = 4170$ MeV, *Phys. Lett. B* **727**, 366 (2013).
- [7] M. Ablikim *et al.* (BESIII Collaboration), Observation of a Charged Charmoniumlike Structure in $e^+e^- \rightarrow (D^*\bar{D}^*)^\pm\pi^\mp$ at $\sqrt{s} = 4.26$ GeV, *Phys. Rev. Lett.* **112**, 132001 (2014).
- [8] M. Ablikim *et al.* (BESIII Collaboration), Observation of a Charged Charmoniumlike Structure $Z_c(4020)$ and Search for the $Z_c(3900)$ in $e^+e^- \rightarrow \pi^+\pi^-h_c$, *Phys. Rev. Lett.* **111**, 242001 (2013).
- [9] R. Aaij *et al.* (LHCb Collaboration), Observation of $J/\psi p$ Resonances Consistent with Pentaquark States in $\Lambda_b^0 \rightarrow J/\psi p K^-$ Decays, *Phys. Rev. Lett.* **115**, 072001 (2015).
- [10] R. Aaij *et al.* (LHCb Collaboration), Observation of a Narrow Pentaquark State, $P_c(4312)^+$, and of Two-Peak Structure of the $P_c(4450)^+$, *Phys. Rev. Lett.* **122**, 222001 (2019).
- [11] M. Ablikim *et al.* (BESIII Collaboration), Observation of a Near-Threshold Structure in the K^+ Recoil-Mass Spectra in $e^+e^- \rightarrow K^+(D_s^-D^{*0} + D_s^{*-}D^0)$, *Phys. Rev. Lett.* **126**, 102001 (2021).
- [12] R. Aaij *et al.* (LHCb Collaboration), Observation of New Resonances Decaying to $J/\psi K^+$ and $J/\psi\phi$, *Phys. Rev. Lett.* **127**, 082001 (2021).
- [13] R. Aaij *et al.* (LHCb Collaboration), Model-Independent Study of Structure in $B^+ \rightarrow D^+D^-K^+$ Decays, *Phys. Rev. Lett.* **125**, 242001 (2020).
- [14] R. Aaij *et al.* (LHCb Collaboration), Amplitude analysis of the $B^+ \rightarrow D^+D^-K^+$ decay, *Phys. Rev. D* **102**, 112003 (2020).
- [15] R. Aaij *et al.* (LHCb Collaboration), Observation of an exotic narrow doubly charmed tetraquark, *Nat. Phys.* **18**, 751 (2022).
- [16] R. Aaij *et al.* (LHCb Collaboration), Study of the doubly charmed tetraquark T_{cc}^+ , *Nat. Commun.* **13**, 3351 (2022).
- [17] Q.-R. Gong, Z.-H. Guo, C. Meng, G.-Y. Tang, Y.-F. Wang, and H.-Q. Zheng, $Z_c(3900)$ as a $D\bar{D}^*$ molecule from the pole counting rule, *Phys. Rev. D* **94**, 114019 (2016).
- [18] Q.-R. Gong, J.-L. Pang, Y.-F. Wang, and H.-Q. Zheng, The $Z_c(3900)$ peak does not come from the “triangle singularity”, *Eur. Phys. J. C* **78**, 276 (2018).
- [19] Q.-F. Cao, H. Chen, H.-R. Qi, and H.-Q. Zheng, Some remarks on $X(6900)$, *Chin. Phys. C* **45**, 103102 (2021).
- [20] H. Chen, H.-R. Qi, and H.-Q. Zheng, $X_1(2900)$ as a \bar{D}_1K molecule, *Eur. Phys. J. C* **81**, 812 (2021).
- [21] L. Maiani, V. Riquer, R. Faccini, F. Piccinini, A. Pilloni, and A. D. Polosa, A $J^{PG} = 1^{++}$ charged resonance in the $Y(4260) \rightarrow \pi^+\pi^-J/\psi$ decay?, *Phys. Rev. D* **87**, 111102(R) (2013).
- [22] J. M. Dias, F. S. Navarra, M. Nielsen, and C. M. Zanetti, $Z_c(3900)^+$ decay width in QCD sum rules, *Phys. Rev. D* **88**, 016004 (2013).
- [23] L. Maiani, F. Piccinini, A. D. Polosa, and V. Riquer, Diquark-antidiquarks with hidden or open charm and the nature of $X(3872)$, *Phys. Rev. D* **71**, 014028 (2005).
- [24] S. Dubynskiy, A. Gorsky, and M. B. Voloshin, Holographic hadro-quarkonium, *Phys. Lett. B* **671**, 82 (2009).
- [25] M. Alberti, G. S. Bali, S. Collins, F. Knechtli, G. Moir, and W. Söldner, Hadroquarkonium from lattice QCD, *Phys. Rev. D* **95**, 074501 (2017).
- [26] D. V. Bugg, An explanation of Belle states $Z_b(10610)$ and $Z_b(10650)$, *Europhys. Lett.* **96**, 11002 (2011).
- [27] F.-K. Guo, X.-H. Liu, and S. Sakai, Threshold cusps and triangle singularities in hadronic reactions, *Prog. Part. Nucl. Phys.* **112**, 103757 (2020).
- [28] K. Abe *et al.* (Belle Collaboration), Observation of a Near-Threshold $\omega J/\psi$ Mass Enhancement in Exclusive $B \rightarrow K\omega J/\psi$ Decays, *Phys. Rev. Lett.* **94**, 182002 (2005).
- [29] P. del Amo Sanchez *et al.* (BABAR Collaboration), Evidence for the decay $X(3872) \rightarrow J/\psi\omega$, *Phys. Rev. D* **82**, 011101 (2010).
- [30] S. Uehara *et al.* (Belle Collaboration), Observation of a Charmonium-Like Enhancement in the $\gamma\gamma \rightarrow \omega J/\psi$ Process, *Phys. Rev. Lett.* **104**, 092001 (2010).
- [31] J. P. Lees *et al.* (BABAR Collaboration), Study of $X(3915) \rightarrow J/\psi\omega$ in two-photon collisions, *Phys. Rev. D* **86**, 072002 (2012).
- [32] P. A. Zyla *et al.* (Particle Data Group), Review of particle physics, *Prog. Theor. Exp. Phys.* **2020**, 083C01 (2020).
- [33] T. Barnes, S. Godfrey, and E. S. Swanson, Higher charmonia, *Phys. Rev. D* **72**, 054026 (2005).
- [34] S. F. Radford and W. W. Repko, Potential model calculations and predictions for heavy quarkonium, *Phys. Rev. D* **75**, 074031 (2007).
- [35] B.-Q. Li and K.-T. Chao, Higher charmonia and X, Y, Z states with screened potential, *Phys. Rev. D* **79**, 094004 (2009).
- [36] H. Wang, Y. Yang, and J. Ping, Strong decays of $\chi_{cJ}(2P)$ and $\chi_{cJ}(3P)$, *Eur. Phys. J. A* **50**, 76 (2014).
- [37] F.-K. Guo and U.-G. Meissner, Where is the $\chi_{c0}(2P)$?, *Phys. Rev. D* **86**, 091501(R) (2012).
- [38] R. F. Lebed and A. D. Polosa, $\chi_{c0}(3915)$ As the lightest $c\bar{c}s\bar{s}$ state, *Phys. Rev. D* **93**, 094024 (2016).
- [39] W. Chen, H.-X. Chen, X. Liu, T. G. Steele, and S.-L. Zhu, Mass spectra for $qc\bar{q}\bar{c}$, $sc\bar{s}\bar{c}$, $qb\bar{q}\bar{b}$, $sb\bar{s}\bar{b}$, tetraquark states with $J^{PC} = 0^{++}$ and 2^{++} , *Phys. Rev. D* **96**, 114017 (2017).
- [40] S. Prelovsek, S. Collins, D. Mohler, M. Padmanath, and S. Piemonte, Charmonium-like resonances with $J^{PC} = 0^{++}, 2^{++}$ in coupled $D\bar{D}, D_s\bar{D}_s$ scattering on the lattice, *J. High Energy Phys.* **06** (2021) 035.

- [41] X. Liu, H. Huang, J. Ping, D. Chen, and X. Zhu, The explanation of some exotic states in the $cs\bar{c}\bar{s}$ tetraquark system, *Eur. Phys. J. C* **81**, 950 (2021).
- [42] L. Meng, B. Wang, and S.-L. Zhu, Predicting the $\bar{D}_s^{(*)}D_s^{(*)}$ bound states as the partners of $X(3872)$, *Sci. Bull.* **66**, 1288 (2021).
- [43] X.-K. Dong, F.-K. Guo, and B.-S. Zou, A survey of heavy-antiheavy hadronic molecules, *Progr. Phys.* **41**, 65 (2021).
- [44] G. Pakhlova *et al.* (Belle Collaboration), Measurement of $e^+e^- \rightarrow D_s^{(*)+}D_s^{(*)-}$ cross sections near threshold using initial-state radiation, *Phys. Rev. D* **83**, 011101 (2011).
- [45] R. Aaij *et al.* (LHCb Collaboration), companion paper, First observation of the $B^+ \rightarrow D_s^+D_s^-K^+$ decay, *Phys. Rev. D* **108**, 034012 (2023).
- [46] Q.-F. Cao, H.-R. Qi, G.-Y. Tang, Y.-F. Xue, and H.-Q. Zheng, On leptonic width of $X(4260)$, *Eur. Phys. J. C* **81**, 83 (2021).
- [47] Q.-F. Cao, H.-R. Qi, Y.-F. Wang, and H.-Q. Zheng, Discussions on the line-shape of the $X(4660)$ resonance, *Phys. Rev. D* **100**, 054040 (2019).
- [48] W. D. Hulsbergen, Decay chain fitting with a Kalman filter, *Nucl. Instrum. Methods Phys. Res., Sect. A* **552**, 566 (2005).
- [49] M. Pivk and F. R. Le Diberder, sPlot: A statistical tool to unfold data distributions, *Nucl. Instrum. Methods Phys. Res., Sect. A* **555**, 356 (2005).
- [50] Y. Xie, *sFit*: A method for background subtraction in maximum likelihood fit, [arXiv:0905.0724](https://arxiv.org/abs/0905.0724).
- [51] S. U. Chung, Spin formalisms, Report No. CERN-71-08, <https://cds.cern.ch/record/186421>; J. D. Richman, An experimenter's guide to the helicity formalism, Report No. CALT-68-1148, <https://lib-extop.c.ksu.edu/preprints/PDF/1984/8409/8409198.pdf>; M. Jacob and G. C. Wick, On the general theory of collisions for particles with spin, *Ann. Phys. (N.Y.)* **7**, 404 (1959).
- [52] R. Aaij *et al.* (LHCb Collaboration), Dalitz plot analysis of $B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$ decays, *Phys. Rev. D* **90**, 072003 (2014).
- [53] R. Aaij *et al.* (LHCb Collaboration), Amplitude analysis of $B^- \rightarrow D^+ \pi^- \pi^-$ decays, *Phys. Rev. D* **94**, 072001 (2016).
- [54] R. Aaij *et al.* (LHCb Collaboration), Amplitude analysis of the $B^+ \rightarrow \pi^+ \pi^+ \pi^-$ decay, *Phys. Rev. D* **101**, 012006 (2020).
- [55] S. M. Flatté, Coupled-channel analysis of the $\pi\eta$ and $K\bar{K}$ systems near $K\bar{K}$ threshold, *Phys. Lett.* **63B**, 224 (1976).
- [56] V. V. Anisovich and A. V. Sarantsev, K -matrix analysis of the $(IJ^{PC} = 00^{++})$ -wave in the mass region below 1900 MeV, *Eur. Phys. J. A* **16**, 229 (2003).
- [57] J. M. Blatt and V. F. Weisskopf, *Theoretical Nuclear Physics* (Springer, New York, 1952), [10.1007/978-1-4612-9959-2](https://doi.org/10.1007/978-1-4612-9959-2).
- [58] F. Von Hippel and C. Quigg, Centrifugal-barrier effects in resonance partial decay widths, shapes, and production amplitudes, *Phys. Rev. D* **5**, 624 (1972).
- [59] S. U. Chung *et al.*, Partial wave analysis in K matrix formalism, *Ann. Phys. (N.Y.)* **4**, 404 (1995).
- [60] M. Tanabashi *et al.* (Particle Data Group), Review of particle physics, *Phys. Rev. D* **98**, 030001 (2018).
- [61] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.131.071901> for a summary of systematic uncertainties, and main results and fit projections from both two-channel Flatté-like parametrization and K -matrix model.
- [62] B. Efron, Bootstrap methods: Another look at the jackknife, *Ann. Statist.* **7**, 1 (1979).
- [63] R. Aaij *et al.* (LHCb Collaboration), Determination of quantum numbers for several excited charmed mesons observed in $B^- \rightarrow D^{*+} \pi^- \pi^-$ decays, *Phys. Rev. D* **101**, 032005 (2020).
- [64] R. Aaij *et al.* (LHCb Collaboration), First observation and amplitude analysis of the $B^- \rightarrow D^+ K^- \pi^-$ decay, *Phys. Rev. D* **91**, 092002 (2015); **93**, 119901(E) (2016).
- [65] E. Wang, W.-H. Liang, and E. Oset, Analysis of the $e^+e^- \rightarrow J/\psi D\bar{D}$ reaction close to the threshold concerning claims of a $\chi_{c0}(2P)$ state, *Eur. Phys. J. A* **57**, 38 (2021).
- [66] S. Okubo, Φ meson and unitary symmetry model, *Phys. Lett.* **5**, 165 (1963).
- [67] J. Iizuka, Systematics and phenomenology of meson family, *Prog. Theor. Phys. Suppl.* **37**, 21 (1966).
- [68] I. J. R. Aitchison, The K -matrix formalism for overlapping resonances, *Nucl. Phys.* **A189**, 417 (1972).
- [69] I. J. R. Aitchison, Unitarity, analyticity and crossing symmetry in two- and three-hadron final state interactions, [arXiv:1507.02697](https://arxiv.org/abs/1507.02697).
- [70] T. Gershon (LHCb Collaboration), Exotic hadron naming convention, Report No. LHCb-PUB-2022-013, 2022.

R. Aaij³², A. S. W. Abdelmotteleb⁵⁰, C. Abellan Beteta⁴⁴, F. Abudinén⁵⁰, T. Ackernley⁵⁴, B. Adeva⁴⁰, M. Adinolfi⁴⁸, H. Afsharnia⁹, C. Agapopoulou¹³, C. A. Aidala⁷⁷, S. Aiola²⁵, Z. Ajaltouni⁹, S. Akar⁵⁹, K. Akiba³², J. Albrecht¹⁵, F. Alessio⁴², M. Alexander⁵³, A. Alfonso Albergo³⁹, Z. Aliouche⁵⁶, P. Alvarez Cartelle⁴⁹, R. Amalric¹³, S. Amato², J. L. Amey⁴⁸, Y. Amhis^{11,42}, L. An⁴², L. Anderlini²², M. Andersson⁴⁴, A. Andreianov³⁸, M. Andreotti²¹, D. Andreou⁶², D. Ao⁶, F. Archilli¹⁷, A. Artamonov³⁸, M. Artuso⁶², E. Aslanides¹⁰, M. Atzeni⁴⁴, B. Audurier¹², S. Bachmann¹⁷, M. Bachmayer⁴³, J. J. Back⁵⁰, A. Bailly-reyre¹³, P. Baladron Rodriguez⁴⁰, V. Balagura¹², W. Baldini²¹, J. Baptista de Souza Leite¹, M. Barbetti^{22,b}, R. J. Barlow⁵⁶, S. Barsuk¹¹, W. Barter⁵⁵, M. Bartolini⁴⁹, F. Baryshnikov³⁸, J. M. Basels¹⁴, G. Bassi^{29,c}, B. Batsukh⁴, A. Battig¹⁵, A. Bay⁴³, A. Beck⁵⁰, M. Becker¹⁵, F. Bedeschi²⁹, I. B. Bediaga¹, A. Beiter⁶², V. Belavin³⁸, S. Belin⁴⁰, V. Bellee⁴⁴, K. Belous³⁸, I. Belov³⁸, I. Belyaev³⁸, G. Benane¹⁰, G. Bencivenni²³, E. Ben-Haim¹³, A. Bereznoy³⁸, R. Bernet⁴⁴, S. Bernet Andres⁷⁵, D. Berninghoff¹⁷, H. C. Bernstein⁶², C. Bertella⁵⁶, A. Bertolin²⁸

C. Betancourt⁴⁴ F. Betti⁴² Ia. Bezshyiko⁴⁴ S. Bhasin⁴⁸ J. Bhom³⁵ L. Bian⁶⁸ M. S. Bieker¹⁵ N. V. Biesuz²¹ S. Bifani⁴⁷ P. Billoir¹³ A. Biolchini³² M. Birch⁵⁵ F. C. R. Bishop⁴⁹ A. Bitadze⁵⁶ A. Bizzeti¹² M. P. Blago⁴⁹ T. Blake⁵⁰ F. Blanc⁴³ S. Blusk⁶² D. Bobulska^{21,d} J. A. Boelhauve⁵³ O. Boente Garcia¹² T. Boettcher⁵⁹ A. Boldyrev³⁸ C. S. Bolognani⁷⁴ R. Bolzonella^{21,d} N. Bondar^{38,42} F. Borgato²⁸ S. Borghi⁵⁶ M. Borsato¹⁷ J. T. Borsuk³⁵ S. A. Bouchiba⁴³ T. J. V. Bowcock⁵⁴ A. Boyer⁴² C. Bozzi²¹ M. J. Bradley⁵⁵ S. Braun⁶⁰ A. Brea Rodriguez⁴⁰ J. Brodzicka³⁵ A. Brossa Gonzalo⁴⁰ J. Brown⁵⁴ D. Brundu²⁷ A. Buonauro⁴⁴ L. Buonincontri²⁸ A. T. Burke⁵⁶ C. Burr⁴² A. Bursche⁶⁶ A. Butkevich³⁸ J. S. Butter³² J. Buytaert⁴² W. Byczynski⁴² S. Cadeddu²⁷ H. Cai⁶⁸ R. Calabrese^{21,d} L. Calefice¹⁵ S. Cali²³ R. Calladine⁴⁷ M. Calvi^{26,e} M. Calvo Gomez⁷⁵ P. Campana²³ D. H. Campora Perez⁷⁴ A. F. Campoverde Quezada⁶ S. Capelli^{26,e} L. Capriotti^{20,f} A. Carbone^{20,f} G. Carboni³¹ R. Cardinale^{24,g} A. Cardini²⁷ I. Carli⁴ P. Carniti^{26,e} L. Carus¹⁴ A. Casais Vidal⁴⁰ R. Caspary¹⁷ G. Casse⁵⁴ M. Cattaneo⁴² G. Cavallero⁴² V. Cavallini^{21,d} S. Celani⁴³ J. Cerasoli¹⁰ D. Cervenkov⁵⁷ A. J. Chadwick⁵⁴ M. G. Chapman⁴⁸ M. Charles¹³ Ph. Charpentier⁴² C. A. Chavez Barajas⁵⁴ M. Chefdeville⁸ C. Chen³ S. Chen⁴ A. Chernov³⁵ S. Chernyshenko⁴⁶ V. Chobanova⁴⁰ S. Cholak⁴³ M. Chrzaszcz³⁵ A. Chubykin³⁸ V. Chulikov³⁸ P. Ciambrone²³ M. F. Cicala⁵⁰ X. Cid Vidal⁴⁰ G. Ciezarek⁴² G. Ciullo^{21,d} P. E. L. Clarke⁵² M. Clemencic⁴² H. V. Cliff⁴⁹ J. Closier⁴² J. L. Cobbedick⁵⁶ V. Coco⁴² J. A. B. Coelho¹¹ J. Cogan¹⁰ E. Cogneras⁹ L. Cojocariu³⁷ P. Collins⁴² T. Colombo⁴² L. Congedo¹⁹ A. Contu²⁷ N. Cooke⁴⁷ I. Corredoira⁴⁰ G. Corti⁴² B. Couturier⁴² D. C. Craik⁵⁸ J. Crkovská⁶¹ M. Cruz Torres^{1,h} R. Currie⁵² C. L. Da Silva⁶¹ S. Dadabaev³⁸ L. Dai⁶⁵ X. Dai⁵ E. Dall'Occo¹⁵ J. Dalseno⁴⁰ C. D'Ambrosio⁴² J. Daniel⁹ A. Danilina³⁸ P. d'Argent¹⁵ J. E. Davies⁵⁶ A. Davis⁵⁶ O. De Aguiar Francisco⁵⁶ J. de Boer⁴² K. De Bruyn⁷³ S. De Capua⁵⁶ M. De Cian⁴³ U. De Freitas Carneiro Da Graca¹ E. De Lucia²³ J. M. De Miranda¹ L. De Paula² M. De Serio^{19,i} D. De Simone⁴⁴ P. De Simone²³ F. De Vellis¹⁵ J. A. de Vries⁷⁴ C. T. Dean⁶¹ F. Debernardis^{19,i} D. Decamp⁸ V. Dedu¹⁰ L. Del Buono¹³ B. Delaney⁵⁸ H.-P. Dembinski¹⁵ V. Denysenko⁴⁴ O. Deschamps⁹ F. Dettori^{27,j} B. Dey⁷¹ A. Di Cicco²³ P. Di Nezza²³ I. Diachkov³⁸ S. Didenko³⁸ L. Dieste Maronas⁴⁰ S. Ding⁶² V. Dobishuk⁴⁶ A. Dolmatov³⁸ C. Dong³ A. M. Donohoe¹⁸ F. Dordei²⁷ A. C. dos Reis¹ L. Douglas⁵³ A. G. Downes⁸ M. W. Dudek³⁵ L. Dufour⁴² V. Duk⁷² P. Durante⁴² J. M. Durham⁶¹ D. Dutta⁵⁶ A. Dziurda³⁵ A. Dzyuba³⁸ S. Easo⁵¹ U. Egede⁶³ V. Egorychev³⁸ S. Eidelman^{38,a} C. Eirea Orro⁴⁰ S. Eisenhardt⁵² E. Ejopu⁵⁶ S. Ek-In⁴³ L. Eklund⁷⁶ S. Ely⁶² A. Ene³⁷ E. Epple⁶¹ S. Escher¹⁴ J. Eschle⁴⁴ S. Esen⁴⁴ T. Evans⁵⁶ F. Fabiano^{27,j} L. N. Falcao¹ Y. Fan⁶ B. Fang⁶⁸ S. Farry⁵⁴ D. Fazzini^{26,e} M. Feo⁴² M. Fernandez Gomez⁴⁰ A. D. Fernandez⁶⁰ F. Ferrari²⁰ L. Ferreira Lopes⁴³ F. Ferreira Rodrigues² S. Ferreres Sole³² M. Ferrillo⁴⁴ M. Ferro-Luzzi⁴² S. Filippov³⁸ R. A. Fini¹⁹ M. Fiorini^{21,d} M. Firlej³⁴ K. M. Fischer⁵⁷ D. S. Fitzgerald⁷⁷ C. Fitzpatrick⁵⁶ T. Fiutowski³⁴ F. Fleuret¹² M. Fontana¹³ F. Fontanelli^{24,g} R. Forty⁴² D. Foulds-Holt⁴⁹ V. Franco Lima⁵⁴ M. Franco Sevilla⁶⁰ M. Frank⁴² E. Franzoso^{21,d} G. Frau¹⁷ C. Frei⁴² D. A. Friday⁵³ J. Fu⁶ Q. Fuehring¹⁵ T. Fulghesu¹³ E. Gabriel³² G. Galati^{19,i} M. D. Galati⁷³ A. Gallas Torreira⁴⁰ D. Galli^{20,f} S. Gambetta^{52,42} Y. Gan³ M. Gandelman² P. Gandini²⁵ Y. Gao⁵ M. Garau^{27,j} L. M. Garcia Martin⁵⁰ P. Garcia Moreno³⁹ J. García Pardiñas^{26,e} B. Garcia Plana⁴⁰ F. A. Garcia Rosales¹² L. Garrido³⁹ C. Gaspar⁴² R. E. Geertsema³² D. Gerick¹⁷ L. L. Gerken¹⁵ E. Gersabeck⁵⁶ M. Gersabeck⁵⁶ T. Gershon⁵⁰ L. Giambastiani²⁸ V. Gibson⁴⁹ H. K. Giemza³⁶ A. L. Gilman⁵⁷ M. Giovannetti^{23,k} A. Gioventù⁴⁰ P. Gironella Gironell³⁹ C. Giugliano^{21,d} M. A. Giza³⁵ K. Gizdov⁵² E. L. Gkougkousis⁴² V. V. Gligorov^{13,42} C. Göbel⁶⁴ E. Golobardes⁷⁵ D. Golubkov³⁸ A. Golutvin^{55,38} A. Gomes^{1,l} S. Gomez Fernandez³⁹ F. Goncalves Abrantes⁵⁷ M. Goncerz³⁵ G. Gong³ I. V. Gorelov³⁸ C. Gotti²⁶ J. P. Grabowski¹⁷ T. Grammatico¹³ L. A. Granado Cardoso⁴² E. Graugés³⁹ E. Graverini⁴³ G. Graziani¹ A. T. Grecu³⁷ L. M. Greeven³² N. A. Grieser⁴ L. Grillo⁵³ S. Gromov³⁸ B. R. Gruberg Cazon⁵⁷ C. Gu³ M. Guarise^{21,d} M. Guittiere¹¹ P. A. Günther¹⁷ E. Gushchin³⁸ A. Guth¹⁴ Y. Guz³⁸ T. Gys⁴² T. Hadavizadeh⁶³ G. Haefeli⁴³ C. Haen⁴² J. Haimberger⁴² S. C. Haines⁴⁹ T. Halewood-leagas⁵⁴ M. M. Halvorsen⁴² P. M. Hamilton⁶⁰ J. Hammerich⁵⁴ Q. Han⁷ X. Han¹⁷ E. B. Hansen⁵⁶ S. Hansmann-Menzemer^{17,42} L. Hao⁶ N. Harnew⁵⁷ T. Harrison⁵⁴ C. Hasse⁴² M. Hatch⁴² J. He^{6,m} K. Heijhoff³² K. Heinicke¹⁵ C. Henderson⁵⁹ R. D. L. Henderson^{63,50} A. M. Hennequin⁵⁸ K. Hennessy⁵⁴ L. Henry⁴² J. H Herd⁵⁵ J. Heuel¹⁴ A. Hicheur² D. Hill⁴³ M. Hilton⁵⁶ S. E. Hollitt¹⁵

J. Horswill⁵⁶, R. Hou⁷, Y. Hou⁸, J. Hu¹⁷, J. Hu⁶⁶, W. Hu⁵, X. Hu³, W. Huang⁶, X. Huang⁶⁸, W. Hulsbergen³², R. J. Hunter⁵⁰, M. Hushchyn³⁸, D. Hutchcroft⁵⁴, P. Ibis¹⁵, M. Idzik³⁴, D. Ilin³⁸, P. Ilten⁵⁹, A. Inglessi³⁸, A. Iniukhin³⁸, A. Ishteev³⁸, K. Ivshin³⁸, R. Jacobsson⁴², H. Jage¹⁴, S. J. Jaimes Elles⁴¹, S. Jakobsen⁴², E. Jans³², B. K. Jashal⁴¹, A. Jawahery⁶⁰, V. Jevtic¹⁵, E. Jiang⁶⁰, X. Jiang^{4,6}, Y. Jiang⁶, M. John⁵⁷, D. Johnson⁵⁸, C. R. Jones⁴⁹, T. P. Jones⁵⁰, B. Jost⁴², N. Jurik⁴², I. Juszczak³⁵, S. Kandybei⁴⁵, Y. Kang³, M. Karacson⁴², D. Karpenkov³⁸, M. Karpov³⁸, J. W. Kautz⁵⁹, F. Keizer⁴², D. M. Keller⁶², M. Kenzie⁵⁰, T. Ketel³², B. Khanji¹⁵, A. Kharisova³⁸, S. Kholodenko³⁸, G. Khreich¹¹, T. Kim¹⁴, V. S. Kirsebom⁴³, O. Kitouni⁵⁸, S. Klaver³³, N. Kleijne^{29,c}, K. Klimaszewski³⁶, M. R. Kmiec³⁶, S. Koliiev⁴⁶, A. Kondybayeva³⁸, A. Konoplyannikov³⁸, P. Kopciwicz³⁴, R. Kopečna¹⁷, P. Koppenburg³², M. Korolev³⁸, I. Kostiuk^{32,46}, O. Kot⁴⁶, S. Kotriakhova⁴⁶, A. Kozachuk³⁸, P. Kravchenko³⁸, L. Kravchuk³⁸, R. D. Krawczyk⁴², M. Kreps⁵⁰, S. Kretzschmar¹⁴, P. Krokovny³⁸, W. Krupa³⁴, W. Krzemien³⁶, J. Kubat¹⁷, W. Kucewicz^{35,34}, M. Kucharczyk³⁵, V. Kudryavtsev³⁸, G. J. Kunde⁶¹, A. Kupsc⁷⁶, D. Lacarrere⁴², G. Lafferty⁵⁶, A. Lai²⁷, A. Lampis^{27,j}, D. Lancierini⁴⁴, C. Landesa Gomez⁴⁰, J. J. Lane⁵⁶, R. Lane⁴⁸, G. Lanfranchi²³, C. Langenbruch¹⁴, J. Langer¹⁵, O. Lantwin³⁸, T. Latham⁵⁰, F. Lazzari^{29,n}, M. Lazzaroni²⁵, R. Le Gac¹⁰, S. H. Lee⁷⁷, R. Lefèvre⁹, A. Leflat³⁸, S. Legotin³⁸, P. Lenisa^{21,d}, O. Leroy¹⁰, T. Lesiak³⁵, B. Leverington¹⁷, A. Li³, H. Li⁶⁶, K. Li⁷, P. Li¹⁷, P.-R. Li⁶⁷, S. Li⁷, T. Li⁶⁶, Y. Li⁴, Z. Li⁶², X. Liang⁶², C. Lin⁶, T. Lin⁵¹, R. Lindner⁴², V. Lisovskyi¹⁵, R. Litvinov^{27,j}, G. Liu⁶⁶, H. Liu⁶, Q. Liu⁶, S. Liu^{4,6}, A. Lobo Salvia³⁹, A. Loi²⁷, R. Lollini⁷², J. Lomba Castro⁴⁰, I. Longstaff⁵³, J. H. Lopes², A. Lopez Huertas³⁹, S. López Soliño⁴⁰, G. H. Lovell⁴⁹, Y. Lu^{4,0}, C. Lucarelli^{22,b}, D. Lucchesi^{28,p}, S. Luchuk³⁸, M. Lucio Martinez⁷⁴, V. Lukashenko^{32,46}, Y. Luo³, A. Lupato⁵⁶, E. Luppi^{21,d}, A. Lusiani^{29,c}, K. Lynch¹⁸, X.-R. Lyu⁶, L. Ma⁴, R. Ma⁶, S. Maccolini²⁰, F. Machefert¹¹, F. Maciuc³⁷, I. Mackay⁵⁷, V. Macko⁴³, P. Mackowiak¹⁵, L. R. Madhan Mohan⁴⁸, A. Maevskiy³⁸, D. Maisuzenko³⁸, M. W. Majewski³⁴, J. J. Malczewski³⁵, S. Malde⁵⁷, B. Malecki^{35,42}, A. Malinin³⁸, T. Maltsev³⁸, G. Manca^{27,j}, G. Mancinelli¹⁰, C. Mancuso^{11,25,q}, D. Manuzzi²⁰, C. A. Manzari⁴⁴, D. Marangotto^{25,q}, J. M. Maratas^{9,r}, J. F. Marchand⁸, U. Marconi²⁰, S. Mariani^{22,b}, C. Marin Benito³⁹, J. Marks¹⁷, A. M. Marshall⁴⁸, P. J. Marshall⁵⁴, G. Martelli^{72,s}, G. Martellotti³⁰, L. Martinazzoli^{42,e}, M. Martinelli^{26,e}, D. Martinez Santos⁴⁰, F. Martinez Vidal⁴¹, A. Massafferri¹, M. Materok¹⁴, R. Matev⁴², A. Mathad⁴⁴, V. Matiunin³⁸, C. Matteuzzi²⁶, K. R. Mattioli⁷⁷, A. Mauri³², E. Maurice¹², J. Mauricio³⁹, M. Mazurek⁴², M. McCann⁵⁵, L. McConnell¹⁸, T. H. McGrath⁵⁶, N. T. McHugh⁵³, A. McNab⁵⁶, R. McNulty¹⁸, J. V. Mead⁵⁴, B. Meadows⁵⁹, G. Meier¹⁵, D. Melnychuk³⁶, S. Meloni^{26,e}, M. Merk^{32,74}, A. Merli^{25,q}, L. Meyer Garcia², D. Miao^{4,6}, M. Mikhasenko^{70,t}, D. A. Milanese⁶⁹, E. Millard⁵⁰, M. Milovanovic⁴², M.-N. Minard^{8,8,a}, A. Minotti^{26,e}, T. Miralles⁹, S. E. Mitchell⁵², B. Mitreska⁵⁶, D. S. Mitzel¹⁵, A. Mödden¹⁵, R. A. Mohammed⁵⁷, R. D. Moise¹⁴, S. Mokhnenko³⁸, T. Mombächer⁴⁰, M. Monk^{50,63}, I. A. Monroy⁶⁹, S. Monteil⁹, M. Morandin²⁸, G. Morello²³, M. J. Morello^{29,c}, J. Moron³⁴, A. B. Morris⁷⁰, A. G. Morris⁵⁰, R. Mountain⁶², H. Mu³, E. Muhammad⁵⁰, F. Muheim⁵², M. Mulder⁷³, K. Müller⁴⁴, C. H. Murphy⁵⁷, D. Murray⁵⁶, R. Murta⁵⁵, P. Muzzetto^{27,j}, P. Naik⁴⁸, T. Nakada⁴³, R. Nandakumar⁵¹, T. Nanut⁴², I. Nasteva², M. Needham⁵², N. Neri^{25,q}, S. Neubert⁷⁰, N. Neufeld⁴², P. Neustroev³⁸, R. Newcombe⁵⁵, J. Nicolini^{15,11}, E. M. Niel⁴³, S. Nieswand¹⁴, N. Nikitin³⁸, N. S. Nolte⁵⁸, C. Normand^{8,27,j}, J. Novoa Fernandez⁴⁰, C. Nunez⁷⁷, A. Oblakowska-Mucha³⁴, V. Obraztsov³⁸, T. Oeser¹⁴, D. P. O'Hanlon⁴⁸, S. Okamura^{21,d}, R. Oldeman^{27,j}, F. Oliva⁵², M. E. Olivares⁶², C. J. G. Onderwater⁷³, R. H. O'Neil⁵², J. M. Otalora Goicochea², T. Ovsianikova³⁸, P. Owen⁴⁴, A. Oyanguren⁴¹, O. Ozelik⁵², K. O. Padeken⁷⁰, B. Pagare⁵⁰, P. R. Pais⁴², T. Pajero⁵⁷, A. Palano¹⁹, M. Palutan²³, Y. Pan⁵⁶, G. Panshin³⁸, L. Paolucci⁵⁰, A. Papanestis⁵¹, M. Pappagallo^{19,i}, L. L. Pappalardo^{21,d}, C. Pappenheimer⁵⁹, W. Parker⁶⁰, C. Parkes⁵⁶, B. Passalacqua^{21,d}, G. Passaleva²², A. Pastore¹⁹, M. Patel⁵⁵, C. Patrignani^{20,f}, C. J. Pawley⁷⁴, A. Pearce⁴², A. Pellegrino³², M. Pepe Altarelli⁴², S. Perazzini²⁰, D. Pereima³⁸, A. Pereiro Castro⁴⁰, P. Perret⁹, M. Petric⁵³, K. Petridis⁴⁸, A. Petrolini^{24,g}, A. Petrov³⁸, S. Petrucci⁵², M. Petruzzo²⁵, H. Pham⁶², A. Philippov³⁸, R. Piandani⁶, L. Pica^{29,c}, M. Piccini⁷², B. Pietrzyk⁸, G. Pietrzyk¹¹, M. Pili⁵⁷, D. Pinci³⁰, F. Pisani⁴², M. Pizzichemi^{26,42,e}, V. Placinta³⁷, J. Plews⁴⁷, M. Plo Casasus⁴⁰, F. Polci^{13,42}, M. Poli Lener²³, M. Poliakov⁶², A. Poluektov¹⁰, N. Polukhina³⁸, I. Polyakov⁴², E. Polycarpo², S. Ponce⁴², D. Popov^{6,42}, S. Popov³⁸, S. Poslavskii³⁸, K. Prasanth³⁵, L. Promberger⁴², C. Prouve⁴⁰, V. Pugatch⁴⁶, V. Puill¹¹, G. Punzi^{29,u}, H. R. Qi³, W. Qian⁶, N. Qin³, S. Qu³, R. Quagliani⁴³, N. V. Raab¹⁸, R. I. Rabadan Trejo⁶, B. Rachwal³⁴

J. H. Rademacker⁴⁸, R. Rajagopalan,⁶² M. Rama²⁹, M. Ramos Pernas⁵⁰, M. S. Rangel², F. Ratnikov³⁸, G. Raven^{33,42}, M. Rebollo De Miguel⁴¹, F. Redi⁴², J. Reich⁴⁸, F. Reiss⁵⁶, C. Remon Alepuz,⁴¹ Z. Ren³, V. Renaudin⁵⁷, P. K. Resmi¹⁰, R. Ribatti^{29,c}, A. M. Ricci²⁷, S. Ricciardi⁵¹, K. Richardson⁵⁸, M. Richardson-Slipper⁵², K. Rinnert⁵⁴, P. Robbe¹¹, G. Robertson⁵², A. B. Rodrigues⁴³, E. Rodrigues⁵⁴, E. Rodriguez Fernandez⁴⁰, J. A. Rodriguez Lopez⁶⁹, E. Rodriguez Rodriguez⁴⁰, A. Rollings⁵⁷, P. Roloff⁴², V. Romanovskiy³⁸, M. Romero Lamas⁴⁰, A. Romero Vidal⁴⁰, J. D. Roth,^{77,a} M. Rotondo²³, M. S. Rudolph⁶², T. Ruf⁴², R. A. Ruiz Fernandez⁴⁰, J. Ruiz Vidal,⁴¹ A. Ryzhikov³⁸, J. Ryzka³⁴, J. J. Saborido Silva⁴⁰, N. Sagidova³⁸, N. Sahoo⁴⁷, B. Saitta^{27,j}, M. Salomoni⁴², C. Sanchez Gras³², I. Sanderswood⁴¹, R. Santacesaria³⁰, C. Santamarina Rios⁴⁰, M. Santimaria²³, E. Santovetti^{31,k}, D. Saranin³⁸, G. Sarpis¹⁴, M. Sarpis⁷⁰, A. Sarti³⁰, C. Satriano^{30,v}, A. Satta³¹, M. Saur¹⁵, D. Savrina³⁸, H. Sazak⁹, L. G. Scantlebury Smead⁵⁷, A. Scarabotto¹³, S. Schael¹⁴, S. Scherl⁵⁴, M. Schiller⁵³, H. Schindler⁴², M. Schmelling¹⁶, B. Schmidt⁴², S. Schmitt¹⁴, O. Schneider⁴³, A. Schopper⁴², M. Schubiger³², S. Schulte⁴³, M. H. Schune¹¹, R. Schwemmer⁴², B. Sciascia^{23,42}, A. Sciucati⁴², S. Sellam⁴⁰, A. Semennikov³⁸, M. Senghi Soares³³, A. Sergi^{24,g}, N. Serra⁴⁴, L. Sestini²⁸, A. Seuthe¹⁵, Y. Shang⁵, D. M. Shangase⁷⁷, M. Shapkin³⁸, I. Shchemerov³⁸, L. Shchutska⁴³, T. Shears⁵⁴, L. Shekhtman³⁸, Z. Shen⁵, S. Sheng^{4,6}, V. Shevchenko³⁸, B. Shi⁶, E. B. Shields^{26,e}, Y. Shimizu¹¹, E. Shmanin³⁸, J. D. Shupperd⁶², B. G. Siddi^{21,d}, R. Silva Coutinho⁴⁴, G. Simi²⁸, S. Simone^{19,i}, M. Singla⁶³, N. Skidmore⁵⁶, R. Skuza¹⁷, T. Skwarnicki⁶², M. W. Slater⁴⁷, J. C. Smallwood⁵⁷, J. G. Smeaton⁴⁹, E. Smith⁴⁴, K. Smith⁶¹, M. Smith⁵⁵, A. Snoch³², L. Soares Lavra⁹, M. D. Sokoloff⁵⁹, F. J. P. Soler⁵³, A. Solomin^{38,48}, A. Solovev³⁸, I. Solovyevev³⁸, R. Song⁶³, F. L. Souza De Almeida², B. Souza De Paula², B. Spaan,^{15,a} E. Spadaro Norella^{25,q}, E. Spiridenkov,³⁸ P. Spradlin⁵³, V. Sriskaran⁴², F. Stagni⁴², M. Stahl⁵⁹, S. Stahl⁴², S. Stanislaus⁵⁷, E. N. Stein⁴², O. Steinkamp⁴⁴, O. Stenyakin,³⁸ H. Stevens¹⁵, S. Stone^{62,a}, D. Strelakina³⁸, F. Suljik⁵⁷, J. Sun²⁷, L. Sun⁶⁸, Y. Sun⁶⁰, P. Svihra⁵⁶, P. N. Swallow⁴⁷, K. Swientek³⁴, A. Szabelski³⁶, T. Szumlak³⁴, M. Szymanski⁴², Y. Tan³, S. Taneja⁵⁶, A. R. Tanner,⁴⁸ M. D. Tat⁵⁷, A. Terentev³⁸, F. Teubert⁴², E. Thomas⁴², D. J. D. Thompson⁴⁷, K. A. Thomson⁵⁴, H. Tilquin⁵⁵, V. Tisserand⁹, S. T'Jampens⁸, M. Tobin⁴, L. Tomassetti^{21,d}, G. Tonani^{25,q}, X. Tong⁵, D. Torres Machado¹, D. Y. Tou³, E. Trifonova,³⁸ S. M. Trilov⁴⁸, C. Trippel⁴³, G. Tuci⁶, A. Tully⁴³, N. Tuning³², A. Ukleja³⁶, D. J. Unverzagt¹⁷, E. Ursov³⁸, A. Usachov³², A. Ustyuzhanin³⁸, U. Uwer¹⁷, A. Vagner³⁸, V. Vagnoni²⁰, A. Valassi⁴², G. Valenti²⁰, N. Valls Canudas⁷⁵, M. van Beuzekom³², M. Van Dijk⁴³, H. Van Hecke⁶¹, E. van Herwijnen³⁸, C. B. Van Hulse^{40,w}, M. van Veghel⁷³, R. Vazquez Gomez³⁹, P. Vazquez Regueiro⁴⁰, C. Vázquez Sierra⁴², S. Vecchi²¹, J. J. Velthuis⁴⁸, M. Veltri^{22,x}, A. Venkateswaran⁴³, M. Veronesi³², M. Vesterinen⁵⁰, D. Vieira⁵⁹, M. Vieites Diaz⁴³, X. Vilasis-Cardona⁷⁵, E. Vilella Figueras⁵⁴, A. Villa²⁰, P. Vincent¹³, F. C. Volle¹¹, D. vom Bruch¹⁰, A. Vorobyev,³⁸ V. Vorobyev,³⁸ N. Voropaev³⁸, K. Vos⁷⁴, C. Vrahas⁵², R. Waldi¹⁷, J. Walsh²⁹, G. Wan⁵, C. Wang¹⁷, J. Wang⁵, J. Wang⁴, J. Wang³, J. Wang⁶⁸, M. Wang⁵, R. Wang⁴⁸, X. Wang⁶⁶, Y. Wang⁷, Z. Wang⁴⁴, Z. Wang³, Z. Wang⁶, J. A. Ward^{50,63}, N. K. Watson⁴⁷, D. Websdale⁵⁵, Y. Wei⁵, C. Weisser,⁵⁸ B. D. C. Westhenry⁴⁸, D. J. White⁵⁶, M. Whitehead⁵³, A. R. Wiederhold⁵⁰, D. Wiedner¹⁵, G. Wilkinson⁵⁷, M. K. Wilkinson⁵⁹, I. Williams,⁴⁹ M. Williams⁵⁸, M. R. J. Williams⁵², R. Williams⁴⁹, F. F. Wilson⁵¹, W. Wislicki³⁶, M. Witek³⁵, L. Witola¹⁷, C. P. Wong⁶¹, G. Wormser¹¹, S. A. Wotton⁴⁹, H. Wu⁶², K. Wyllie⁴², Z. Xiang⁶, D. Xiao⁷, Y. Xie⁷, A. Xu⁵, J. Xu⁶, L. Xu³, L. Xu³, M. Xu⁵⁰, Q. Xu⁶, Z. Xu⁹, Z. Xu⁶, D. Yang³, S. Yang⁶, Y. Yang⁶, Z. Yang⁵, Z. Yang⁶⁰, L. E. Yeomans⁵⁴, V. Yeroshenko¹¹, H. Yeung⁵⁶, H. Yin⁷, J. Yu⁶⁵, X. Yuan⁶², E. Zaffaroni⁴³, M. Zavertyaev¹⁶, M. Zdybal³⁵, O. Zenaiev⁴², M. Zeng³, C. Zhang⁵, D. Zhang⁷, L. Zhang³, S. Zhang⁶⁵, S. Zhang⁵, Y. Zhang⁵, Y. Zhang⁵⁷, A. Zharkova³⁸, A. Zhelezov¹⁷, Y. Zheng⁶, T. Zhou⁵, X. Zhou⁶, Y. Zhou⁶, V. Zhovkovska¹¹, X. Zhu³, X. Zhu⁷, Z. Zhu⁶, V. Zhukov^{14,38}, Q. Zou^{4,6}, S. Zucchelli^{20,f}, D. Zuliani²⁸, and G. Zunica⁵⁶

(LHCb Collaboration)

¹Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil²Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil³Center for High Energy Physics, Tsinghua University, Beijing, China

- ⁴*Institute Of High Energy Physics (IHEP), Beijing, China*
- ⁵*School of Physics State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China*
- ⁶*University of Chinese Academy of Sciences, Beijing, China*
- ⁷*Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China*
- ⁸*Université Savoie Mont Blanc, CNRS, IN2P3-LAPP, Annecy, France*
- ⁹*Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France*
- ¹⁰*Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France*
- ¹¹*Université Paris-Saclay, CNRS/IN2P3, IJCLab, Orsay, France*
- ¹²*Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France*
- ¹³*LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France*
- ¹⁴*I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany*
- ¹⁵*Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany*
- ¹⁶*Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany*
- ¹⁷*Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany*
- ¹⁸*School of Physics, University College Dublin, Dublin, Ireland*
- ¹⁹*INFN Sezione di Bari, Bari, Italy*
- ²⁰*INFN Sezione di Bologna, Bologna, Italy*
- ²¹*INFN Sezione di Ferrara, Ferrara, Italy*
- ²²*INFN Sezione di Firenze, Firenze, Italy*
- ²³*INFN Laboratori Nazionali di Frascati, Frascati, Italy*
- ²⁴*INFN Sezione di Genova, Genova, Italy*
- ²⁵*INFN Sezione di Milano, Milano, Italy*
- ²⁶*INFN Sezione di Milano-Bicocca, Milano, Italy*
- ²⁷*INFN Sezione di Cagliari, Monserrato, Italy*
- ²⁸*Università degli Studi di Padova, Università e INFN, Padova, Padova, Italy*
- ²⁹*INFN Sezione di Pisa, Pisa, Italy*
- ³⁰*INFN Sezione di Roma La Sapienza, Roma, Italy*
- ³¹*INFN Sezione di Roma Tor Vergata, Roma, Italy*
- ³²*Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands*
- ³³*Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, Netherlands*
- ³⁴*AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland*
- ³⁵*Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*
- ³⁶*National Center for Nuclear Research (NCBJ), Warsaw, Poland*
- ³⁷*Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*
- ³⁸*Affiliated with an institute covered by a cooperation agreement with CERN*
- ³⁹*ICCUB, Universitat de Barcelona, Barcelona, Spain*
- ⁴⁰*Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain*
- ⁴¹*Instituto de Física Corpuscular, Centro Mixto Universidad de Valencia - CSIC, Valencia, Spain*
- ⁴²*European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- ⁴³*Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- ⁴⁴*Physik-Institut, Universität Zürich, Zürich, Switzerland*
- ⁴⁵*NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- ⁴⁶*Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- ⁴⁷*University of Birmingham, Birmingham, United Kingdom*
- ⁴⁸*H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- ⁴⁹*Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- ⁵⁰*Department of Physics, University of Warwick, Coventry, United Kingdom*
- ⁵¹*STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- ⁵²*School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- ⁵³*School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- ⁵⁴*Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- ⁵⁵*Imperial College London, London, United Kingdom*
- ⁵⁶*Department of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- ⁵⁷*Department of Physics, University of Oxford, Oxford, United Kingdom*
- ⁵⁸*Massachusetts Institute of Technology, Cambridge, Massachusetts, USA*
- ⁵⁹*University of Cincinnati, Cincinnati, Ohio, USA*
- ⁶⁰*University of Maryland, College Park, Maryland, USA*
- ⁶¹*Los Alamos National Laboratory (LANL), Los Alamos, New Mexico, USA*
- ⁶²*Syracuse University, Syracuse, New York, USA*

- ⁶³*School of Physics and Astronomy, Monash University, Melbourne, Australia
(associated with Institution Department of Physics, University of Warwick, Coventry, United Kingdom)*
- ⁶⁴*Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil
(associated with Institution Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil)*
- ⁶⁵*Physics and Micro Electronic College, Hunan University, Changsha City, China
(associated with Institution Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China)*
- ⁶⁶*Guangdong Provincial Key Laboratory of Nuclear Science, Guangdong-Hong Kong Joint Laboratory of Quantum Matter,
Institute of Quantum Matter, South China Normal University, Guangzhou, China
(associated with Institution Center for High Energy Physics, Tsinghua University, Beijing, China)*
- ⁶⁷*Lanzhou University, Lanzhou, China (associated with Institution Institute Of High Energy Physics (IHEP), Beijing, China)*
- ⁶⁸*School of Physics and Technology, Wuhan University, Wuhan, China
(associated with Institution Center for High Energy Physics, Tsinghua University, Beijing, China)*
- ⁶⁹*Departamento de Física, Universidad Nacional de Colombia, Bogota, Colombia
(associated with Institution LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France)*
- ⁷⁰*Universität Bonn - Helmholtz-Institut für Strahlen und Kernphysik, Bonn, Germany
(associated with Institution Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany)*
- ⁷¹*Eotvos Lorand University, Budapest, Hungary
(associated with Institution European Organization for Nuclear Research (CERN), Geneva, Switzerland)*
- ⁷²*INFN Sezione di Perugia, Perugia, Italy (associated with Institution INFN Sezione di Ferrara, Ferrara, Italy)*
- ⁷³*Van Swinderen Institute, University of Groningen, Groningen, Netherlands
(associated with Institution Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands)*
- ⁷⁴*Universiteit Maastricht, Maastricht, Netherlands
(associated with Institution Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands)*
- ⁷⁵*DS4DS, La Salle, Universitat Ramon Llull, Barcelona, Spain
(associated with Institution ICCUB, Universitat de Barcelona, Barcelona, Spain)*
- ⁷⁶*Department of Physics and Astronomy, Uppsala University, Uppsala, Sweden
(associated with Institution School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom)*
- ⁷⁷*University of Michigan, Ann Arbor, Michigan, USA (associated with Institution Syracuse University, Syracuse, New York, USA)*

^aDeceased.

^bAlso at Università di Firenze, Firenze, Italy.

^cAlso at Scuola Normale Superiore, Pisa, Italy.

^dAlso at Università di Ferrara, Ferrara, Italy.

^eAlso at Università di Milano Bicocca, Milano, Italy.

^fAlso at Università di Bologna, Bologna, Italy.

^gAlso at Università di Genova, Genova, Italy.

^hAlso at Universidad Nacional Autónoma de Honduras, Tegucigalpa, Honduras.

ⁱAlso at Università di Bari, Bari, Italy.

^jAlso at Università di Cagliari, Cagliari, Italy.

^kAlso at Università di Roma Tor Vergata, Roma, Italy.

^lAlso at Universidade de Brasília, Brasília, Brazil.

^mAlso at Hangzhou Institute for Advanced Study, UCAS, Hangzhou, China.

ⁿAlso at Università di Siena, Siena, Italy.

^oAlso at Central South U., Changsha, China.

^pAlso at Università di Padova, Padova, Italy.

^qAlso at Università degli Studi di Milano, Milano, Italy.

^rAlso at MSU—Iligan Institute of Technology (MSU-IIT), Iligan, Philippines.

^sAlso at Università di Perugia, Perugia, Italy.

^tAlso at Excellence Cluster ORIGINS, Munich, Germany.

^uAlso at Università di Pisa, Pisa, Italy.

^vAlso at Università della Basilicata, Potenza, Italy.

^wAlso at Universidad de Alcalá, Alcalá de Henares, Spain.

^xAlso at Università di Urbino, Urbino, Italy.