


Analytic Solutions for the Motion of Spinning Particles near Spherically Symmetric Black Holes and Exotic Compact Objects

Vojtěch Witzany^{1,*} and Gabriel Andres Piovano^{2,†}

¹*Institute of Theoretical Physics, Faculty of Mathematics and Physics, Charles University, CZ-180 00 Prague, Czech Republic*

²*School of Mathematics and Statistics, University College Dublin, Belfield, Dublin 4, Ireland*

 (Received 8 August 2023; revised 15 January 2024; accepted 26 March 2024; published 23 April 2024)

Rapidly rotating bodies moving in curved space-time experience the so-called spin-curvature force, which becomes important for the motion of compact objects in gravitational-wave inspirals. As a first approximation, this effect is captured in the motion of a spinning test particle. We solve the equations of motion of a spinning particle to leading order in spin in arbitrary static and spherically symmetric space-times in terms of one-dimensional closed-form integrals. This solves the problem and proves its integrability in a wide range of modified gravities and near exotic compact objects. Then, by specializing to the case of bound orbits in Schwarzschild space-time, we demonstrate how to express the solution in the form of Jacobi elliptic functions.

DOI: [10.1103/PhysRevLett.132.171401](https://doi.org/10.1103/PhysRevLett.132.171401)

Introduction.—After a track record of spectacular successes in their first three observing runs [1–3], in May 2023 the LIGO and Virgo instruments were joined for the first time by the Japanese detector KAGRA to start the fourth run of observing coalescing compact objects through gravitational waves [4]. The so-called LVK Collaboration is expected to detect more events in this run than have been amassed to date. Next-generation detectors on Earth and in space promise to multiply the sensitivity and reach of the detectors even further [5,6]. As the number of events grow, we will observe binaries with different mass ranges, mass ratios, and dynamical setups and, consequently, we will need a more faithful and complete picture of the two-body dynamics in order to detect and interpret these signals correctly.

In particular, the space-based observatory Laser Interferometer Space Antenna (LISA) will detect inspirals of stellar-mass compact objects (henceforth dubbed as secondary) onto supermassive black holes [7]. This will allow us to accurately test whether the supermassive objects in the centers of galaxies are truly black holes or perhaps some exotic compact objects, or whether their gravitational field is described by Einstein gravity [8]. These so-called extreme-mass-ratio inspirals (EMRIs) are best modeled in a mass-ratio expansion [9]. At leading order, the two-body dynamics is approximated by the adiabatic inspiral of a test particle in the space-time of the massive compact object, where the inspiral is driven by fluxes sourced by the secondary [10,11]. Subleading corrections to the motion are collectively called post-1-adiabatic and include the effects of second order in the mass-ratio fluxes, conservative self-force, and the spin of the smaller companion [12–14]. Post-1-adiabatic corrections are fundamental to model waveforms suited for data analysis for LISA [7,15], and

the spin of the secondary plays a crucial role [16–18]. The effects of the latter are fully captured by the motion of a spinning test particle in the space-time of the massive compact object (computed to linear order to spin) and the outgoing gravitational-wave flux it sources [19–27]. Here we present an analytical solution of this motion near spherically symmetric compact objects in terms of quadratures. In particular, our analysis implies that the spinning particle motion is integrable at linear order in spin in any static, spherically symmetric space-time. We then specialize to Schwarzschild space-time and reexpress the solution in terms of Jacobi elliptic integrals. Even though a number of works have treated this and similar topics previously (see, e.g., Refs. [27–32]), this is the first time an analytic, closed-form solution of generic bound motion of a spinning test particle in Schwarzschild space-time is presented.

We use the $G=c=1$ geometrized units and the $(-+++)$ signature of the metric. Greek indices label coordinate components. The Riemann tensor is defined by $a_{\nu;\kappa\lambda} - a_{\nu;\lambda\kappa} \equiv R^{\mu}_{\nu\kappa\lambda} a_{\mu}$, where “;” denotes the covariant derivative and a_{μ} is an arbitrary form.

Spinning particle.—The motion of a spinning particle is given by the equations of motion [33–35]

$$\frac{DP^{\mu}}{d\tau} = -\frac{1}{2}R^{\mu}_{\nu\kappa\lambda}\dot{x}^{\nu}S^{\kappa\lambda}, \quad (1)$$

$$\frac{DS^{\mu\nu}}{d\tau} = P^{\mu}\dot{x}^{\nu} - P^{\nu}\dot{x}^{\mu}, \quad (2)$$

where P^{μ} and \dot{x}^{ν} are the 4-momentum and tangent vector to the worldline of the spinning test particle, while $S^{\mu\nu} = -S^{\nu\mu}$ is the spin tensor. We are only interested in the dynamics to linear order in spin $S^{\mu\nu}$ and, unless specified, all formulas

are to be assumed at most $\mathcal{O}(S)$ accurate. This is justified by considering a comparably light compact object in the field of a heavy black hole, since the truncated terms can be shown to scale as higher order in the mass ratio, as discussed in Supplemental Material [36]. We fix the relation of the centroid $x^\mu(\tau)$ and the momentum P^μ by the Tulczyjew-Dixon supplemental spin condition $S^{\mu\nu}P_\nu = 0$, to obtain $P^\mu = m\dot{x}^\mu + \mathcal{O}(S^2)$ where m is the particle mass [35,47,48]. The spin tensor can then be expressed as $S^{\mu\nu} = m\epsilon^{\mu\nu\kappa\lambda}\dot{x}_\kappa s_\lambda/2$, where s^μ is the specific spin vector and $\epsilon^{\mu\nu\kappa\lambda}$ is the Levi-Civita pseudotensor. The equations then reduce to

$$\frac{D^2x^\mu}{d\tau^2} = -\frac{1}{4}R^\mu_{\nu\gamma\delta}\epsilon^{\gamma\delta\kappa\lambda}\dot{x}^\nu\dot{x}^\kappa s^\lambda, \quad (3)$$

$$\frac{Ds^\lambda}{d\tau} = 0. \quad (4)$$

Motion in static, spherically symmetric metric.—A general static, spherically symmetric space-time metric can be locally expressed in the form

$$ds^2 = -f(r)dt^2 + h(r)dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2). \quad (5)$$

The Schwarzschild metric is contained within this class by setting $f(r)$, $h(r)$ to $f(r) = 1/h(r) = 1 - 2M/r$. The Killing vectors of this metric are generators of time translations and rotations around the x , y , and z axes,

$$\xi_{(t)} = \frac{\partial}{\partial t}, \quad (6)$$

$$\xi_{(x)} = -\sin\phi\frac{\partial}{\partial\theta} - \cos\phi\cot\theta\frac{\partial}{\partial\phi}, \quad (7)$$

$$\xi_{(y)} = \cos\phi\frac{\partial}{\partial\theta} - \sin\phi\cot\theta\frac{\partial}{\partial\phi}, \quad (8)$$

$$\xi_{(z)} = \frac{\partial}{\partial\phi}. \quad (9)$$

Dixon [49] showed that for any Killing vector the spinning particle motion has a constant of the form $C_{(\xi)} = P^\mu\xi_\mu - \xi_{\rho;\sigma}S^{\rho\sigma}/2$. In this case, the Killing vectors correspond to a conserved specific spin-orbital energy and a formal angular momentum vector (we normalize each $C_{(\xi)}$ by m),

$$\mathcal{E} = -f\dot{t} + \frac{r^2\sin\theta f' s^\phi\dot{\theta} - s^\theta\dot{\phi}}{2\sqrt{fh}}, \quad (10)$$

$$\begin{aligned} \mathcal{J}_x &= -r^2(\sin\phi\dot{\theta} + \cos\phi\cos\theta\sin\theta\dot{\phi}) \\ &+ \sqrt{\frac{f}{h}}[\sin\theta\cos\phi h(s^t\dot{r} - s^r\dot{t}) \\ &+ r\sin\phi\sin\theta(s^\phi\dot{t} - s^t\dot{\phi}) \\ &+ r\cos\phi\cos\theta(s^t\dot{\theta} - s^\theta\dot{t})], \end{aligned} \quad (11)$$

$$\begin{aligned} \mathcal{J}_y &= r^2(\cos\phi\dot{\theta} - \sin\phi\cos\theta\sin\theta\dot{\phi}) \\ &+ \sqrt{\frac{f}{h}}[\sin\theta\sin\phi h(s^t\dot{r} - s^r\dot{t}) \\ &+ r\cos\phi\sin\theta(s^t\dot{\phi} - s^\phi\dot{t}) \\ &+ r\sin\phi\cos\theta(s^t\dot{\theta} - s^\theta\dot{t})], \end{aligned} \quad (12)$$

$$\begin{aligned} \mathcal{J}_z &= r^2\sin^2\theta\dot{\phi} \\ &+ \sqrt{\frac{f}{h}}[\cos\theta h(s^t\dot{r} - s^r\dot{t}) \\ &+ r\sin\theta(s^\theta\dot{t} - s^t\dot{\theta})]. \end{aligned} \quad (13)$$

The linearization in spin means that these integrals of motion are conserved up to $\mathcal{O}(s^2)$. Another integral of motion is obtained by noticing that the following vector is parallel transported along geodesics in general spherically symmetric static space-times,

$$l = \frac{r\dot{\theta}}{\sin\theta}\frac{\partial}{\partial\phi} - r\sin\theta\dot{\phi}\frac{\partial}{\partial\theta}, \quad (14)$$

and for spinning particles $DI^\mu/d\tau = \mathcal{O}(s)$. As a result, the aligned component of the spin vector is an approximate constant of motion,

$$s_{\parallel} \equiv \frac{l^\mu s_\mu}{\sqrt{l^\nu l_\nu}}, \quad \frac{ds_{\parallel}}{d\tau} = 0 + \mathcal{O}(s^2). \quad (15)$$

Angular momentum aligned coordinates.—For general initial conditions, the formal angular-momentum vector $\vec{\mathcal{J}} = (\mathcal{J}_x, \mathcal{J}_y, \mathcal{J}_z)$ keeps pointing into a constant direction. Additionally, the orbital plane is always almost orthogonal to this vector up to $\mathcal{O}(s)$ corrections. Hence, without loss of generality, we can rotate into a new coordinate system $\theta, \phi \rightarrow \vartheta, \varphi$ such that its axis of φ rotation ($\vartheta = 0, \pi$) points in the direction of $\vec{\mathcal{J}}$. When defined with respect to this new system, we obtain $\mathcal{J}'_x = \mathcal{J}'_y = 0$, $\mathcal{J}'_z = \sqrt{\mathcal{J}_x^2 + \mathcal{J}_y^2 + \mathcal{J}_z^2} \equiv \mathcal{J}$, and the position of the particle will fulfil $\vartheta(\tau) = \pi/2 + \delta\vartheta(\tau)$, where

$$\delta\vartheta = \frac{\sqrt{fh}(s^r \dot{t} - s^t \dot{r})}{r^2 \dot{\varphi}} + \mathcal{O}(s^2). \quad (16)$$

In other words, the ϑ motion of the particle is automatically expressed in terms of the other variables. This transformation was already implicitly used in the numerical studies of Refs. [29,50], even though the authors did not realize it also allows one to analytically solve for the ϑ degree of freedom at $\mathcal{O}(s)$. More details of the computation can be found in Supplemental Material [36].

The equations of motion for the other orbital variables $r(\tau)$, $t(\tau)$, and $\varphi(\tau)$ can now be expressed in first-order form up to $\mathcal{O}(s^2)$,

$$\dot{r}^2 = \frac{1}{h} \left(-1 + \frac{\mathcal{E}^2}{f} - \frac{\mathcal{J}^2}{r^2} \right) + \frac{s_{\parallel} \mathcal{E} \mathcal{J} (2f - rf')}{(fh)^{3/2} r^2}, \quad (17)$$

$$\dot{t} = \frac{\mathcal{E}}{f} + \frac{s_{\parallel} \mathcal{J} f'}{2fr\sqrt{fh}}, \quad (18)$$

$$\dot{\varphi} = \frac{\mathcal{J}}{r^2} + \frac{s_{\parallel} \mathcal{E}}{r^2 \sqrt{fh}}. \quad (19)$$

Solving parallel transport.—As for the spin degree of freedom, we need to solve the parallel transport equation at leading (geodesic) order. To do so, we construct a parallel-transported tetrad $e_{(A)}^{\mu}$, $A = 0, \dots, 3$ inspired by Marck [51]. We start with the zeroth and third leg (component order t, r, φ, ϑ),

$$e_{(0)}^{\mu} = (\mathcal{E}/f, \dot{r}, \mathcal{J}/r^2, 0), \quad (20)$$

$$e_{(3)}^{\mu} = \left(0, 0, 0, -\frac{1}{r} \right), \quad (21)$$

where \dot{r} is given by Eq. (17). Note that the full expression for \dot{r} including $\mathcal{O}(s)$ corrections avoids singularities in the definition of the tetrad at radial turning points. Now the first and second legs are given as $e_{(1)}^{\mu} = \cos\psi \tilde{e}_{(1)}^{\mu} + \sin\psi \tilde{e}_{(2)}^{\mu}$ and $e_{(2)}^{\mu} = -\sin\psi \tilde{e}_{(1)}^{\mu} + \cos\psi \tilde{e}_{(2)}^{\mu}$, where

$$\tilde{e}_{(1)}^{\mu} = \left(\frac{r\dot{r}\sqrt{h}}{\sqrt{f(\mathcal{J}^2 + r^2)}}, \frac{\mathcal{E}r}{\sqrt{fh(\mathcal{J}^2 + r^2)}}, 0, 0 \right), \quad (22)$$

$$\tilde{e}_{(2)}^{\mu} = \left(\frac{\mathcal{E}\mathcal{J}}{f\sqrt{\mathcal{J}^2 + r^2}}, \frac{\mathcal{J}\dot{r}}{r}, \frac{\sqrt{\mathcal{J}^2 + r^2}}{r^2}, 0 \right), \quad (23)$$

and the precession angle $\psi(\tau)$ is obtained by integrating

$$\dot{\psi} = \frac{\mathcal{E}\mathcal{J}}{\mathcal{J}^2 + r^2}. \quad (24)$$

The components of the spin with respect to the parallel-transported tetrad $e_{(A)}^{\mu}$ are constant at leading order.

Additionally, since the spin magnitude $s \equiv \sqrt{s^{\mu} s_{\mu}}$ is conserved and $s_{\mu} e_{(3)}^{\mu} = s_{\parallel} + \mathcal{O}(s^2)$, the solution for the evolution of the spin vector can be expressed as

$$s^t = \sqrt{\frac{s^2 - s_{\parallel}^2}{f(\mathcal{J}^2 + r^2)}} \left(\frac{\mathcal{E}\mathcal{J} \cos\psi}{\sqrt{f}} + r\dot{r} \sin\psi \right), \quad (25a)$$

$$s^r = \sqrt{s^2 - s_{\parallel}^2} \left(\frac{\mathcal{J}\dot{r} \cos\psi}{r} + \frac{\mathcal{E}r \sin\psi}{\sqrt{fh(\mathcal{J}^2 + r^2)}} \right), \quad (25b)$$

$$s^{\varphi} = \frac{\sqrt{(s^2 - s_{\parallel}^2)(\mathcal{J}^2 + r^2)} \cos\psi}{r^2}, \quad (25c)$$

$$s^{\vartheta} = -\frac{s_{\parallel}}{r}. \quad (25d)$$

One can now also reexpress $\delta\vartheta$ from Eq. (16) as

$$\delta\vartheta = \frac{\sqrt{(s^2 - s_{\parallel}^2)(\mathcal{J}^2 + r^2)} \sin\psi}{\mathcal{J}r}. \quad (26)$$

Solution by quadrature.—The motion of the spinning test particle in any static, spherically symmetric metric is solved by quadrature as follows. First, one integrates

$$\tau(r) - \tau(r_0) = \pm \int_{r_0}^r \frac{dr'}{\sqrt{\mathcal{R}(r')}}, \quad (27)$$

$$\mathcal{R}(r) \equiv \frac{1}{h} \left(-1 + \frac{\mathcal{E}^2}{f} - \frac{\mathcal{J}^2}{r^2} \right) + \frac{s_{\parallel} \mathcal{E} \mathcal{J} (2f - rf')}{(fh)^{3/2} r^2} \quad (28)$$

and inverts this relation to obtain $r(\tau)$. This is then substituted into the r -parametrized solutions

$$t(r) - t(r_0) = \pm \int_{r_0}^r \frac{dr'}{\sqrt{\mathcal{R}(r')}} \left[\frac{\mathcal{E}}{f} + \frac{s_{\parallel} \mathcal{J} f'}{2f r' \sqrt{fh}} \right], \quad (29)$$

$$\varphi(r) - \varphi(r_0) = \pm \int_{r_0}^r \frac{dr'}{\sqrt{\mathcal{R}(r')}} \left[\frac{\mathcal{J}}{r'^2} + \frac{s_{\parallel} \mathcal{E}}{r'^2 \sqrt{fh}} \right], \quad (30)$$

$$\psi(r) - \psi(r_0) = \pm \int_{r_0}^r \frac{dr'}{\sqrt{\mathcal{R}(r')}} \left[\frac{\mathcal{E}\mathcal{J}}{\mathcal{J}^2 + r'^2} \right]. \quad (31)$$

The aligned coordinate position $\vartheta = \pi/2 + \delta\vartheta$ and the spin components s^{μ} are then obtained by substituting the solutions for $r(\tau)$ and $\psi(\tau)$ into Eqs. (25) and (26).

Schwarzschild space-time.—Let us demonstrate how to use this solution near a Schwarzschild black hole by setting $f = 1/h = 1 - 2M/r$. We focus on bound motion between radial turning points $r_1 > r_2$ which are real roots of $\mathcal{R}(r)$. Following Darwin's treatment of geodesics [52], we parametrize the motion by eccentricity e and semilatus rectum

p such that $r_1 = p/(1 - e)$, $r_2 = p/(1 + e)$. The relation between energy, angular momentum, and the orbital elements e , p is obtained by inserting the geodesic relations from Darwin into the equation $\mathcal{R}(r) = 0$ and computing the spin corrections

$$\mathcal{E}^2 = \frac{(p-2M)^2 - 4M^2 e^2}{p[p-M(3+e^2)]} + s_{\parallel} \frac{(e^2-1)^2 M \sqrt{Mp[p^2-4Mp-4M^2(e^2-1)]}}{p^2[p-M(3+e^2)]^2}, \quad (32)$$

$$\mathcal{J}^2 = \frac{Mp^2}{p-M(3+e^2)} - s_{\parallel} \frac{[2p-3M(3+e^2)] \sqrt{Mp[p^2-4Mp-4M^2(e^2-1)]}}{[p-M(3+e^2)]^2}. \quad (33)$$

Interestingly, these shifts exactly agree with those for a particle with spin fully aligned with orbital angular momentum when $s_{\parallel} = s$ (cf. Refs. [53,54]). The bound motion exists for $e \in [0, 1)$ and $p \in (p_c(e), \infty)$, where the last stable orbits $p_c(e)$ are determined by the vanishing of the Jacobian of the transform in Eqs. (32) and (33), which yields (cf. the geodesic case in Ref. [55])

$$p_c = (6 + 2e)M + 2s_{\parallel} \sqrt{\frac{2(1+e)}{3+e}}. \quad (34)$$

This generalizes the results for innermost stable circular orbits of particles with aligned spin from Refs. [29,30,56,57] to fully generic motion. The function $\mathcal{R}(r)$ can now be reexpressed as

$$\mathcal{R} = \frac{1 - \mathcal{E}^2}{r^4} (r_1 - r)(r - r_2)(r - r_3)(r - r_4), \quad (35)$$

$$r_1 > r_2 > r_3 > r_4, \quad r_{1,2} = \frac{p}{1 \mp e}, \quad r_4 = 0, \quad (36)$$

$$r_3 = \frac{2Mp}{p-4M} + \frac{2s_{\parallel} \sqrt{Mp[p^2-4Mp+4M^2(1-e^2)]}}{(p-4M)^2}. \quad (37)$$

Expression as Jacobi elliptic integrals.—In Schwarzschild space-time, the most elegant parametrization of the motion is through Carter-Mino time [58,59] λ , $d\tau/d\lambda = r^2$ so that

$$\lambda(r) - \lambda(r_2) = \int_{r_2}^r \frac{dr'}{\sqrt{R(r')}} = \frac{2F(\chi, k)}{\sqrt{(1 - \mathcal{E}^2)(r_1 - r_3)r_2}}, \quad (38)$$

$$\sin \chi \equiv \sqrt{\frac{(r_1 - r_3)(r - r_2)}{(r_1 - r_2)(r - r_3)}}, \quad (39)$$

$$k^2 = \frac{(r_1 - r_2)r_3}{(r_1 - r_3)r_2}, \quad (40)$$

$$R(r) \equiv (1 - \mathcal{E}^2)(r_1 - r)(r - r_2)(r - r_3)r, \quad (41)$$

where $F(\chi, k)$ is the elliptic integral of the first kind. All elliptic integrals F , K , E , Π and their inverses am , sn , cn will be defined in the angle-modulus convention of Byrd and Friedman [60] (see also Supplemental Material [36]). The $\lambda(r)$ function can be inverted by the same arguments as given by Van de Meent [61] for corresponding formulas for Kerr geodesics to yield

$$r(\lambda) = \frac{r_3(r_1 - r_2)\text{sn}^2\left(\frac{K(k)}{\pi} q^r, k\right) - r_2(r_1 - r_3)}{(r_1 - r_2)\text{sn}^2\left(\frac{K(k)}{\pi} q^r, k\right) - (r_1 - r_3)}, \quad (42)$$

$$q^r \equiv \Upsilon^r \lambda + q_0^r, \quad (43)$$

$$\Upsilon^r \equiv \frac{\pi \sqrt{(1 - \mathcal{E}^2)(r_1 - r_3)r_2}}{2K(k)}, \quad (44)$$

where $K(k)$ is the complete elliptic integral of the first kind and q_0^r is an integration constant determined by initial conditions. Again, following closely the notation and approach of Ref. [61], we express the other orbital variables as

$$t(\lambda) = q^t + \tilde{T}_r \left(\text{am} \left(\frac{q^r}{\pi} K(k), k \right) \right) - \frac{\tilde{T}_r(\pi)}{2\pi} q^r, \quad (45)$$

$$\varphi(\lambda) = q^\varphi + \tilde{\Phi}_r \left(\text{am} \left(\frac{q^r}{\pi} K(k), k \right) \right) - \frac{\tilde{\Phi}_r(\pi)}{2\pi} q^r, \quad (46)$$

$$\psi(\lambda) = q^\psi + \tilde{\Psi}_r \left(\text{am} \left(\frac{q^r}{\pi} K(k), k \right) \right) - \frac{\tilde{\Psi}_r(\pi)}{2\pi} q^r, \quad (47)$$

$$q^t \equiv \Upsilon^t \lambda + q_0^t, \quad q^\varphi \equiv \Upsilon^\varphi \lambda + q_0^\varphi, \quad q^\psi \equiv \Upsilon^\psi \lambda + q_0^\psi, \quad (48)$$

where $q_0^{t,\varphi,\psi}$ are again integration constants and

$$\begin{aligned} \tilde{T}_r(\chi) = & \frac{\mathcal{E}}{\sqrt{(1 - \mathcal{E}^2)r_2(r_1 - r_3)}} \left[\frac{2M(r_1 - r_3)(r_2 - r_3) + r_3(r_3(r_2 + r_3) - r_1(r_2 - r_3)) + 2\frac{\mathcal{J}}{\mathcal{E}}Ms_{\parallel} F(\chi, k)}{r_3 - 2M} \right. \\ & + (r_1 + r_2 + r_3 + 4M)(r_2 - r_3) \Pi \left(\chi, \frac{r_1 - r_2}{r_1 - r_3}, k \right) + r_2(r_1 - r_3) E(\chi, k) \\ & \left. - \frac{2M(r_2 - r_3)(8M^2 + \frac{\mathcal{J}}{\mathcal{E}}s_{\parallel})}{(r_2 - 2M)(r_3 - 2M)} \Pi \left(\chi, \frac{(r_3 - 2M)(r_1 - r_2)}{(r_2 - 2M)(r_1 - r_3)}, k \right) - \frac{(r_1 - r_2) \sin 2\chi \sqrt{r_2(r_1 - r_3)[r_2(r_1 - r_3) - r_3(r_1 - r_2) \sin^2 \chi]}}{r_1 + r_2 - 2r_3 + (r_1 - r_2) \cos 2\chi} \right], \end{aligned} \quad (49)$$

$$\tilde{\Phi}_r(\chi) = \frac{2(\mathcal{J} + s_{\parallel}\mathcal{E})}{\sqrt{(1-\mathcal{E}^2)r_2(r_1-r_3)}} F(\chi, k), \quad (50)$$

$$\begin{aligned} \tilde{\Psi}_r(\chi) = & \frac{2\mathcal{E}\mathcal{J}}{\sqrt{(1-\mathcal{E}^2)(r_1-r_3)r_2(\mathcal{J}^2+r_3^2)}} \left[r_3^2 F(\chi, k) + \frac{\mathcal{J}^2(r_2^2-r_3^2)}{\mathcal{J}^2+r_2^2} \operatorname{Re} \left(\Pi \left(\chi, \frac{r_3-i\mathcal{J}r_1-r_2}{r_2-i\mathcal{J}r_1-r_3}, k \right) \right) \right. \\ & \left. - \frac{\mathcal{J}(r_2-r_3)(\mathcal{J}^2-r_2r_3)}{\mathcal{J}^2+r_2^2} \operatorname{Im} \left(\Pi \left(\chi, \frac{r_3-i\mathcal{J}r_1-r_2}{r_2-i\mathcal{J}r_1-r_3}, k \right) \right) \right], \end{aligned} \quad (51)$$

$$\begin{aligned} \Upsilon^t = \frac{\Upsilon^r}{2\pi} \tilde{T}_r(\pi) = & \frac{\mathcal{E}}{2K(k)} \left[\frac{2M(r_1-r_3)(r_2-r_3) + r_3(r_3(r_2+r_3) - r_1(r_2-r_3))}{r_3-2M} K(k) \right. \\ & + (r_1+r_2+r_3+4M)(r_2-r_3) \Pi \left(\frac{r_1-r_2}{r_1-r_3}, k \right) + r_2(r_1-r_3)E(k) \\ & \left. - \frac{(r_2-r_3)16M^3}{(r_2-2M)(r_3-2M)} \Pi \left(\frac{(r_3-2M)(r_1-r_2)}{(r_2-2M)(r_1-r_3)}, k \right) - \frac{\mathcal{J}s_{\parallel}}{\mathcal{E}} \left(1 - \frac{r_1}{r_1-2M} \Pi \left(\frac{2M(r_1-r_2)}{r_2(r_1-2M)}, k \right) \right) \right], \end{aligned} \quad (52)$$

$$\Upsilon^\varphi = \frac{\Upsilon^r}{2\pi} \tilde{\Phi}_r(\pi) = \mathcal{J} + s_{\parallel}\mathcal{E}, \quad (53)$$

$$\begin{aligned} \Upsilon^\psi = \frac{\Upsilon^r}{2\pi} \tilde{\Psi}_r(\pi) = & \frac{\mathcal{E}\mathcal{J}r_3^2}{\mathcal{J}^2+r_3^2} + \frac{\mathcal{E}\mathcal{J}^2(r_2-r_3)}{K(k)(\mathcal{J}^2+r_2^2)(\mathcal{J}^2+r_3^2)} \left[\mathcal{J}(r_2+r_3) \operatorname{Re} \left(\Pi \left(\frac{r_3-i\mathcal{J}r_1-r_2}{r_2-i\mathcal{J}r_1-r_3}, k \right) \right) \right. \\ & \left. - (\mathcal{J}^2-r_2r_3) \operatorname{Im} \left(\Pi \left(\frac{r_3-i\mathcal{J}r_1-r_2}{r_2-i\mathcal{J}r_1-r_3}, k \right) \right) \right]. \end{aligned} \quad (54)$$

This allows us to plot fully explicit orbital solutions, as demonstrated in Fig. 1. It should be noted that Υ^ψ , $\tilde{\Psi}_r(\chi)$ can be evaluated at $s_{\parallel} = 0$ at leading order (with no spin correction to r_3 , \mathcal{E} , \mathcal{J}). On the other hand, $\Upsilon^{r,t,\varphi}$, \tilde{T}_r , and $\tilde{\Phi}_r$ need to be explicitly expanded in s_{\parallel} due to the spin corrections to r_3 , \mathcal{E} , and \mathcal{J} at fixed p , e [see Eqs. (32), (33), and (37)]. This procedure is straightforward, but the results are lengthy and thus relegated to Supplemental




FIG. 1. A generic inclined spinning particle orbit (left) and its description in angular momentum aligned coordinates (right) with $p = 7M$, $e = 0.3$, $s_{\parallel} = 0$, $s = 0.3M$.

Material [36]. From the ratios of these frequencies, one can compute coordinate-time frequencies $\Omega^{r,\varphi,\psi} \equiv \Upsilon^{r,\varphi,\psi}/\Upsilon^t$ or the nodal and periastron precession rates, which is also discussed in Supplemental Material [36].

Discussion and outlook.—The presented analytical solution for the motion of spinning test particles in Schwarzschild space-time can be taken “as is” to source gravitational-wave fluxes or to compute the gravitational self-force on the particle [18–27]. Another useful output will be to extend the solution to compute action variables and scattering orbits in order to compare to effective one-body models [62–65] and other approaches to the relativistic two-body problem [66,67]. The formulas provided in the notebooks along with this Letter can also be straightforwardly implemented into existing code bases such as the BHPToolkit [68], thus providing an immediate impact in gravitational-wave modeling. In particular, this solution is a springboard to understand the radiation reaction on the fully generic motion of spinning particles in Kerr space-time [27,32,69]. Indeed, we have recently constructed semianalytical orbits of spinning particles near Kerr black holes by using the Hamilton-Jacobi formalism presented in Ref. [69], and it is proving important to have independently derived analytical formulas for validation, even if just for the Schwarzschild case. An important application will also be to use the quadratures in Eqs. (27)–(31) to study deviations of the particle motion in modified gravities such as in Einstein-Gauss-Bonnet gravity [70,71], Horndeski

and beyond-Horndeski theories [72,73], or near boson stars [74–77] and other exotic compact objects [78].

However, we can already use our results to draw general conclusions about the detectability of secondary spin in large-mass-ratio inspirals onto spherically symmetric compact objects. Our solution demonstrates that time-averaged observables depend only on the aligned component of spin s_{\parallel} , independent of the compact-object model, and the contributions of the orthogonal component $s_{\perp}, s_{\perp}^2 \equiv s^2 - s_{\parallel}^2$ to any dynamical variable will oscillate with the characteristic frequency Ω^{ψ} generically different from the orbital frequencies. The spin contributions to the post-1-adiabatic phase of the waveform depend only on the spin shifts to quantities such as average orbital frequencies and the average energy- and angular-momentum fluxes, which are independent of s_{\perp} (see also Ref. [24]). Furthermore, the Ω^{ψ} oscillation due to $s_{\perp} \lesssim m$ only appears in the $\mathcal{O}(m/M)$ subdominant corrections to the waveform amplitude [79,80]. As a result, waveform templates that neglect s_{\perp} contributions will have a mismatch with models that include s_{\perp} of order $\mathcal{O}(m^2/M^2)$. Hence, according to the Lindblom criterion, the waveforms that either include or neglect s_{\perp} for EMRIs will be indistinguishable from each other in a matched filtering analysis unless the signal-to-noise ratio of the EMRI signal reaches the order of the large mass ratio $M/m \sim 10^4 - 10^6$ [81,82]. This provides a key insight that will vastly simplify the treatment of spin in modified-gravity waveforms at large mass ratios.

Another consequence of this Letter is that the motion of spinning particles in any static spherically symmetric space-time is integrable to linear order in spin. In other words, if there is any chaos to be observed by the numerical integration of spinning particle motion [29,31], it is an $\mathcal{O}(s^2)$ effect that goes beyond the validity of the equations themselves, which is in line with the quantitative numerical scalings of resonances observed in Schwarzschild space-time [31].

This work makes use of the Wolfram *Mathematica* package KerrGeodesics [83], which is a part of the BHPTToolkit [68].

We would like to thank Viktor Skoupý and Josh Mathews for careful reading of an earlier draft of this Letter. V. W. is grateful for the support from the Charles University *Primus* program. G. A. P. acknowledges support from an Irish Research Council Fellowship under Grant No. GOIPD/2022/496.

*vojtech.witzany@mattfyz.cuni.cz

†gabriel.piovano@ucd.ie

[1] B. P. Abbott *et al.* (LIGO Scientific, Virgo Collaborations), *Phys. Rev. X* **9**, 031040 (2019).

- [2] R. Abbott *et al.* (LIGO Scientific, Virgo Collaborations), *Phys. Rev. X* **11**, 021053 (2021).
- [3] R. Abbott *et al.* (KAGRA, VIRGO, LIGO Scientific Collaborations), *Phys. Rev. X* **13**, 041039 (2023).
- [4] B. P. Abbott *et al.* (KAGRA, LIGO Scientific, Virgo, VIRGO Collaborations), *Living Rev. Relativity* **21**, 3 (2018).
- [5] V. Kalogera *et al.*, arXiv:2111.06990.
- [6] LISA Consortium, arXiv:1702.00786.
- [7] S. Babak, J. Gair, A. Sesana, E. Barausse, C. F. Sopuerta, C. P. L. Berry, E. Berti, P. Amaro-Seoane, A. Petiteau, and A. Klein, *Phys. Rev. D* **95**, 103012 (2017).
- [8] K. G. Arun *et al.* (LISA), *Living Rev. Relativity* **25**, 4 (2022).
- [9] L. Barack and A. Pound, *Rep. Prog. Phys.* **82**, 016904 (2019).
- [10] S. A. Hughes, N. Warburton, G. Khanna, A. J. K. Chua, and M. L. Katz, *Phys. Rev. D* **103**, 104014 (2021); **107**, 089901 (E) (2023).
- [11] S. A. Hughes, S. Drasco, E. E. Flanagan, and J. Franklin, *Phys. Rev. Lett.* **94**, 221101 (2005).
- [12] M. van de Meent, *Phys. Rev. D* **97**, 104033 (2018).
- [13] N. Warburton, A. Pound, B. Wardell, J. Miller, and L. Durkan, *Phys. Rev. Lett.* **127**, 151102 (2021).
- [14] B. Wardell, A. Pound, N. Warburton, J. Miller, L. Durkan, and A. Le Tiec, *Phys. Rev. Lett.* **130**, 241402 (2023).
- [15] L. Barack *et al.*, *Classical Quantum Gravity* **36**, 143001 (2019).
- [16] E. A. Huerta, J. R. Gair, and D. A. Brown, *Phys. Rev. D* **85**, 064023 (2012).
- [17] E. A. Huerta and J. R. Gair, *Phys. Rev. D* **84**, 064023 (2011).
- [18] G. A. Piovano, R. Brito, A. Maselli, and P. Pani, *Phys. Rev. D* **104**, 124019 (2021).
- [19] N. Warburton, T. Osburn, and C. R. Evans, *Phys. Rev. D* **96**, 084057 (2017).
- [20] S. Akcay, S. R. Dolan, C. Kavanagh, J. Moxon, N. Warburton, and B. Wardell, *Phys. Rev. D* **102**, 064013 (2020).
- [21] J. Mathews, A. Pound, and B. Wardell, *Phys. Rev. D* **105**, 084031 (2022).
- [22] V. Skoupý and G. Lukes-Gerakopoulos, *Phys. Rev. D* **103**, 104045 (2021).
- [23] V. Skoupý and G. Lukes-Gerakopoulos, *Phys. Rev. D* **105**, 084033 (2022).
- [24] V. Skoupý, G. Lukes-Gerakopoulos, L. V. Drummond, and S. A. Hughes, *Phys. Rev. D* **108**, 044041 (2023).
- [25] G. A. Piovano, A. Maselli, and P. Pani, *Phys. Rev. D* **102**, 024041 (2020).
- [26] G. A. Piovano, A. Maselli, and P. Pani, *Phys. Rev. D* **107**, 024021 (2023).
- [27] L. V. Drummond, P. Lynch, A. G. Hanselman, D. R. Becker, and S. A. Hughes, *Phys. Rev. D* **109**, 064030 (2024).
- [28] R. H. Rietdijk and J. W. van Holten, *Classical Quantum Gravity* **10**, 575 (1993).
- [29] S. Suzuki and K. I. Maeda, *Phys. Rev. D* **58**, 023005 (1998).
- [30] P. I. Jefremov, O. Y. Tsupko, and G. S. Bisnovaty-Kogan, *Phys. Rev. D* **91**, 124030 (2015).
- [31] O. Zelenka, G. Lukes-Gerakopoulos, V. Witzany, and O. Kopáček, *Phys. Rev. D* **101**, 024037 (2020).

- [32] L. V. Drummond and S. A. Hughes, *Phys. Rev. D* **105**, 124041 (2022).
- [33] M. Mathisson, *Gen. Relativ. Gravit.* **42**, 1011 (2010).
- [34] A. Papapetrou, *Proc. R. Soc. A* **209**, 248 (1951).
- [35] W. Dixon, *Nuovo Cimento (1955–1965)* **34**, 317 (1964).
- [36] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.132.171401>, for further details of the derivations in this Letter, which includes additional Refs. [37–46].
- [37] I. Bailey and W. Israel, *Commun. Math. Phys.* **42**, 65 (1975).
- [38] J. Ehlers and E. Rudolph, *Gen. Relativ. Gravit.* **8**, 197 (1977).
- [39] T. Binnington and E. Poisson, *Phys. Rev. D* **80**, 084018 (2009).
- [40] J. Steinhoff and D. Puetzfeld, *Phys. Rev. D* **86**, 044033 (2012).
- [41] J. Steinhoff, T. Hinderer, A. Buonanno, and A. Taracchini, *Phys. Rev. D* **94**, 104028 (2016).
- [42] J. Steinhoff and G. Schäfer, *Europhys. Lett.* **87**, 50004 (2009).
- [43] S. Chakrabarti, T. Delsate, N. Gürlebeck, and J. Steinhoff, *Phys. Rev. Lett.* **112**, 201102 (2014).
- [44] C. Møller, On the Definition of the Centre of Gravity of an Arbitrary Closed System in the Theory of Relativity, *Communications of the Dublin Institute for Advanced Studies* (1949), <https://dair.dias.ie/id/eprint/20/>.
- [45] A. Morbidelli, *Modern Celestial Mechanics: Aspects of Solar System Dynamics* (Taylor & Francis, London, 2002).
- [46] R. Fujita and W. Hikida, *Classical Quantum Gravity* **26**, 135002 (2009).
- [47] W. Tulczyjew, *Acta Phys. Pol.* **18**, 393 (1959).
- [48] O. Semerák, *Mon. Not. R. Astron. Soc.* **308**, 863 (1999).
- [49] W. G. Dixon, *Proc. R. Soc. A* **314**, 499 (1970).
- [50] O. Zelenka, G. Lukes-Gerakopoulos, V. Witzany, and O. Kopáček, *Phys. Rev. D* **101**, 024037 (2020).
- [51] J.-A. Marck, *Proc. R. Soc. A* **385**, 431 (1983).
- [52] C. G. Darwin, *Proc. R. Soc. A* **249**, 180 (1959).
- [53] L. V. Drummond and S. A. Hughes, *Phys. Rev. D* **105**, 124040 (2022).
- [54] S. Mukherjee and S. Tripathy, *Phys. Rev. D* **101**, 124047 (2020).
- [55] C. Cutler, D. Kennefick, and E. Poisson, *Phys. Rev. D* **50**, 3816 (1994).
- [56] M. Favata, *Phys. Rev. D* **83**, 024028 (2011).
- [57] O. Y. Tsupko, G. Bisnovatyi-Kogan, and P. Jefremov, *Gravitation Cosmol.* **22**, 138 (2016).
- [58] B. Carter, *Phys. Rev.* **174**, 1559 (1968).
- [59] Y. Mino, *Phys. Rev. D* **67**, 084027 (2003).
- [60] P. F. Byrd and M. D. Friedman, *Handbook of Elliptic Integrals for Engineers and Physicists* (Springer, New York, 2013), Vol. 67.
- [61] M. van de Meent, *Classical Quantum Gravity* **37**, 145007 (2020).
- [62] A. Buonanno and T. Damour, *Phys. Rev. D* **59**, 084006 (1999).
- [63] S. Akcay, R. Gamba, and S. Bernuzzi, *Phys. Rev. D* **103**, 024014 (2021).
- [64] A. Albertini, R. Gamba, A. Nagar, and S. Bernuzzi, *Phys. Rev. D* **109**, 044022 (2024).
- [65] M. Khalil, A. Buonanno, H. Estelles, D. P. Mihaylov, S. Ossokine, L. Pompili, and A. Ramos-Buades, *Phys. Rev. D* **108**, 124036 (2023).
- [66] L. Blanchet, A. Buonanno, and A. Le Tiec, *Phys. Rev. D* **87**, 024030 (2013).
- [67] Z. Bern, A. Luna, R. Roiban, C.-H. Shen, and M. Zeng, *Phys. Rev. D* **104**, 065014 (2021).
- [68] Black Hole Perturbation Toolkit, <http://bhptoolkit.org/>.
- [69] V. Witzany, *Phys. Rev. D* **100**, 104030 (2019).
- [70] D. Glavan and C. Lin, *Phys. Rev. Lett.* **124**, 081301 (2020).
- [71] Y.-P. Zhang, S.-W. Wei, and Y.-X. Liu, *Universe* **6**, 103 (2020).
- [72] E. Babichev, C. Charmousis, and A. Lehébel, *J. Cosmol. Astropart. Phys.* **04** (2017) 027.
- [73] A. Bakopoulos, C. Charmousis, P. Kanti, and N. Lecoeur, *J. High Energy Phys.* **08** (2022) 055.
- [74] M. Kesden, J. Gair, and M. Kamionkowski, *Phys. Rev. D* **71**, 044015 (2005).
- [75] C. F. B. Macedo, P. Pani, V. Cardoso, and L. C. B. Crispino, *Phys. Rev. D* **88**, 064046 (2013).
- [76] V. Diemer, K. Eilers, B. Hartmann, I. Schaffer, and C. Toma, *Phys. Rev. D* **88**, 044025 (2013).
- [77] Y. Brihaye, V. Diemer, and B. Hartmann, *Phys. Rev. D* **89**, 084048 (2014).
- [78] V. Cardoso and P. Pani, *Living Rev. Relativity* **22**, 4 (2019).
- [79] E. E. Flanagan and S. A. Hughes, *Phys. Rev. D* **57**, 4535 (1998).
- [80] A. Pound and B. Wardell, [arXiv:2101.04592](https://arxiv.org/abs/2101.04592).
- [81] L. Lindblom, B. J. Owen, and D. A. Brown, *Phys. Rev. D* **78**, 124020 (2008).
- [82] B. F. Schutz, in *The Detection of Gravitational Waves* (Cambridge University Press, Cambridge, England, 1991), pp. 406–451.
- [83] N. Warburton, B. Wardell, O. Long, S. Upton, P. Lynch, Z. Nasipak, L. C. Stein, *KerrGeodesics*, Version 0.9.0 (2023), [10.5281/zenodo.8108265](https://zenodo.org/record/8108265).