

**Fast Rare Events in Exit Times Distributions of Jump Processes**Alessandro Vezzani 

*Istituto dei Materiali per l'Elettronica ed il Magnetismo (IMEM-CNR), Parco Area delle Scienze, 37/A-43124 Parma, Italy;*  
*Dipartimento di Scienze Matematiche, Fisiche e Informatiche, Università degli Studi di Parma,*  
*Parco Area delle Scienze, 7/A 43124 Parma, Italy;*  
*and INFN, Gruppo Collegato di Parma, Parco Area delle Scienze 7/A, 43124 Parma, Italy*

Raffaella Burioni

*Dipartimento di Scienze Matematiche, Fisiche e Informatiche, Università degli Studi di Parma,*  
*Parco Area delle Scienze, 7/A 43124 Parma, Italy*  
*and INFN, Gruppo Collegato di Parma, Parco Area delle Scienze 7/A, 43124 Parma, Italy*

 (Received 19 September 2023; revised 11 March 2024; accepted 4 April 2024; published 2 May 2024)

Rare events in the first-passage distributions of jump processes are capable of triggering anomalous reactions or series of events. Estimating their probability is particularly important when the jump probabilities have broad-tailed distributions, and rare events are therefore not so rare. We formulate a general approach for estimating the contribution of fast rare events to the exit probabilities in the presence of fat-tailed distributions. Using this approach, we study three jump processes that are used to model a wide class of phenomena ranging from biology to transport in disordered systems, ecology, and finance: discrete time random walks, Lévy walks, and the Lévy-Lorentz gas. We determine the exact form of the scaling function for the probability distribution of fast rare events, in which the jump process exits from an interval in a very short time at a large distance opposite to the starting point. In particular, we show that events occurring on timescales orders of magnitude smaller than the typical timescale of the process can make a significant contribution to the exit probability. Our results are confirmed by extensive numerical simulations.

DOI: [10.1103/PhysRevLett.132.187101](https://doi.org/10.1103/PhysRevLett.132.187101)

A stochastic process that reaches a certain threshold value for the first time can trigger many events: a chemical reaction occurs [1], a target is reached [2–7], the dynamics of a financial process starts [8], biological and ecological processes take place [9,10]. The study of these triggering events and the probability of their occurrence is based on the knowledge of first-passage probabilities [11]. As history-dependent quantities, first-passage probabilities are difficult to determine: general results are typically available for the average first-passage time and its lower-order moments [11,12], but knowledge of complete first-passage distributions is generally limited. Given the “trigger” property of exit times, the tails of these distributions are particularly important as they allow the probability of rare anomalous events to be estimated, such as, for example, an exit in a very short time [13]. These estimates are particularly relevant in the case of broad-tailed distributions, where rare events are not so rare.

In particular, a few results are available for first-passage probabilities of jump processes, which are stochastic processes that involve random jumps between different states or positions, occurring at random times and with random magnitudes [14]. Jump processes are so important in modeling the dynamics of stochastic processes in many fields that even their one-dimensional formulation is of

great relevance [15–17]. In the one-dimensional case, exit time probabilities refer to the process leaving a particular state or interval within a specified time. In stochastic processes with two alternative outcomes, the exit side can also be of importance [10], for example to quantify observables such as transmission or backscattering probabilities [18]. A well-known example of jump processes is that of random walks (RWs), in discrete and continuous time [19,20], for which recent results have been obtained [21–23]. An interesting question is to determine the exit probabilities from the side of a domain opposite to the starting point. A rare event in this case corresponds to a fast walker leaving the interval in a very short time, i.e., in a time where the typical distances covered are still much shorter than the size of the domain. To exit in a short time, the trajectory of the walker should correspond to a long jump, allowing it to travel far from the starting point. When fat-tailed distributions for jumps are present in the stochastic process, these very fast events may occur in timescales orders of magnitude shorter than expected, proving crucial in predicting anomalous behaviors.

Recently, we have investigated the role played in rare events by the so-called big jump principle [24]. The principle explains extreme events in a wide class of systems with

heavy-tailed distributions not in terms of an accumulation of many small subevents but as an effect of only the biggest event, the big jump [25]. The principle has been successfully applied to characterize the tail of the probability distribution, at distances much larger than their scaling length, for a wide class of jump processes for RWs involving Lévy and subexponential statistics for space and time [24,26–28], even in disordered settings such as the Lévy-Lorentz gas [29,30].

In this Letter, we first extend the big jump approach to first-passage probabilities and formulate the principle in a general way that makes it applicable to a large set of processes with jumps following subexponential statistics, in the sense of [25,31]. In particular, for a walker starting at the origin, we analytically determine the probability of reaching for the first time a distance  $X$  without being absorbed at the origin, in the asymptotic limit where  $X$  is much larger than the scaling length of the process (i.e., in the limit of very fast atypical events). We consider three well-known models of one-dimensional jump processes: the discrete time RW, the most basic jump process [20]; the Lévy walk, widely used to model animal movement and search patterns [32,33]; and the Lévy-Lorentz gas, a relevant model in the study of transport in complex and disordered media [34]. For this last model, there is currently no known first-passage probability estimate [35,36]. Taking advantage of the new formulation, we estimate the exit probability from one side of a domain and show that, interestingly, fast rare events can make a significant contribution to the exit time probability even when the scaling length is diffusive and the bulk of the distribution is Gaussian. We test our predictions by comparing them with extensive numerical simulations, with very good agreement. Our results show that the big jump principle can also be applied to first-passage problems, whereas previously it has only been used for probability densities. This broad applicability is based on the fact that the principle provides insight into the physical process that leads the walker to cover very large distances.

*The big jump for exit time probabilities.*—In a stochastic jump process, we call  $P_X(T)$  the probability density of reaching a distance  $X$  from the origin for the first time at time  $T$  without being absorbed by the starting point,  $x = 0$ . The characteristic length  $\ell(T) \sim T^r$  is defined by the scaling form of  $P_X(T)$  such that, for  $X \sim \ell(T)$ ,  $P_X(T) \sim f[X/\ell(T)]$  [24]. If there are independent jumps (renewals) in the process, drawn from a subexponential distribution [25,31], that allow reaching a distance  $X \gg \ell(T)$ , then the big jump principle is expected to hold [24]. In particular, the principle states that a process reaching a distance  $X \gg \ell(T)$  is triggered by a single very large jump, drawn from the subexponential distribution, while shorter distances of the order of  $\ell(T)$  are reached by processes involving multiple jumps. This physical insight suggests that we can focus on the big jump and for  $X \gg \ell(T)$  write the probability density function (PDF)  $P_X(T)$  as

$$P_X(T) \sim R(T_w) S(T_w) p_s(X). \quad (1)$$

Here,  $p_s(X)$  is the probability that the walker will reach a distance greater than  $X$  in a single jump;  $T_w$  is the time at which the big jump occurs, i.e.,  $T_w = T - \tau(X)$  where  $\tau(X)$  is the time at which the walker reaches the distance  $X$  in a single jump; the survival probability  $S(T_w)$  is the probability that the walker is not absorbed before time  $T_w$ ; and finally,  $R(T_w)$  is the rate at which jumps are made at time  $T_w$ . The  $p_s(X)$  and  $\tau(X)$  can be calculated directly by knowing the dynamics of the single jump, while the rate  $R(T_w)$  and the survival probability  $S(T_w)$  depend on the overall stochastic process. This factorization allows us to separate the calculation into quantities that depend on different characteristics of the stochastic process. We note that the renewals, which allow the application of the big jump, may not coincide with all stochastic events in the dynamics (see, for example, the difference between scattering events and renewals in the Lévy Lorentz gas).

*The discrete time random walk.*—In discrete time RWs [20], at each discrete time step the walker moves with probability  $1/2$  to the right or to the left, with the step length  $r$  drawn from the PDF  $p(r)$ . A finite second moment of  $p(r)$  implies standard diffusion while, if the second moment diverges, the walker performs a symmetric Lévy flight. We consider a  $p(r)$  with a power law decay at large  $r$ ,

$$p(r) \sim \frac{\alpha r_0^\alpha}{r^{1+\alpha}}, \quad (2)$$

emphasizing that the principle holds for any subexponential  $p(r)$  [26]. Here, time is defined by the number of steps  $n$  and  $x(n)$  is the position of the walker after  $n$  steps [ $x(0) = 0$ ]. This process features standard diffusion with a scaling length  $\ell(n) \sim n^{1/2}$  for  $\alpha > 2$  and a Lévy superdiffusive dynamics with  $\ell(n) \sim n^{1/\alpha}$ , for  $0 < \alpha < 2$ . We focus on the exit time probability  $P_X(n)$ , i.e., the probability that the walker at step  $n$  reaches a distance greater than  $X$  for the first time without being absorbed at the origin. The numerical results in Fig. 1 show  $P_X(n)$  as a function of the exit time  $n/X^2$ , with the time rescaling determined by the growth of  $\ell(n)$  [for  $\alpha = 2.5$ ,  $X \sim \ell(n) \sim n^{1/2}$ ]. The scaling form of  $P_X(n)$  and the analytical plot in the dashed line were obtained in [23] using a continuous limit. Very fast events, occurring on much shorter timescales (i.e., for  $n^{1/2} \ll X$ ) are also observed with non-negligible probability. These are the rare events that we want to describe.

The big jump principle states that the RW exits at time  $n$  at a distance  $X \gg \ell(n)$  if it is not absorbed at the origin, and if exactly at step  $n$  the walker makes a jump in the positive direction, of length greater than  $X$ . The distance covered in the previous steps can be neglected. Now, in discrete time RW, jumps are instantaneous, so  $\tau(X) = 0$  and  $T_w = T = n$ . Moreover,  $S(T_w) = S(n)$  can be obtained by the Sparre-Andersen theorem [37–39], which states that,




FIG. 1. The exit probability  $P_X(n)$  for a Lévy flight is plotted as a function of the rescaled exit time  $n/X^2$  ( $\alpha > 2$ ). We use a step length distribution  $p(r)$  with a cutoff at short distances  $r_0 = 1$ . The typical dynamics is diffusive and, at large number of steps  $n$ , converges slowly to the theoretical prediction with no free parameters [Eqs. (1),(5),(6) in [23]], which also provides the scaling factor  $X^3$ . On the left, very fast events are present with significant probability.

for a general jump processes in a semi-infinite line, the survival probability after  $n$  steps decays asymptotically as  $S(n) \sim (\pi n)^{-1/2}$ . The probability of making a jump greater than  $X$  is  $p_s(X) = \frac{1}{2} \int_X^\infty dr p(r)$  where the factor  $1/2$  accounts for the  $1/2$  probability of forward jumps. The renewal rate is trivially 1, so for  $n \gg 1$  and  $X \gg \ell(n)$  Eq. (1) reads

$$P_X(n) \sim S(n) \frac{1}{2} \int_X^\infty dr p(r) \sim \frac{1}{2(\pi n)^{1/2}} \frac{r_0^\alpha}{X^\alpha}. \quad (3)$$

Equation (3) is compared with numerical simulations in Fig. 2, showing an excellent agreement in the asymptotic regime at large  $X$ , both for the Lévy flights  $\alpha < 2$  and for normal diffusion  $\alpha > 2$ . On the other hand, the peaks of the distributions at shorter  $X$  occur when the system size is comparable to the scaling length  $\ell(n)$ , according to the analytical predictions in [23], as shown in Fig. 1 for  $\alpha > 2$ .

*The Lévy walk.*—The Lévy walk [20,40] is a continuous-time process in which a walker takes random steps of duration  $t$  at constant velocity  $v$ . In one dimension, each step is covered with equal probability in the positive and the negative direction, and the duration of the steps follows the PDF  $p(t)$ . We consider systems in which  $p(t)$  at large  $t$  is again a power law:

$$p(t) \sim \frac{\alpha t_0^\alpha}{t^{1+\alpha}}. \quad (4)$$

The scaling length of the stochastic process is  $\ell(t) \sim t^\gamma$ , with  $\gamma = 1$  for  $\alpha < 1$ ,  $\gamma = 1/\alpha$  for  $1 < \alpha < 2$ , and  $\gamma = 1/2$





FIG. 2. Exit probability  $P_X(n)$  for a Lévy flight.  $P_X(n)$  is multiplied by  $n^{1/2}$  and plotted as a function of the distance  $X$ . Upper panel  $\alpha = 1.2$ ; lower panel  $\alpha = 2.5$ . The dashed line shows the asymptotic analytical prediction in Eq. (3).

for  $\alpha > 2$  [40]. In addition, the single step contains another scaling length growing linearly with time due to the constant velocity motion. Therefore, for  $\alpha < 1$  in a single step the walker cannot cover a distance much larger than  $\ell(t)$  and the single big jump approach cannot be applied. On the other hand, for  $\alpha > 1$  as well as for other subexponential PDFs (e.g., Weibull), the principle can be used efficiently [24,26].

In Lévy walks the motion within a step is ballistic, so  $\tau(X) = X/v$  and  $T_w = T - X/v$ . The number of steps in a time  $T_w$  is  $n = T_w/\langle t \rangle$ , where  $\langle t \rangle = \int dt t' p(t')$  is the average duration of the steps. The Sparre-Andersen theorem can again be used and gives  $S(T_w) \sim (\pi T_w / \langle t \rangle)^{-1/2}$ , and the jump rate is constant,  $R(T_w) = 1/\langle t \rangle$  [24,26]. Finally,  $P_s(X) = \frac{1}{2} \int_{X/v}^\infty dt p(t)$  where again we take into account the ballistic motion within a step and the  $1/2$  probability of forward jumps. Summing up from Eq. (1) we obtain




FIG. 3. Exit probability  $P_X(T)$  for a Lévy walk.  $P_X(T)$  is multiplied by  $(vT)^{1/2+\alpha}$  and plotted as a function of the rescaled distance  $X/(vT)$ ,  $\alpha = 1.5$ . The continuous line shows the asymptotic analytical prediction in Eq. (5).

$$P_X(T) \sim \frac{1}{(vT)^{\alpha+1/2}} \frac{\langle t \rangle^{-1/2} t_0^\alpha}{2[\pi(1-y)]^{1/2} y^\alpha}, \quad (5)$$

where  $y = X/(vT) < 1$ . The ballistic motion in the big jump naturally introduces a new length scale,  $vT$ , which is the maximum distance that can be traveled in a time  $T$ . Notice that  $\langle t \rangle$  in Eq. (5) depends on the whole shape of  $p(t)$  and not on its far tail only. In the simulations we consider a distribution with a cutoff at  $t = t_0 = 1$ . In Fig. 3 we show that  $P_X(T)$  plotted against  $X/(vT)$  indeed converges to Eq. (5) at large times and distances, showing that the big jump estimate is correct.

*The Lévy Lorentz gas.*—The Lévy Lorentz gas [29,30] is a system of scatterers randomly placed in one dimension. The distances between scatterers are drawn again from a power law PDF  $p(l)$  at large  $l$ :

$$p(l) \sim \frac{\alpha l_0^\alpha}{l^{1+\alpha}}. \quad (6)$$

A RW is naturally defined on the Lévy Lorentz gas: the walker moves at constant speed  $v$  and is reflected with probability  $\epsilon$  ( $0 < \epsilon < 1$ ) when it hits a scatterer [41]. We focus on the case where the walker starts at  $t = 0$  in a scattering site placed at  $x = 0$  [29,30]. For this model, the PDF of the walker position has recently been studied [30,41–44]: the scaling length reads  $\ell(T) \sim T^{1/(1+\tilde{\alpha})}$ , where  $\tilde{\alpha} = \alpha$  if  $\alpha < 1$  and  $\tilde{\alpha} = 1$  if  $\alpha > 1$  [30] and the behavior of the PDF at large distances has been estimated using the big jump principle [24,30,45]. However, no results are yet known for the exit time probabilities.

Here,  $n(T_w)$  is the number of times (scattering events) in which a walker is reflected or transmitted by a scatterer. A renewal only occurs when a walker is transmitted for the first time by a scattering point, since in this case the distance





FIG. 4. Exit probability  $P_X(n)$  for a Lévy Lorentz gas.  $P_X(n)$  is multiplied by  $(vT)^{1+\alpha}$  and plotted as a function of the distance  $X/(vT)$ . Upper panel  $\alpha = 0.6$ ; lower panel  $\alpha = 1.5$ . The continuous line shows the asymptotic analytical prediction in Eq. (7). In this case a multiplicative constant has been optimized to reproduce numerical data. For large times the probabilities are small and the stochastic fluctuations become large, especially for  $\alpha > 1$ .

of the next scatterer is a random variable, independent of the RW history. Then, the big jump principle [24] applies and the walker can overcome the distance  $X \gg \ell(T)$  only by entering a region where the two scatterers are separated by a distance greater than  $X$ . Since the motion is ballistic, the large jump occurs at  $T_w = T - X/v$  and the rate of independent renewals is  $R(T_w) \sim T_w^{-[1/(1+\tilde{\alpha})]}$  [24]. On the other hand, the Sparre-Andersen theorem gives  $S(T_w) \sim n(T_w)^{-1/2}$ . Without considering the lattice spacing, the dynamics is a RW on a lattice, and the number of different scatterers visited in  $n(T_w)$  steps is  $v(T_w) \sim n(T_w)^{1/2}$ . Note that  $v(T_w)$  is also the number of scatterers within a distance  $\ell(T_w)$ ; so the result in [46] gives  $v(T_w) \sim \ell(T_w)^{\tilde{\alpha}} \sim T_w^{[\tilde{\alpha}/(1+\tilde{\alpha})]}$  and  $n(T_w) \sim T_w^{[2\tilde{\alpha}/(1+\tilde{\alpha})]}$  (for  $\alpha \geq 1$  we recover that the number of scattering events is proportional to the elapsed time). Since two

scatterers are separated by a distance greater than  $X$  with probability  $P_s(X) = \int_X^\infty dl p(l)$ , Eq. (1) gives

$$P_X(T) \sim \frac{1}{T_w^{1+\alpha}} \frac{1}{T_w^{\frac{\alpha}{1+\alpha}}} \int_X^\infty dl p(l) \sim \frac{1}{T^{\alpha+1}} \frac{1}{(1-y)y^\alpha}, \quad (7)$$

where again the rescaled distance  $y = X/(vT) \leq 1$  is induced by the ballistic motion in the single jump. Equation (7) contains a multiplicative constant since the rate and the number of jumps have been calculated only by scaling arguments. Figure 4 compares Eq. (7) to numerical simulation, showing an excellent agreement.

**Conclusions.**—We formulate a big jump approach for first-passage probabilities and use it to derive the analytical form of the tail in the exit distributions for three fundamental jump processes: the discrete time RW, the Lévy walk, and the Lévy-Lorentz gas. In practice, the physical information about how this single large jump unfolds is used to give an exact analytical estimate by separating the dynamics at the jump from the whole stochastic process, which can also be very complex. Our result shows that, in the presence of power law jump distributions, anomalous exit events can occur on timescales that are orders of magnitude smaller than typical exit times, even when the process is Gaussian. The result is heuristic and based on the application of the big jump principle, yet comparison with detailed numerical simulations suggests that the estimate is essentially correct, thus opening the way to a rigorous derivation. In this perspective, a fundamental question should be to identify what are the general characteristics of a stochastic process with subexponential distribution, allowing the application of the single big jump principle and in particular of Eq. (1). In its general form, the principle can also be applied to continuous-time processes and stochastic differential equation, provided that one can again identify the set of subexponential “jumps,” as has been successfully done in the case of the dynamics of cold atoms [24]. In that case, the big jump could be related to instantonic solutions [47–49]. Finally, we obtained our result for power law distributions but we expect such estimates to hold for a large class of subexponential jump processes [25,31], providing a boost to the study of rare events in first-passage probabilities.

We warmly thank Olivier Bénichou and Jérémie Klinger for interesting discussions. R. B. is supported by the Project funded under the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.4—Call for tender No. 3138 of 16/12/2021 of Italian Ministry of University and Research funded by the European Union—NextGenerationEU, Award Number: Project code CN00000023, Concession Decree No. 1033 of 17/06/2022 adopted by the Italian Ministry of University and Research, CUP D93C22000400001, “Sustainable Mobility Center” (CNMS), Spoke 9.

- [1] C. W. Gardiner, *Handbook of Stochastic Methods: for Physics, Chemistry and the Natural Sciences*, 3rd ed. (Springer, Berlin, 2004).
- [2] O. Bénichou, M. Coppey, M. Moreau, P.-H. Suet, and R. Voituriez, Optimal search strategies for hidden targets, *Phys. Rev. Lett.* **94**, 198101 (2005).
- [3] E. Agliari, R. Burioni, and A. Manzotti, Effective target arrangement in a deterministic scale-free graph, *Phys. Rev. E* **82**, 011118 (2010).
- [4] O. Bénichou, C. Loverdo, M. Moreau, and R. Voituriez, Intermittent search strategies, *Rev. Mod. Phys.* **83**, 81 (2011).
- [5] A. Godec and R. Metzler, Universal proximity effect in target search kinetics in the few-encounter limit, *Phys. Rev. X* **6**, 041037 (2016).
- [6] F. Thiel, E. Barkai, and D. A. Kessler, First detected arrival of a quantum walker on an infinite line, *Phys. Rev. Lett.* **120**, 040502 (2018).
- [7] H. Meyer and H. Rieger, Optimal non-Markovian search strategies with  $n$ -step memory, *Phys. Rev. Lett.* **127**, 070601 (2021).
- [8] R. Chichportiche and J.-P. Bouchaud, Some applications of first-passage ideas to finance, in *First-Passage Phenomena and Their Applications*, edited by R. Metzler, G. Oshanin, and S. Redner (World Scientific, Singapore, 2014), Chap. 1, pp. 447–476.
- [9] O. Pulkkinen and R. Metzler, Distance matters: The impact of gene proximity in bacterial gene regulation, *Phys. Rev. Lett.* **110**, 198101 (2013).
- [10] B. M. S. Arani, S. R. Carpenter, L. Lahti, E. H. van Nes, and M. Scheffer, Exit time as a measure of ecological resilience, *Science* **372**, eaay 4895 (2021).
- [11] S. Redner, *A Guide to First-Passage Processes* (Cambridge University Press, Cambridge, England, 2013).
- [12] O. Bénichou and R. Voituriez, From first-passage times of random walks in confinement to geometry-controlled kinetics, *Phys. Rep.* **539**, 225 (2014).
- [13] S. D. Lawley, Extreme first-passage times for random walks on networks, *Phys. Rev. E* **102**, 062118 (2020).
- [14] N. G. Van Kampen, *Stochastic Processes in Physics and Chemistry, Third Edition* (North-Holland Personal Library, Amsterdam, 2013).
- [15] A. V. Chechkin, R. Metzler, J. Klafter, and V. Y. Gonchar, Introduction to the theory of Lévy flights, in *Anomalous Transport*, edited by R. Klages, G. Radons, and I. M. Sokolov (John Wiley & Sons, Ltd., New York, 2008), Chap. 5, pp. 129–162.
- [16] O. K. Dudko, G. Hummer, and A. Szabo, Intrinsic rates and activation free energies from single-molecule pulling experiments, *Phys. Rev. Lett.* **96**, 108101 (2006).
- [17] U. M. B. Marconi, A. Puglisi, L. Rondoni, and A. Vulpiani, Fluctuation-dissipation: Response theory in statistical physics, *Phys. Rep.* **461**, 111 (2008).
- [18] S. Rotter and S. Gigan, Light fields in complex media: Mesoscopic scattering meets wave control, *Rev. Mod. Phys.* **89**, 015005 (2017).
- [19] G. H. Weiss, *Aspects and Applications of the Random Walk* (North Holland, Amsterdam, 1994).
- [20] J. Klafter and I. Sokolov, *First Steps in Random Walks: From Tools to Applications* (Oxford University Press, New York, 2011).

- [21] N. Levernier, O. Bénichou, and R. Voituriez, Universality classes of hitting probabilities of jump processes, *Phys. Rev. Lett.* **126**, 100602 (2021).
- [22] J. Klinger, R. Voituriez, and O. Bénichou, Splitting probabilities of symmetric jump processes, *Phys. Rev. Lett.* **129**, 140603 (2022).
- [23] J. Klinger, R. Voituriez, and O. Bénichou, Leftward, rightward, and complete exit-time distributions of jump processes, *Phys. Rev. E* **107**, 054109 (2023).
- [24] A. Vezzani, E. Barkai, and R. Burioni, Single-big-jump principle in physical modeling, *Phys. Rev. E* **100**, 012108 (2019).
- [25] S. Foss, D. Korshunov, and S. Zachary, *An Introduction to Heavy-Tailed and Subexponential Distributions* (Springer, New York, NY, 2013).
- [26] R. Burioni and A. Vezzani, Rare events in stochastic processes with sub-exponential distributions and the big jump principle, *J. Stat. Mech.* (2020) 034005.
- [27] A. Vezzani, E. Barkai, and R. Burioni, Rare events in generalized Lévy walks and the big jump principle, *Sci. Rep.* **10**, 2732 (2020).
- [28] W. Wang, A. Vezzani, R. Burioni, and E. Barkai, Transport in disordered systems: The single big jump approach, *Phys. Rev. Res.* **1**, 033172 (2019).
- [29] E. Barkai, V. Fleurov, and J. Klafter, One-dimensional stochastic Lévy-lorentz gas, *Phys. Rev. E* **61**, 1164 (2000).
- [30] R. Burioni, L. Caniparoli, and A. Vezzani, Lévy walks and scaling in quenched disordered media, *Phys. Rev. E* **81**, 060101(R) (2010).
- [31] C. M. Goldie and C. Klüppelberg, Subexponential distributions, in *A Practical Guide to Heavy Tails: Statistical Techniques and Applications*, edited by R. J. Adler, R. E. Feldman, and M. S. Taqqu (Birkhauser Boston Inc., Boston, USA, 1998), pp. 435–459.
- [32] M. A. Lomholt, K. Tal, R. Metzler, and J. Klafter, Lévy strategies in intermittent search processes are advantageous, *Proc. Natl. Acad. Sci. U.S.A.* **105**, 11055 (2008).
- [33] G. M. Viswanathan, M. G. E. da Luz, E. P. Raposo, and H. E. Stanley, *The Physics of Foraging: An Introduction to Random Searches and Biological Encounters*. (Cambridge University Press, Cambridge, England, 2011).
- [34] P. Barthélemy, J. Bertolotti, and D. S. Wiersma, A Lévy flight for light, *Nature (London)* **453**, 495 (2008).
- [35] J. Ślęzak and S. Burov, From diffusion in compartmentalized media to non-Gaussian random walks, *Sci. Rep.* **11**, 5101 (2021).
- [36] C. Bousige, P. Levitz, and B. Coasne, Bridging scales in disordered porous media by mapping molecular dynamics onto intermittent Brownian motion, *Nat. Commun.* **12**, 1043 (2021).
- [37] E. Sparre Andersen, On the fluctuations of sums of random variables II, *Math. Scand.* **2**, 194 (1954).
- [38] E. Sparre Andersen, On sums of symmetrically dependent random variables, *Scand. Actuarial J.* **1953**, 123 (1953).
- [39] S. N. Majumdar, P. Mounaix, and G. Schehr, Survival probability of random walks and Lévy flights on a semi-infinite line, *J. Phys. A* **50**, 465002 (2017).
- [40] V. Zaburdaev, S. Denisov, and J. Klafter, Lévy walks, *Rev. Mod. Phys.* **87**, 483 (2015).
- [41] R. Burioni, S. di Santo, S. Lepri, and A. Vezzani, Scattering lengths and universality in superdiffusive Lévy materials, *Phys. Rev. E* **86**, 031125 (2012).
- [42] R. Artuso, G. Cristadoro, M. Onofri, and M. Radice, Non-homogeneous persistent random walks and Lévy-Lorentz gas, *J. Stat. Mech.* (2018) 083209.
- [43] A. Bianchi, G. Cristadoro, M. Lenci, and M. Ligabò, Random walks in a one-dimensional Lévy random environment, *J. Stat. Phys.* **163**, 22 (2020).
- [44] A. Bianchi, M. Lenci, and F. Pène, Continuous-time random walk between Lévy-spaced targets in the real line, *Stoch. Proc. Appl.* **130**, 708 (2020).
- [45] M. Zamparo, Large fluctuations and transport properties of the Lévy-Lorentz gas, *Ann. Inst. Henri Poincaré, Probab. Stat.* **59**, 621 (2023).
- [46] C. W. J. Beenakker, C. W. Groth, and A. R. Akhmerov, Nonalgebraic length dependence of transmission through a chain of barriers with a Lévy spacing distribution, *Phys. Rev. B* **79**, 024204 (2009).
- [47] M. Alqahtani and T. Grafke, Instantons for rare events in heavy-tailed distributions, *J. Phys. A* **54**, 175001 (2021).
- [48] N. R. Smith, Anomalous scaling and first-order dynamical phase transition in large deviations of the Ornstein-Uhlenbeck process, *Phys. Rev. E* **105**, 014120 (2022).
- [49] Ž. Krajnik, E. Ilievski, and T. Prosen, Absence of normal fluctuations in an integrable magnet, *Phys. Rev. Lett.* **128**, 090604 (2022).