


Efficient Quantum Algorithms for Stabilizer Entropies

Tobias Haug^{1,2,*}, Soovin Lee^{2,†} and M. S. Kim²

¹Quantum Research Center, Technology Innovation Institute, Abu Dhabi, UAE

²Blackett Laboratory, Imperial College London, SW7 2AZ, United Kingdom

 (Received 4 June 2023; revised 8 May 2024; accepted 10 May 2024; published 13 June 2024)

Stabilizer entropies (SEs) are measures of nonstabilizerness or “magic” that quantify the degree to which a state is described by stabilizers. SEs are especially interesting due to their connections to scrambling, localization and property testing. However, applications have been limited so far as previously known measurement protocols for SEs scale exponentially with the number of qubits. Here, we efficiently measure SEs for integer Rényi index $n > 1$ via Bell measurements. The SE of N -qubit quantum states can be measured with $O(n)$ copies and $O(nN)$ classical computational time, where for even n we additionally require the complex conjugate of the state. We provide efficient bounds of various nonstabilizerness monotones that are intractable to compute beyond a few qubits. Using the IonQ quantum computer, we measure SEs of random Clifford circuits doped with non-Clifford gates and give bounds for the stabilizer fidelity, stabilizer extent, and robustness of magic. We provide efficient algorithms to measure Clifford-averaged $4n$ -point out-of-time-order correlators and multifractal flatness. With these measures we study the scrambling time of doped Clifford circuits and random Hamiltonian evolution depending on nonstabilizerness. Counterintuitively, random Hamiltonian evolution becomes less scrambled at long times, which we reveal with the multifractal flatness. Our results open up the exploration of nonstabilizerness with quantum computers.

DOI: [10.1103/PhysRevLett.132.240602](https://doi.org/10.1103/PhysRevLett.132.240602)

Stabilizer states and Clifford operations are essential to quantum information and quantum computing [1–3]. They are the cornerstone to run quantum algorithms on most fault-tolerant quantum computers, where Clifford operations are intertwined with non-Clifford gates [4,5]. To characterize the amount of non-Clifford resources needed to realize quantum states and operations the resource theory of nonstabilizerness has been put forward [6–14]. Stabilizer entropies (SEs) [15] are measures of nonstabilizerness with efficient algorithms for matrix product states [16–18] that have enabled the study of nonstabilizerness in many-body systems [16–24].

Recently, SEs have also been related to various important properties of quantum systems. SEs probe error-correction [25] and measurement-induced phase transitions [26,27], as well as relate to the entanglement spectrum [28] and property testing [29,30]. SEs are also connected to the participation entropy [31], which is helpful to understand Anderson [32] and many-body localization [33]. Further, recent works established a fruitful connection between out-of-time-order correlators (OTOCs) and

nonstabilizerness [30,34,35]. OTOCs describe scrambling in quantum systems [36,37]. However, OTOCs are challenging to measure directly and often require an inverse of the time evolution [38]. Higher-order OTOCs and nonstabilizerness have been related to quantum chaos [34] and state certification [30].

The aforementioned properties make SEs highly interesting for experimental studies of quantum computers and simulators. However, the progress has so far been limited as all previously known measurement protocols for SEs scale exponentially with the number of qubits [14,39].

Here, we efficiently measure SEs with integer index $n > 1$ on quantum computers and simulators via Bell measurements on two copies of N -qubit quantum states. Our algorithms are practical to implement with $O(n)$ copies and $O(nN)$ classical postprocessing time, where even n requires also access to the complex conjugate of the state. We devise an efficient protocol to measure Clifford-averaged multifractal flatness and $4n$ -point OTOCs where for odd n we do not require an inverse time evolution. We study the interplay of nonstabilizerness and scrambling and show that the number of Clifford gates needed for OTOCs to converge depends on the number of T gates. Further, we use the multifractal flatness to show that random Hamiltonian evolution stops being random for long evolution times. We also provide efficiently computable bounds to other nonstabilizerness monotones, which are otherwise intractable beyond a few qubits. Finally,

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

we measure the Tsallis SE on the IonQ quantum computer and demonstrate SEs as efficient bounds for the robustness of magic, stabilizer extent, and stabilizer fidelity. Our work introduces methods to uncover the key features that characterize the power of quantum computers and simulators.

SE.—For an N -qubit state $|\psi\rangle$, the Rényi- n SE is given by [15]

$$M_n(|\psi\rangle) = (1-n)^{-1} \ln \left(\sum_{\sigma \in \mathcal{P}} 2^{-N} \langle \psi | \sigma | \psi \rangle^{2n} \right), \quad (1)$$

where n is the index of the SE and \mathcal{P} is the set of 4^N Pauli strings. The Pauli strings are N -qubit tensor products $\sigma_r = \bigotimes_{j=1}^N \sigma_{r_{2j-1}r_{2j}}$ with $r \in \{0, 1\}^{2N}$, where $\sigma_{00} = I_1$, $\sigma_{01} = \sigma^x$, $\sigma_{10} = \sigma^z$, and $\sigma_{11} = \sigma^y$ with ℓ -qubit identity matrix I_ℓ and Pauli matrices σ^k , $k \in \{x, y, z\}$. M_n is a faithful measure of nonstabilizerness for pure states, i.e., $M_n(|\psi_{\text{STAB}}\rangle) = 0$ only for pure stabilizer states $|\psi_{\text{STAB}}\rangle$, and greater zero else [15]. Further, SEs are invariant under Clifford unitaries U_C with $M_n(U_C|\psi\rangle) = M_n(|\psi\rangle)$, where Clifford unitaries map any Pauli string σ to another Pauli string σ' via $U_C \sigma U_C^\dagger = \sigma'$. Further, M_n is additive with $M_n(|\psi\rangle \otimes |\phi\rangle) = M_n(|\psi\rangle) + M_n(|\phi\rangle)$. M_n for $n < 2$ is not a monotone under channels that can map a pure state to another pure state, while the case $n \geq 2$ remains an open problem.

Evaluating Eq. (1) requires an efficient measurement protocol for the n th moment of the Pauli spectrum

$$A_n(|\psi\rangle) = 2^{-N} \sum_{\sigma \in \mathcal{P}} \langle \psi | \sigma | \psi \rangle^{2n}, \quad (2)$$

which on first glance appears challenging due to the summation over exponentially many Pauli strings.

Algorithms.—We now provide two algorithms to efficiently measure $A_n(|\psi\rangle)$. First, we introduce Algorithm 1, which is efficient for odd $n > 1$. We write A_n as the expectation value of an observable $\Gamma_n^{\otimes N}$ acting on $2n$ copies of $|\psi\rangle$ via the replica trick [16]

$$A_n = 2^{-N} \sum_{\sigma \in \mathcal{P}} \langle \psi | \sigma | \psi \rangle^{2n} = \langle \psi | \otimes^{2n} \Gamma_n^{\otimes N} | \psi \rangle^{\otimes 2n}, \quad (3)$$

where $\Gamma_n = \frac{1}{2} \sum_{k=0}^3 (\sigma^k)^{\otimes 2n}$. For even $n > 1$, $2^{-N} \Gamma_n^{\otimes N}$ is a projector with two possible eigenvalues $\omega \in \{0, 2^N\}$ as shown in the Supplemental Material (SM) A [40]. In contrast, for odd $n > 1$ it is unitary and hermitian with eigenvalues $\omega \in \{-1, 1\}$. This fact was previously pointed out in Ref. [49] for stabilizer testing.

To measure the operator $\Gamma_n^{\otimes N}$ we transform the operator into a diagonal eigenbasis. We first recall the Bell transformation acting on two qubits $U_{\text{Bell}} = (H \otimes I_1) \text{CNOT}$, where $H = (1/\sqrt{2})(\sigma^x + \sigma^z)$ is the Hadamard gate, and $\text{CNOT} = \exp[i(\pi/4)(I_1 - \sigma^z) \otimes (I_1 - \sigma^x)]$. It turns out Γ_n is diagonalized by U_{Bell}

ALGORITHM 1. SE without complex conjugate.

Input: Integer $n > 1$; L repetitions;
Output: $A_n(|\psi\rangle)$

- 1 $A_n = 0$
- 2 **for** $k = 1, \dots, L$ **do**
- 3 **for** $j = 1, \dots, n$ **do**
- 4 Prepare $|\eta\rangle = U_{\text{Bell}}^{\otimes N} |\psi\rangle \otimes |\psi\rangle$
- 5 Sample in computational basis $r^{(j)} \sim |\langle r | \eta \rangle|^2$
- 6 **end**
- 7 $b = 1$
- 8 **for** $\ell = 1, \dots, N$ **do**
- 9 $\nu_1 = \bigoplus_{j=1}^n r_{2\ell-1}^{(j)}$; $\nu_2 = \bigoplus_{j=1}^n r_{2\ell}^{(j)}$
- 10 **if** n odd **then**
- 11 $b = b \cdot (-2\nu_1 \cdot \nu_2 + 1)$
- 12 **else**
- 13 $b = b \cdot 2(\nu_1 - 1) \cdot (\nu_2 - 1)$
- 14 **end**
- 15 **end**
- 16 $A_n = A_n + b/L$
- 17 **end**

$$A_n = \langle \psi | \otimes^{2n} \left\{ U_{\text{Bell}}^{\otimes n \dagger} \frac{1}{2} [(I_1 \otimes I_1)^{\otimes n} + (\sigma^z \otimes I_1)^{\otimes n} + (I_1 \otimes \sigma^z)^{\otimes n} + (-1)^n (\sigma^z \otimes \sigma^z)^{\otimes n}] U_{\text{Bell}}^{\otimes n} \right\}^{\otimes N} | \psi \rangle^{\otimes 2n}. \quad (4)$$

Algorithm 1 utilizes this fact to provide an unbiased estimator for A_n , where \bigoplus denotes binary addition. While Eq. (4) involves $2n$ copies of $|\psi\rangle$, the Bell transformation Eq. (4) can be written as tensor products. Thus, A_n is evaluated using only Bell measurements on two copies of the quantum state, which requires only a $2N$ -qubit quantum computer. Then, via postprocessing A_n is computed as the parity of odd and even qubit index Bell measurement outcomes as derived in SM B [40].

We apply Hoeffding's inequality to bound the number of copies as $C = O(n\Delta\omega_n\epsilon^{-2})$, where ϵ is the error and $\Delta\omega_n$ the range of eigenvalues of $\Gamma_n^{\otimes N}$. For odd $n > 1$, we have $\Delta\omega_n = 2$ and $C = O(n\epsilon^{-2})$. For even n , the eigenvalue spectrum of $\Gamma_n^{\otimes N}$ diverges and we require an exponential number of measurements. In SM C [40] we extend our algorithm to get gradients $\partial_k A_n$ via the shift rule for variational quantum algorithms.

Now, we provide Algorithm 2, which is efficient for any integer $n > 1$ but requires access to the complex conjugate $|\psi^*\rangle$. We rewrite the SE as a sampling problem,

$$A_n = \mathbb{E}_{\sigma \sim \Xi(\sigma)} [\langle \psi | \sigma | \psi \rangle^{2n-2}], \quad (5)$$

where $\Xi(\sigma) = 2^{-N} \langle \psi | \sigma | \psi \rangle^2$ is the probability distribution of Pauli strings σ . The circuit for the algorithm is shown in Fig. 1. First, we prepare $|\psi^*\rangle \otimes |\psi\rangle$ on the quantum




FIG. 1. Measurement protocol for Algorithm 2.

computer and transform into the Bell basis $|\eta\rangle = U_{\text{Bell}}^{\otimes N} |\psi^*\rangle \otimes |\psi\rangle$. Next, we sample from $2N$ -qubit state $|\eta\rangle$ in the computational basis, gaining outcome $\mathbf{r} \in \{0, 1\}^{2N}$. As shown in [50,51], we have $\Xi(\sigma_{\mathbf{r}}) = |\langle \mathbf{r} | \eta \rangle|^2$, where $|\mathbf{r}\rangle$ is the computational basis state corresponding to bitstring \mathbf{r} . Thus, sampling \mathbf{r} from $|\eta\rangle$ corresponds to sampling Pauli strings $\sigma_{\mathbf{r}} \sim \Xi(\sigma_{\mathbf{r}})$. Then, we perform $2n - 2$ measurements on $|\psi\rangle$ in the eigenbasis of the sampled $\sigma_{\mathbf{r}}$ and multiply the measured eigenvalues λ_k , gaining an unbiased estimator of $\langle \psi | \sigma_{\mathbf{r}} | \psi \rangle^{2n-2}$.

The measured eigenvalues $\prod_{k=1}^{2n-2} \lambda_k \in \{+1, -1\}$ have a range $\Delta\omega_n = 2$, thus according to Hoeffding's inequality we require at most $C = O(n\epsilon^{-2})$ copies of $|\psi\rangle$ and $O(\epsilon^{-2})$ copies of $|\psi^*\rangle$ for any integer $n > 1$. Note that $|\psi^*\rangle$ cannot be efficiently prepared in general with only black-box access to $|\psi\rangle$ [52–54]. However, when we have a circuit description of the unitary preparing the state, $|\psi^*\rangle$ is constructed by an element-wise conjugation of the coefficients of the unitary [55].

Tsallis SE.—We now define a measure of nonstabilizer-ness that we call the Tsallis- n SE [56],

$$T_n(|\psi\rangle) = -(1-n)^{-1} \left(1 - \sum_{\sigma \in \mathcal{P}} 2^{-N} \langle \psi | \sigma | \psi \rangle^{2n} \right). \quad (6)$$

They are a generalization of the linear SE T_2 [15] and the von Neumann SE $T_1 = M_1 = -2^{-N} \sum_{\sigma \in \mathcal{P}} \langle \psi | \sigma | \psi \rangle^2 \times \ln(\langle \psi | \sigma | \psi \rangle^2)$ [17]. T_n can be efficiently measured for integer $n > 1$ using our protocols. They are faithful measures of nonstabilizer-ness that are invariant under Clifford unitaries and related to Rényi SEs via $M_n = (1-n)^{-1} \ln[1 + (1-n)T_n]$. Tsallis SEs lack the additive property of the Rényi SE; however our numerics suggest that Tsallis SEs may be a strong monotone that is a not necessary but highly desirable property for resource measures [57]. Within extensive numerical optimization for $N \leq 6$ qubits we were unable to find states that could violate strong monotonicity for the Tsallis- n SE for $n \geq 2$ (see SM D [40]).

Note that measuring the Rényi SE $M_n \sim \ln(A_n)$ with precision ϵ_M requires $O[n \exp(M_n \epsilon_M^{-2})]$ samples due to the logarithm (see SM D [40]). Thus, M_n is efficiently measurable as long as $M_n = O[\log(N)]$.

ALGORITHM 2. SE with complex conjugate.

Input: Integer $n > 1$; L repetitions
Output: $A_n(|\psi\rangle)$

- 1 $A_n = 0$
- 2 **for** $k = 1, \dots, L$ **do**
- 3 Prepare $|\eta\rangle = U_{\text{Bell}}^{\otimes N} |\psi^*\rangle \otimes |\psi\rangle$
- 4 Sample $\mathbf{r} \sim |\langle \mathbf{r} | \eta \rangle|^2$
- 5 $b = 1$
- 6 **for** $\ell = 1, \dots, 2n - 2$ **do**
- 7 Prepare $|\psi\rangle$ and measure in eigenbasis of Paulistring $\sigma_{\mathbf{r}}$ for eigenvalue $\lambda \in \{+1, -1\}$
- 8 $b = b \cdot \lambda$
- 9 **end**
- 10 $A_n = A_n + b/L$
- 11 **end**

Clifford-averaged OTOCs.—We now show how to efficiently measure $4n$ -point OTOCs of unitary U averaged over the Clifford group. The $4n$ -point OTOC for N -qubit Pauli strings σ and σ' is given by [30]

$$\text{otoc}_{4n}(U, \sigma, \sigma') = (2^{-N} \text{tr}(\sigma U \sigma' U^\dagger))^{2n}. \quad (7)$$

We find that otoc_{4n} averaged over the group of Clifford unitaries \mathcal{C}_N can be related to SE of U , which we define via the Choi state $|U\rangle = I_N \otimes U|\Phi\rangle$, where $|\Phi\rangle = 2^{-N/2} \sum_{i=0}^{2^N-1} |i\rangle \otimes |i\rangle$. In particular, we have

$$\mathbb{E}_{U_C, U'_C \in \mathcal{C}_N} [\text{otoc}_{4n}(U_C U U'_C, \sigma, \sigma')] = \frac{A_n(|U\rangle) 4^N - 1}{(4^N - 1)^2}, \quad (8)$$

where the Pauli strings $\sigma, \sigma' \in \mathcal{P} \setminus \{I_N\}$ exclude the identity and the proof is found in SM K [40] using results of Ref. [30]. For odd $n > 1$, we can efficiently measure Eq. (8) via Algorithm 1. For even $n > 1$, we additionally require the complex conjugate $|U^*\rangle$ for Algorithm 2. The complex conjugate of the Choi state can be efficiently prepared with access to U^* or U^\dagger due to the ricochet property $|U^*\rangle = I_N \otimes U^*|\Phi\rangle = U^\dagger \otimes I_N|\Phi\rangle$ [55].

Multifractal flatness.—The participation entropy is given by $\mathcal{I}_q(|\psi\rangle) = \sum_k |\langle k | \psi \rangle|^{2q}$, where $|k\rangle$ are computational basis states, $q > 0$ and $0 \leq \mathcal{I}_q \leq 1$ [32]. The participation entropy quantifies the spread of the wave function over basis states, i.e., $\mathcal{I}_q = 1$ for computational basis states, while it is small when the state is delocalized over many computational basis states. The multifractal flatness $\mathcal{F}(|\psi\rangle) = \mathcal{I}_3(|\psi\rangle) - \mathcal{I}_2^2(|\psi\rangle)$ measures the flatness of the distribution $|\langle k | \psi \rangle|^2$, i.e., we have $\mathcal{F} = 0$ when the distribution $|\langle k | \psi \rangle|^2$ is constant over its support, else we have $\mathcal{F} > 0$. In particular, stabilizer states have $\mathcal{F} = 0$ [58].

Recently, \mathcal{F} averaged over \mathcal{C}_N has been proposed as $\bar{\mathcal{F}}$ [31]. This quantity describes the participation ratio

averaged over all possible choices of basis states. $\bar{\mathcal{F}}$ has been connected to SEs as follows [31]:

$$\bar{\mathcal{F}}(|\psi\rangle) = \mathbb{E}_{U_C \in \mathcal{C}_N} [\mathcal{F}(U_C|\psi\rangle)] = \frac{2[1 - A_2(|\psi\rangle)]}{(2^N + 1)(2^N + 2)}. \quad (9)$$

Thus, Algorithm 2 allows us to efficiently measure $\bar{\mathcal{F}}(|\psi\rangle)$ directly without the need of averaging over \mathcal{C}_N .

Bounds on nonstabilizerness.—We now provide efficient bounds on three magic monotones, namely the robustness of magic R [8], stabilizer extent ξ [59], and the stabilizer fidelity $F_{\text{STAB}} = \max_{|\phi\rangle \in \text{STAB}} |\langle \psi | \phi \rangle|^2$ [59] (see SM E [40]). Computing R , ξ , and F_{STAB} requires solving an optimization program over the set of pure N -qubit stabilizer states. As the number of stabilizer states scales as $O(2^{N^2})$, these three measures in general become numerically infeasible beyond five qubits [8,60].

Our algorithms provide efficient bounds for integer $n > 1$ (see [17] or SM E [40]),

$$R \geq \xi \geq F_{\text{STAB}}^{-1} \geq A_n^{-\frac{1}{2n}}. \quad (10)$$

The bound can be tightened for $n \geq \frac{1}{2}$ to $R \geq A_n^{[1/2(1-n)]}$ [8,15]. With methods from Ref. [49,61], we also prove a lower bound on F_{STAB} for $n > 1$ (see SM F [40]),

$$A_n^{\frac{1}{2n}} \geq F_{\text{STAB}} \geq \frac{A_n - 2^{1-n}}{1 - 2^{1-n}}. \quad (11)$$

The min-relative entropy of magic $D_{\min} = -\ln(F_{\text{STAB}})$ can be seen as the distance to the nearest stabilizer state. We now argue that D_{\min} , Rényi SEs with $n \geq 2$ and the recently introduced additive Bell magic \mathcal{B}_a [14] are closely related. In particular, we find evidence for respective upper and lower bounds independent of qubit number N (see SM G [40]). Via numerical optimization we find $1.7M_2 \gtrsim D_{\min} \gtrsim \frac{1}{4}M_2$ as well as $3.5M_2 \gtrsim \mathcal{B}_a \gtrsim 2.88M_2$ for at least $N \leq 4$, while similar bounds can also be found for larger n . Thus, D_{\min} , $M_{n \geq 2}$, and \mathcal{B}_a can be seen as measures of nonstabilizerness that relate to the distance to the nearest stabilizer state. In contrast, the robustness of magic R and stabilizer extent ξ relate to the degree a state can be approximated by a combination of stabilizer states. They belong to a different class of nonstabilizerness measures as they cannot be upper bounded with D_{\min} or M_n for $n > 1/2$ [17].

Demonstration.—We now study SEs with Bell measurements on the IonQ quantum computer [14] using Algorithm 1 in Fig. 2. We investigate random Clifford circuits U_C doped with N_T non-Clifford gates,

$$|\psi(N_T)\rangle = U_C^{(0)} \prod_{k=1}^{N_T} V_T^{(k)} U_C^{(k)} |0\rangle, \quad (12)$$

where $V_T^{(k)} = \exp[-i(\pi/8)\sigma_{g(k)}^z]$ is the T gate of the k th layer acting on a randomly chosen qubit $g(k)$. With




FIG. 2. Measurement of nonstabilizerness for quantum states generated by Eq. (12) with random Clifford circuits doped with N_T T gates on the IonQ quantum computer. (a) We show Tsallis SE T_3 with and without error mitigation as well as exact simulation. Dashed line is average value for Haar random states. Dots represent mean value and error bars the standard deviation taken over 6 random instances of the circuit. We have $N = 3$ qubits, 10^3 Bell measurements, and a measured depolarization error of $p \approx 0.1$. (b) We show upper and lower bounds on F_{STAB} via Eq. (11) evaluated using the error mitigated T_3 as blue and green dots as well as simulation of the bound as dashed lines. The orange dots show simulations of F_{STAB} .

increasing N_T these states transition from efficiently simulable stabilizer states to intractable quantum states [34,62]. To reduce noise, we compress the circuits into layered circuits composed of single-qubit operations and CNOT gates arranged in a nearest-neighbor chain configuration [14]. The state prepared by the quantum computer is not pure but degraded by noise. However, SEs are faithful measures of nonstabilizerness only for pure states. Using measurements on the noisy state, we mitigate A_n and T_n from measurements on noisy states by assuming a global depolarization error model (see SM H [40]).

In Fig. 2(a), we show T_3 with and without error mitigation for different N_T , where for each value we average over six random instances of Eq. (12). We find that the results on the IonQ quantum computer with error mitigation closely match the simulated values. The Tsallis SE is zero for $N_T = 0$, then increases with N_T until it converges to the average value of Haar random states indicated as black dashed line. In Fig. 2(b), we use the mitigated results for A_3 to compute upper and lower bounds for the stabilizer fidelity F_{STAB} using Eq. (11). The measured result indeed gives valid bounds of the exactly simulated F_{STAB} . We find that the upper bound is relatively tight, while the lower bound is nontrivial only for small N_T . In SM I [40], we provide additional results for the IonQ quantum computer on measures of nonstabilizerness. While our error mitigation scheme assumes global depolarization noise, it works well on actual quantum computers that have more complicated noise profiles. In SM J [40], we simulate our error mitigation scheme for various unital and nonunital noise models, and find very good performance. As SEs are moments of exponentially many Pauli strings, self-averaging effects may explain the good performance.




FIG. 3. (a) $\text{otoc}_8(U, \sigma_1^x, \sigma_1^x)$ against d layers of single-qubit Clifford gates and CNOT gates arranged in a nearest-neighbor chain, doped with N_T T gates and $N = 4$ qubits. Dashed line is the Clifford-averaged OTOC Eq. (8). (b) Multifractal flatness \mathcal{F} for evolution in time t with random Hamiltonian $\exp(-iH_{\text{GUE}}t)|0\rangle$. Dashed line is the Clifford-averaged multifractal flatness Eq. (9).

Scrambling.—We now study scrambling using the multifractal flatness \mathcal{F} and OTOCs. In Fig. 3(a), we show $\text{otoc}_8(U, \sigma_1^x, \sigma_1^x)$ against d layers of Clifford gates doped with N_T T gates. We find that the OTOC decreases with d , converging to a minimum once the unitary is fully scrambled. This minimum is given by the Clifford averaged OTOC Eq. (8) and depends on N_T . The d needed to converge depends on the number of T gates, where for $N_T = 0$ convergence is achieved for $d \sim 10$, while higher N_T requires larger d to converge. We observe similar convergence for \mathcal{F} and other OTOCs in SM K [40].

In Fig. 3(b), we study \mathcal{F} for the evolution of a state $|\psi(t)\rangle = \exp(-iH_{\text{GUE}}t)|0\rangle$ in time t using a random Hamiltonian H_{GUE} drawn from the Gaussian unitary ensemble (GUE). We observe that \mathcal{F} initially increases, reaching a maximum at $t \sim 1$. This is followed by a sudden dip to the Clifford-averaged multifractal flatness Eq. (9). This is hallmark of reaching deep thermalization or unitary design, where the system is indistinguishable from Haar-random dynamics [63]. Counterintuitively, for longer (exponential) times \mathcal{F} ramps up again, converging to a value above the Clifford-average. Here, the system stops being fully random due to dephasing of energy eigenvalues [63]. In SM K [40], we show how to measure \mathcal{F} and approximate GUE Hamiltonians using a Hamiltonian of random Pauli strings that can be implemented in experiment.

Discussions.—We show how to efficiently measure SEs with a cost independent of qubit number N , which is an exponential improvement over previous protocols [14,39]. For integer $n > 1$, our protocol is asymptotically optimal with the number of copies scaling as $O(n\epsilon^{-2})$ and the classical postprocessing time as $O(nN\epsilon^{-2})$ with additive error ϵ . The protocol is easy to implement using Bell measurements that have been demonstrated for quantum computers and simulators [64–66]. We note that our approach is distinct from the previously introduced Bell magic [14] as shown in SM L [40].

Our measurement protocol allows for efficient experimental characterization of different important properties of

quantum states. We demonstrate an efficient bound on nonstabilizerness monotones that otherwise are hard to compute beyond a few qubits. These monotones serve as lower bounds on state preparation complexity and characterize the runtime of classical simulation algorithms [8,59]. Further, we show how to efficiently measure Clifford-averaged $4n$ -point OTOCs. Our protocol has the advantage that it does not require implementing time reversal for odd $n > 1$, which can be a challenge [36]. Our protocol can measure higher order OTOCs that promise to reveal more features compared to the usually considered four-point OTOCs [67,68]. Our methods allow direct experimental study phase transitions in SE that have been found for purity testing [29] and quantum error correction [25]. Finally, we enable certification of magic gates in fault-tolerant quantum computers, where the SE could be directly evaluated from recent experimental data [69].

We use our methods to study scrambling in Clifford circuits doped with T gates and random Hamiltonian evolution. OTOCs not only measure scrambling, but also depend on nonstabilizerness in a nontrivial way [70]. We can disentangle these two effects by measuring the Clifford-averaged OTOC. We study when Clifford circuits doped with T gates become fully scrambled, revealing that the depth depends on the number of T gates. We also study the scrambling with random Hamiltonians. Notably, random Hamiltonian evolution deep thermalizes at intermediate times, becoming indistinguishable from Haar-random unitaries [63] that we observe via the convergence of the multifractal flatness to its Clifford-averaged value. Counterintuitively, the evolution becomes less random again for long times due to the dephasing of its energy eigenstates [63]. While nonlocal OTOCs and SEs lack clear signatures of this effect (see SE L), we find that multifractal flatness and same-site OTOCs exhibit clear gaps to their Clifford-average.

Future work could find efficient protocols for even n without the need of complex conjugation and tighten the lower bound of SEs for the stabilizer fidelity.

Note added.—Before acceptance of this Letter, the monotonicity of Rényi SE and strong monotonicity of Tsallis SE have been proven for $n \geq 2$ [72].

The code for this work is available on GitHub [71].

We thank Hyukjoon Kwon, Ludovico Lami, Lorenzo Leone, Salvatore F. E. Oliviero, Adam Taylor, and especially Lorenzo Piroli for inspiring discussions. We thank IonQ for providing quantum computing resources. This work is supported by a Samsung Global Research Cluster (GRC) project and the UK Hub in Quantum Computing and Simulation, part of the UK National Quantum Technologies Programme with funding from UKRI EPSRC Grant No. EP/T001062/1.

*Corresponding author: tobias.haug@u.nus.edu

†Corresponding author: soovinlee310@gmail.com

- [1] Daniel Gottesman, Stabilizer codes and quantum error correction, Caltech Ph.D., Ph.D. thesis, California Institute of Technology, 1997, [arXiv:quant-ph/9705052](https://arxiv.org/abs/quant-ph/9705052).
- [2] Peter W. Shor, Fault-tolerant quantum computation, in *Proceedings of 37th Conference on Foundations of Computer Science* (IEEE, New York, 1996), pp. 56–65.
- [3] A. Yu Kitaev, Fault-tolerant quantum computation by anyons, *Ann. Phys. (Amsterdam)* **303**, 2 (2003).
- [4] Sergey Bravyi and Alexei Kitaev, Universal quantum computation with ideal clifford gates and noisy ancillas, *Phys. Rev. A* **71**, 022316 (2005).
- [5] Earl T. Campbell, Barbara M. Terhal, and Christophe Vuillot, Roads towards fault-tolerant universal quantum computation, *Nature (London)* **549**, 172 (2017).
- [6] Earl T. Campbell, Catalysis and activation of magic states in fault-tolerant architectures, *Phys. Rev. A* **83**, 032317 (2011).
- [7] Victor Veitch, S. A. Hamed Mousavian, Daniel Gottesman, and Joseph Emerson, The resource theory of stabilizer quantum computation, *New J. Phys.* **16**, 013009 (2014).
- [8] Mark Howard and Earl Campbell, Application of a resource theory for magic states to fault-tolerant quantum computing, *Phys. Rev. Lett.* **118**, 090501 (2017).
- [9] Xin Wang, Mark M. Wilde, and Yuan Su, Quantifying the magic of quantum channels, *New J. Phys.* **21**, 103002 (2019).
- [10] Michael Beverland, Earl Campbell, Mark Howard, and Vadym Kliuchnikov, Lower bounds on the non-clifford resources for quantum computations, *Quantum Sci. Technol.* **5**, 035009 (2020).
- [11] Jiaqing Jiang and Xin Wang, Lower bound for the t count via unitary stabilizer nullity, *Phys. Rev. Appl.* **19**, 034052 (2023).
- [12] Zi-Wen Liu and Andreas Winter, Many-body quantum magic, *PRX Quantum* **3**, 020333 (2022).
- [13] Kaifeng Bu, Roy J. Garcia, Arthur Jaffe, Dax Enshan Koh, and Lu Li, Complexity of quantum circuits via sensitivity, magic, and coherence, [arXiv:2204.12051](https://arxiv.org/abs/2204.12051).
- [14] Tobias Haug and M. S. Kim, Scalable measures of magic resource for quantum computers, *PRX Quantum* **4**, 010301 (2023).
- [15] Lorenzo Leone, Salvatore F.E. Oliviero, and Alioscia Hamma, Stabilizer rényi entropy, *Phys. Rev. Lett.* **128**, 050402 (2022).
- [16] Tobias Haug and Lorenzo Piroli, Quantifying nonstabilizerness of matrix product states, *Phys. Rev. B* **107**, 035148 (2023).
- [17] Tobias Haug and Lorenzo Piroli, Stabilizer entropies and nonstabilizerness monotones, *Quantum* **7**, 1092 (2023).
- [18] Guglielmo Lami and Mario Collura, Nonstabilizerness via perfect pauli sampling of matrix product states, *Phys. Rev. Lett.* **131**, 180401 (2023).
- [19] Salvatore F.E. Oliviero, Lorenzo Leone, and Alioscia Hamma, Magic-state resource theory for the ground state of the transverse-field Ising model, *Phys. Rev. A* **106**, 042426 (2022).
- [20] Jovan Odavić, Tobias Haug, Gianpaolo Torre, Alioscia Hamma, Fabio Franchini, and Salvatore Marco Giampaolo, Complexity of frustration: A new source of non-local non-stabilizerness, *SciPost Phys.* **15**, 131 (2023).
- [21] Liyuan Chen, Roy J. Garcia, Kaifeng Bu, and Arthur Jaffe, Magic of random matrix product states, *Phys. Rev. B* **109**, 174207 (2024).
- [22] Kanato Goto, Tomoki Nosaka, and Masahiro Nozaki, Probing chaos by magic monotones, *Phys. Rev. D* **106**, 126009 (2022).
- [23] Stefano Piemontese, Tommaso Roscilde, and Alioscia Hamma, Entanglement complexity of the Rokhsar-Kivelson-Sign wavefunctions, *Phys. Rev. B* **107**, 134202 (2023).
- [24] Junjie Chen, Yuxuan Yan, and You Zhou, Magic of quantum hypergraph states, [arXiv:2308.01886](https://arxiv.org/abs/2308.01886).
- [25] Pradeep Niroula, Christopher David White, Qingfeng Wang, Sonika Johri, Daiwei Zhu, Christopher Monroe, Crystal Noel, and Michael J. Gullans, Phase transition in magic with random quantum circuits, [arXiv:2304.10481](https://arxiv.org/abs/2304.10481).
- [26] M. Bejan, C. McLauchlan, and B. Béri, Dynamical magic transitions in monitored clifford + t circuits, [arXiv:2312.00132](https://arxiv.org/abs/2312.00132).
- [27] Gerald E. Fux, Emanuele Tirrito, Marcello Dalmonte, and Rosario Fazio, Entanglement-magic separation in hybrid quantum circuits, [arXiv:2312.02039](https://arxiv.org/abs/2312.02039).
- [28] Emanuele Tirrito, Poetri Sonya Tarabunga, Guglielmo Lami, Titas Chanda, Lorenzo Leone, Salvatore F.E. Oliviero, Marcello Dalmonte, Mario Collura, and Alioscia Hamma, Quantifying non-stabilizerness through entanglement spectrum flatness, *Phys. Rev. A* **109**, L040401 (2024).
- [29] Lorenzo Leone, Salvatore F. E. Oliviero, Gianluca Esposito, and Alioscia Hamma, Phase transition in stabilizer entropy and efficient purity estimation, *Phys. Rev. A* **109**, 032403 (2024).
- [30] Lorenzo Leone, Salvatore F.E. Oliviero, and Alioscia Hamma, Nonstabilizerness determining the hardness of direct fidelity estimation, *Phys. Rev. A* **107**, 022429 (2023).
- [31] Xhek Turkeshi, Marco Schirò, and Piotr Sierant, Measuring nonstabilizerness via multifractal flatness, *Phys. Rev. A* **108**, 042408 (2023).
- [32] C. Castellani and L. Peliti, Multifractal wavefunction at the localisation threshold, *J. Phys. A* **19**, L429 (1986).
- [33] Jean-Marie Stéphan, Shunsuke Furukawa, Grégoire Misguich, and Vincent Pasquier, Shannon and entanglement entropies of one- and two-dimensional critical wave functions, *Phys. Rev. B* **80**, 184421 (2009).
- [34] Lorenzo Leone, Salvatore F.E. Oliviero, You Zhou, and Alioscia Hamma, Quantum chaos is quantum, *Quantum* **5**, 453 (2021).
- [35] Roy J Garcia, Kaifeng Bu, and Arthur Jaffe, Resource theory of quantum scrambling, *Proc. Natl. Acad. Sci. U.S.A.* **120**, e2217031120 (2023).
- [36] Shenglong Xu and Brian Swingle, Scrambling dynamics and out-of-time ordered correlators in quantum many-body systems: A tutorial, *PRX Quantum* **5**, 010201 (2024).
- [37] Neil Dowling, Pavel Kos, and Kavan Modi, Scrambling is necessary but not sufficient for chaos, *Phys. Rev. Lett.* **131**, 180403 (2023).
- [38] Jun Li, Ruihua Fan, Hengyan Wang, Bingtian Ye, Bei Zeng, Hui Zhai, Xinhua Peng, and Jiangfeng Du, Measuring out-of-time-order correlators on a nuclear magnetic resonance quantum simulator, *Phys. Rev. X* **7**, 031011 (2017).
- [39] Salvatore F. E. Oliviero, Lorenzo Leone, Alioscia Hamma, and Seth Lloyd, Measuring magic on a quantum processor, *npj Quantum Inf.* **8**, 148 (2022).

- [40] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.132.240602>, which includes Refs. [41–48], for detailed derivations, as well as additional experimental and numerical results.
- [41] Kosuke Mitarai, Makoto Negoro, Masahiro Kitagawa, and Keisuke Fujii, Quantum circuit learning, *Phys. Rev. A* **98**, 032309 (2018).
- [42] Juan Carlos Garcia-Escartin and Pedro Chamorro-Posada, Swap test and hong-ou-mandel effect are equivalent, *Phys. Rev. A* **87**, 052330 (2013).
- [43] Fernando G. S. L. Brandao, Aram W. Harrow, and Michał Horodecki, Local random quantum circuits are approximate polynomial-designs, *Commun. Math. Phys.* **346**, 397 (2016).
- [44] Frank Arute, Kunal Arya, Ryan Babbush, Dave Bacon, Joseph C. Bardin, Rami Barends, Rupak Biswas, Sergio Boixo, Fernando G. S. L. Brandao, David A. Buell *et al.*, Quantum supremacy using a programmable superconducting processor, *Nature (London)* **574**, 505 (2019).
- [45] Marko Žnidarič, Tomaž Prosen, and Peter Prelovšek, Many-body localization in the Heisenberg XXZ magnet in a random field, *Phys. Rev. B* **77**, 064426 (2008).
- [46] Martin Gärtner, Justin G. Bohnet, Arghavan Safavi-Naini, Michael L. Wall, John J. Bollinger, and Ana Maria Rey, Measuring out-of-time-order correlations and multiple quantum spectra in a trapped-ion quantum magnet, *Nat. Phys.* **13**, 781 (2017).
- [47] Xiao Mi, Pedram Roushan, Chris Quintana, Salvatore Mandra, Jeffrey Marshall, Charles Neill, Frank Arute, Kunal Arya, Juan Atalaya, Ryan Babbush *et al.*, Information scrambling in quantum circuits, *Science* **374**, 1479 (2021).
- [48] Philip Daniel Blocher, Serwan Asaad, Vincent Mourik, Mark A. I. Johnson, Andrea Morello, and Klaus Mølmer, Measuring out-of-time-ordered correlation functions without reversing time evolution, *Phys. Rev. A* **106**, 042429 (2022).
- [49] David Gross, Sepehr Nezami, and Michael Walter, Schur–Weyl duality for the clifford group with applications: Property testing, a robust hudson theorem, and de finetti representations, *Commun. Math. Phys.* **385**, 1325 (2021).
- [50] Ashley Montanaro, Learning stabilizer states by Bell sampling, [arXiv:1707.04012](https://arxiv.org/abs/1707.04012).
- [51] Ching-Yi Lai and Hao-Chung Cheng, Learning quantum circuits of some t gates, *IEEE Trans. Inf. Theory* **68**, 3951 (2022).
- [52] Yuxiang Yang, Giulio Chiribella, and Gerardo Adesso, Certifying quantumness: Benchmarks for the optimal processing of generalized coherent and squeezed states, *Phys. Rev. A* **90**, 042319 (2014).
- [53] Jisho Miyazaki, Akihito Soeda, and Mio Muraō, Complex conjugation supermap of unitary quantum maps and its universal implementation protocol, *Phys. Rev. Res.* **1**, 013007 (2019).
- [54] Tobias Haug, Kishor Bharti, and Dax Enshan Koh, Pseudorandom unitaries are neither real nor sparse nor noise-robust, [arXiv:2306.11677](https://arxiv.org/abs/2306.11677).
- [55] Sumeet Khatri, Ryan LaRose, Alexander Poremba, Lukasz Cincio, Andrew T. Sornborger, and Patrick J. Coles, Quantum-assisted quantum compiling, *Quantum* **3**, 140 (2019).
- [56] Constantino Tsallis, Possible generalization of Boltzmann-Gibbs statistics, *J. Stat. Phys.* **52**, 479 (1988).
- [57] Eric Chitambar and Gilad Gour, Quantum resource theories, *Rev. Mod. Phys.* **91**, 025001 (2019).
- [58] Piotr Sierant and Xhek Turkeshi, Universal behavior beyond multifractality of wave functions at measurement-induced phase transitions, *Phys. Rev. Lett.* **128**, 130605 (2022).
- [59] Sergey Bravyi, Dan Browne, Pádraic Calpin, Earl Campbell, David Gosset, and Mark Howard, Simulation of quantum circuits by low-rank stabilizer decompositions, *Quantum* **3**, 181 (2019).
- [60] Scott Aaronson and Daniel Gottesman, Improved simulation of stabilizer circuits, *Phys. Rev. A* **70**, 052328 (2004).
- [61] Sabee Grewal, Vishnu Iyer, William Kretschmer, and Daniel Liang, Improved stabilizer estimation via bell difference sampling, [arXiv:2304.13915](https://arxiv.org/abs/2304.13915).
- [62] Jonas Haferkamp, Random quantum circuits are approximate unitary t -designs in depth $O(nt^{5+o(1)})$, *Quantum* **6**, 795 (2022).
- [63] Jordan Cotler, Nicholas Hunter-Jones, Junyu Liu, and Beni Yoshida, Chaos, complexity, and random matrices, *J. High Energy Phys.* **11** (2017) 048.
- [64] Rajibul Islam, Ruichao Ma, Philipp M. Preiss, M. Eric Tai, Alexander Lukin, Matthew Rispoli, and Markus Greiner, Measuring entanglement entropy in a quantum many-body system, *Nature (London)* **528**, 77 (2015).
- [65] Hsin-Yuan Huang, Michael Broughton, Jordan Cotler, Sitan Chen, Jerry Li, Masoud Mohseni, Hartmut Neven, Ryan Babbush, Richard Kueng, John Preskill *et al.*, Quantum advantage in learning from experiments, *Science* **376**, 1182 (2022).
- [66] Dolev Bluvstein, Harry Levine, Giulia Semeghini, Tout T. Wang, Sepehr Ebadi, Marcin Kalinowski, Alexander Keesling, Nishad Maskara, Hannes Pichler, Markus Greiner *et al.*, A quantum processor based on coherent transport of entangled atom arrays, *Nature (London)* **604**, 451 (2022).
- [67] Daniel A. Roberts and Beni Yoshida, Chaos and complexity by design, *J. High Energy Phys.* **04** (2017) 121.
- [68] Roy J. Garcia, You Zhou, and Arthur Jaffe, Quantum scrambling with classical shadows, *Phys. Rev. Res.* **3**, 033155 (2021).
- [69] Dolev Bluvstein, Simon J. Evered, Alexandra A. Geim, Sophie H. Li, Hengyun Zhou, Tom Manovitz, Sepehr Ebadi, Madelyn Cain, Marcin Kalinowski, Dominik Hangleiter *et al.*, Logical quantum processor based on reconfigurable atom arrays, *Nature (London)* **626**, 1 (2023).
- [70] Lorenzo Leone, Salvatore F.E. Oliviero, and Alioscia Hamma, Isospectral twirling and quantum chaos, *Entropy* **23**, 1073 (2021).
- [71] Tobias Haug, Code for measuring stabilizer entropy on quantum computers (2024), https://github.com/txhaug/stabilizer_entropy.
- [72] Lorenzo Leone and Lennart Bittel, Stabilizer entropies are monotones for magic-state resource theory, [arXiv:2404.11652](https://arxiv.org/abs/2404.11652).