

**Erratum: Nuclear Charge Radii of Silicon Isotopes  
[Phys. Rev. Lett. 132, 162502 (2024)]**

Kristian König<sup>1</sup>, Julian C. Berengut, Anastasia Borschevsky, Alex Brinson, B. Alex Brown, Adam Dockery, Serdar Elhatisari, Ephraim Eliav, Ronald F. Garcia Ruiz, Jason D. Holt, Bai-Shan Hu, Jonas Karthein, Dean Lee, Yuan-Zhuo Ma, Ulf-G. Meißner, Kei Minamisono, Alexander V. Oleynichenko, Skyy V. Pineda, Sergey D. Prosnjak, Marten L. Reitsma, Leonid V. Skripnikov, Adam Vernon, and Andréi Zaitsevskii



(Received 29 June 2024; published 2 August 2024)

DOI: 10.1103/PhysRevLett.133.059901

In this Letter, we did not clarify the method used in the density functional theory calculations. The calculated charge radii using two functionals, NL3\* [1] and SVmin [2], shown in Figs. 2 and 5, were taken from the database available at [3]. This website was not cited in the published version of this Letter. We would like to thank Witold Nazarewicz for pointing this out and highlighting that the calculations published at [3] do not represent the latest results that can be obtained with density functional theory.

- [1] S. E. Agbemava, A. V. Afanasjev, D. Ray, and P. Ring, *Phys. Rev. C* **89**, 054320 (2014).
- [2] P. Klüpfel, P.-G. Reinhard, T. J. Bürenich, and J. A. Maruhn, *Phys. Rev. C* **79**, 034310 (2009).
- [3] <https://massexplorer.frib.msu.edu>.