

Geometric Formulation of Generalized Root- $T\bar{T}$ Deformations

H. Babaei-Aghbolagh^{1,2,*} Song He^{1,3,4,†} Tommaso Morone^{5,‡} Hao Ouyang^{1,§} and Roberto Tateo^{5,||}

¹Center for Theoretical Physics and College of Physics, Jilin University, Changchun 130012, China

²Department of Physics, University of Mohaghegh Ardabili, P.O. Box 179, Ardabil, Iran

³School of Physical Science and Technology, Ningbo University, Ningbo 315211, China

⁴Max Planck Institute for Gravitational Physics (Albert Einstein Institute), Am Mühlenberg 1, 14476 Golm, Germany

⁵Dipartimento di Fisica, Università di Torino, and INFN Sezione di Torino, Via P. Giuria 1, 10125, Torino, Italy



(Received 13 May 2024; accepted 16 August 2024; published 12 September 2024)

We develop a generic geometric formalism that incorporates both $T\bar{T}$ -like and root- $T\bar{T}$ -like deformations in arbitrary dimensions. This framework applies to a wide family of stress-energy tensor perturbations and encompasses various well-known field theories. Building upon the recently proposed correspondence between Ricci-based gravity and $T\bar{T}$ -like deformations, we further extend this duality to include root- $T\bar{T}$ -like perturbations. This refinement extends the potential applications of our approach and contributes to a deeper exploration of the interplay between stress tensor perturbations and gravitational dynamics. Among the various original outcomes detailed in this Letter, we have also obtained a deformation of the flat Jackiw-Teitelboim gravity action.

DOI: 10.1103/PhysRevLett.133.111602

Introduction—Recent studies concerning deformations of classical and quantum field theories have revealed rich connections between geometry and field dynamics. A prime example is that of $T\bar{T}$ deformations [1,2] of two-dimensional theories, driven by the irrelevant composite operator [3]

$$O_{T\bar{T}} = -\det(T_\nu^\mu) = \frac{1}{2}(T^{\mu\nu}T_{\mu\nu} - T_\mu^\mu T_\nu^\nu). \quad (1)$$

Despite being irrelevant, two-dimensional $T\bar{T}$ deformations remain well-controlled and even solvable at the quantum level. In the deformed theory, various quantities can be computed exactly from their counterparts in the original model. These include the finite-volume spectrum, the S matrix [1,2], the classical Lagrangian [4–6], and the torus partition function [7–10]. $T\bar{T}$ deformations connect with different topics in theoretical physics, such as string theory [11–14], holography [15–26], and quantum gravity [27–32]. We refer the reader to [33] for a pedagogical review on the subject.

Furthermore, the $T\bar{T}$ deformation lends itself to a number of geometric interpretations. It was proposed in

[7] that $T\bar{T}$ perturbing a theory is equivalent to coupling the original theory to a random geometry. $T\bar{T}$ deformations can also be interpreted as coupling the original theory to a flat space Jackiw-Teitelboim-like gravity [27,28], or equivalently, a topological gravity [29,34,35].

Another interesting deformation of two-dimensional field theories, driven by the so-called root- $T\bar{T}$ operator [36–38]

$$R = \sqrt{\frac{1}{2}T^{\mu\nu}T_{\mu\nu} - \frac{1}{4}T_\mu^\mu T_\nu^\nu}, \quad (2)$$

has recently attracted growing attention. While its quantum-mechanical definition remains uncertain, the root- $T\bar{T}$ perturbation displays some surprising properties at the classical level. Notably, it commutes with the $T\bar{T}$, allowing for their simultaneous activation, and for some integrable field theories, it preserves classical integrability [39]. The relation between root- $T\bar{T}$ deformed conformal field theories and ultrarelativistic (BMS₃) field theories was discussed in [40]. Finally, the connection between the root- $T\bar{T}$ deformation and the modified boundary conditions in the holographic dictionary was studied in [41]. These results were later employed to explore the modular properties of deformed holographic conformal field theories in [42,43].

In higher space-time dimensions, stress-energy tensor perturbations give rise to many interesting field theory models [4,7,36]. Extensive research has focused on $T\bar{T}$ -like and root- $T\bar{T}$ -like deformations of four-dimensional Maxwell's theory, exploring the relationship between electromagnetic duality invariance and stress tensor deformations [44–46]. The massive gravity formulation of duality-invariant nonlinear electrodynamics was studied

*Contact author: h.babaei@uma.ac.ir

†Contact author: hesong@jlu.edu.cn

‡Contact author: tommaso.morone@unito.it

§Contact author: haoouyang@jlu.edu.cn

||Contact author: roberto.tateo@unito.it

in [47] and, in three dimensions, it was shown that Born-Infeld theory displays a classical $T\bar{T}$ -like flow, connected to free Maxwell theory [48]. Furthermore, recent studies have explored nonlinear chiral two-form gauge theories in six dimensions as $T\bar{T}$ -like deformations [49].

This Letter introduces a generic geometric approach to encompass a broader class of stress-energy tensor perturbations. We show that a two-dimensional theory deformed by both $T\bar{T}$ and root- $T\bar{T}$ operators is dynamically equivalent to the undeformed theory coupled to a novel gravity action, at least at a classical level. We further generalize the geometric formulation to accommodate various deformations in higher dimensions. While prior studies have investigated geometric formulations of $T\bar{T}$ -like deformations in higher dimensions within the metric approach [36], our formulation is based on the description in terms of eigenvalues of the product of the vielbein. This approach allows us to study stress-energy tensor-related flows within a simple and elegant setup.

The recent work [50] has emphasized that a $T\bar{T}$ -type deformed matter action coupled with the standard Einstein-Hilbert action is equivalent to an undeformed matter theory coupled with a Ricci-based gravity theory [51]. Adopting this perspective, we incorporate this logic into our geometric formulations and introduce Ricci-based gravity actions linked with root- $T\bar{T}$ -like deformations. We develop

a unified framework for $T\bar{T}$ and root- $T\bar{T}$ perturbations in field theories across various space-time dimensions, which may extend the class of exact-solvability preserving deformations and deepen our understanding of the fundamental principles of quantum gravity and string theory.

Unified geometric formulation of $T\bar{T}$ and root- $T\bar{T}$ deformations in $d = 2$ —We denote by $S_0[\phi, e_\mu^a]$ an arbitrary undeformed action, where ϕ indicates a generic collection of matter fields and e_ν^a denotes an auxiliary dynamical zweibein. The associated auxiliary metric is $g_{\mu\nu} = \eta_{ab} e_\mu^a e_\nu^b$. We couple the auxiliary zweibein to a second zweibein f_μ^a , and the metric tensor $h_{\mu\nu} = \eta_{ab} f_\mu^a f_\nu^b$ associated to f_μ^a will eventually emerge as the metric of the manifold on which the deformed theory lives. It is convenient to define two Lorentz invariant variables

$$\begin{aligned} y_1 &= \text{tr}(e^{-1}f) = f_\mu^a e_\mu^a, \\ y_2 &= \text{tr}[(e^{-1}f)^2] = f_\mu^a e_\nu^b f_\nu^c e_\mu^d. \end{aligned} \quad (3)$$

We now show that the combination of $T\bar{T}$ and root- $T\bar{T}$ deformations can be generated from the action

$$S_{\gamma,\lambda}[\phi, e_\mu^a, f_\mu^a] = S_0[\phi, e_\mu^a] + S_{\text{grav}}[e_\mu^a, f_\mu^a], \quad (4)$$

where gravity action S_{grav} is

$$S_{\text{grav}}[e_\mu^a, f_\mu^a] = \frac{1}{2\lambda} \int d^2x \det e \left(2 + y_1^2 - y_2 - 2y_1 \cosh \frac{\gamma}{2} + 2\sqrt{2y_2 - y_1^2} \sinh \frac{\gamma}{2} \right). \quad (5)$$

The parameters λ and γ represent the $T\bar{T}$ and the root- $T\bar{T}$ perturbing parameters, respectively. When $\gamma = 0$, S_{grav} reduces to the topological gravity action associated to the $T\bar{T}$ deformation [29],

$$S_{\text{grav}}[e_\mu^a, f_\mu^a] = \frac{1}{2\lambda} \int d^2x \epsilon^{\mu\nu} \epsilon_{ab} (e_\mu^a - f_\mu^a)(e_\nu^b - f_\nu^b), \quad (6)$$

where ϵ is the Levi-Civita symbol. Our analysis will be carried out using the Euclidean signature, and the generalization to the Lorentzian signature is straightforward.

The deformed action can be obtained by extremizing (4) with respect to the auxiliary zweibein e_μ^a : performing the variation of (4) with respect to e_μ^a , we have

$$\det e (T^{[0]})_\nu^\mu \equiv \frac{\delta S_0}{\delta e_\mu^a} e_\nu^a = -\frac{\delta S_{\text{grav}}}{\delta e_\mu^a} e_\nu^a, \quad (7)$$

where $(T^{[0]})_\nu^\mu$ is the stress-energy tensor of the undeformed theory, computed with respect to e_μ^a . We denote the solution of the equation of motion by e_μ^{*a} . Note that (7) may admit multiple solutions e_μ^{*a} related to the choice of the branch for the square root of the root- $T\bar{T}$ operator: in this Letter, we ignore such branch ambiguities. However, in the quantum theory, we expect one should sum over contributions from

all branches in the path integral. The deformed field theory is obtained substituting e_μ^{*a} back into (4),

$$S_{\text{deformed}}[\phi, f_\mu^a] = S_{\gamma,\lambda}[\phi, e_\mu^{*a}, f_\mu^a]. \quad (8)$$

The stress-energy tensor of the deformed theory can be computed as

$$T_\nu^\mu \equiv \frac{1}{\det f} \frac{\delta S_{\gamma,\lambda}}{\delta f_\mu^a} f_\nu^a = \frac{1}{\det f} \frac{\delta S_{\text{grav}}}{\delta f_\mu^a} f_\nu^a \Big|_{e=e^*}, \quad (9)$$

where we have used the on-shell condition (7) for e_μ^a , so that $S_\lambda[\phi, e_\mu^{*a}, f_\mu^a]$ explicitly depends on f_μ^a alone. To simplify notation, we will not distinguish between e_ν^a and its on-shell value e_ν^{*a} , unless necessary. One can verify that the total action (4) obeys the following flow equations:

$$\frac{\partial S_{\gamma,\lambda}}{\partial \lambda} = - \int d^2x \det f \det(T_\nu^\nu), \quad (10)$$

$$\frac{\partial S_{\gamma,\lambda}}{\partial \gamma} = \int d^2x \det f \sqrt{\frac{1}{2} T_\nu^\mu T_\mu^\nu - \frac{1}{4} (T_\nu^\nu)^2}. \quad (11)$$

Therefore, the action (4) provides a geometric description of the combined $T\bar{T}$ and root- $T\bar{T}$ deformations. Since $S_{\gamma,\lambda}$ is defined as independent of the flow path, and since the

operators do not have explicit λ and γ dependence, the two types of deformations commute with each other.

As discussed in Sec. 2.3 of [29], one can translate the vielbein formulation to the metric formulation by choosing a gauge such that $e^{-1}f = \sqrt{g^{-1}h}$ by using the local Lorentz transformations of e and f , where we have omitted indices to simplify the notation. The validity of the flow equations (10) and (11) can also be verified in the metric formulation, and the details are shown in the Supplemental Material [52].

We now illustrate our methodology, starting from the simple undeformed action of a free scalar,

$$S_0[\phi, e_\mu^a] = \int d^2x \det e \left(\frac{1}{2} \eta^{ab} e_\mu^a e_\nu^b \partial_\mu \phi \partial_\nu \phi \right). \quad (12)$$

The solution of the equation of motion for e_ν^a is

$$\begin{aligned} e_\mu^{*a} &= \frac{1}{2} e^{\pm\gamma} \left(\frac{1}{\sqrt{1-4\lambda e^{\pm\gamma} X}} + 1 \right) f_\mu^a \\ &\mp \left(\frac{\sinh \frac{\gamma}{2} \pm 2\lambda e^{\pm\frac{\gamma}{2}} X}{X\sqrt{1-4\lambda e^{\pm\gamma} X}} + \frac{\sinh \frac{\gamma}{2}}{2X} \right) \eta^{ab} f_b^\nu \partial_\nu \phi \partial_\mu \phi, \end{aligned} \quad (13)$$

where $X = \frac{1}{2} \eta^{ab} f_a^\mu f_b^\nu \partial_\mu \phi \partial_\nu \phi$. Substituting the solution e_μ^{*a} into the action, we get

$$S_{\gamma,\lambda}[\phi, e_\nu^{*a}, f_\nu^a] = \int d^2x \frac{1 - \sqrt{1 - 4e^{\pm\gamma} \lambda X}}{2\lambda}, \quad (14)$$

which reproduces the result obtained in [38].

Uplift to higher dimensions—In this section, we uplift the geometric description to a family of deformations induced by functionals of the stress-energy tensor in higher dimensions. In d space-time dimensions, we consider the following general form for the gravity action:

$$S_{\text{grav}}[e_\mu^a, f_\mu^a] = \int d^d x \det e B(e^{-1}f), \quad (15)$$

where B is a Lorentz invariant function of $(e^{-1}f)_\nu^\mu$. Therefore, B depends only on the Lorentz invariant variables $y_n = \text{tr}[(e^{-1}f)^n]$, $n = 1, \dots, d$. For $n > d$, the y_n are not independent quantities. Since one can express B in terms of the variables y_n , the stress-energy tensor can be computed as

$$T \equiv \frac{1}{\det f} \frac{\delta S_{\text{grav}}}{\delta f} f = \sum_{n=1}^d \frac{n}{\det(e^{-1}f)} (e^{-1}f)^n \partial_{y_n} B, \quad (16)$$

where T denotes the matrix T_ν^μ . To construct higher-dimensional deforming operators, we need to compute Lorentz invariant functionals of the stress-energy tensor (16). Although we can express each invariant $\text{tr}(T^k)$ in terms of the y variables, this approach is quite inefficient in arbitrary dimensions, since there is no simple general formula for y_n when $n > d$.

However, assuming $e^{-1}f$ can be diagonalized by means of some matrix U as $e^{-1}f = U \text{diag}(\alpha_1, \dots, \alpha_d) U^{-1}$, the

function B can be expressed in terms of the eigenvalues α_i , and each y_n reduces to a power sum symmetric polynomial of the α_i . For this reason, working with the eigenvalues of $e^{-1}f$ proves to be a far more convenient strategy. The stress-energy tensor can be expressed as

$$T = \left(\prod_{k=1}^d \alpha_k \right)^{-1} U \text{diag}(\alpha_1 \partial_{\alpha_1} B, \dots, \alpha_d \partial_{\alpha_d} B) U^{-1}, \quad (17)$$

and

$$\text{tr}(T^k) = \left(\prod_{j=1}^d \alpha_j \right)^{-k} \sum_{i=1}^d (\alpha_i \partial_{\alpha_i} B)^k. \quad (18)$$

Expressing y_1 and y_2 in terms of eigenvalues of $e^{-1}f$, the two-dimensional gravity action (5) can be significantly simplified,

$$S_{\text{grav}}[e_\mu^a, f_\mu^a] = \frac{1}{\lambda} \int d^2x \det e (\alpha_1 - e^{\frac{\gamma}{2}})(\alpha_2 - e^{-\frac{\gamma}{2}}). \quad (19)$$

Note that S_{grav} is not a symmetric function of the eigenvalues because of the nonanalyticity of the root- $T\bar{T}$ operator. Exchanging two eigenvalues is equivalent to crossing a branch cut.

Motivated by the expression (19), we propose a generalization in arbitrary d space-time dimensions,

$$B = \frac{1}{\lambda^{\Sigma-1}} \prod_{k=1}^d (\alpha_k^{p_k} - \beta_k^{p_k})^{1/p_k}, \quad (20)$$

where λ and β_k are perturbing parameters, p_k are numbers characterizing the deformation, and $\Sigma = \sum_{k=1}^d p_k^{-1}$. When $d = 2$ and $p_k = 1$, the action (20) reduces to (19) if we identify $\beta_1 = e^{(\gamma/2)}$ and $\beta_2 = e^{-(\gamma/2)}$. We will show that the parameters λ and $\log \beta_k$ emerge as higher-dimensional analogs of the two-dimensional $T\bar{T}$ and root- $T\bar{T}$ deformation parameters, respectively. With the ansatz (20), the eigenvalues of the stress-energy tensor can be computed as

$$\tau_i = \alpha_i \partial_{\alpha_i} B \prod_{j=1}^d \alpha_j^{-1} = \frac{\alpha_i^{p_i}}{\alpha_i^{p_i} - \beta_i^{p_i}} B \prod_{j=1}^d \alpha_j^{-1}. \quad (21)$$

We also find

$$\prod_{k=1}^d \tau_k^{1/p_k} = \frac{1}{\lambda^{\Sigma-1}} B^{\Sigma-1} \left(\prod_{k=1}^d \alpha_k \right)^{1-\Sigma}. \quad (22)$$

Therefore, the flow equation for λ is

$$\frac{\delta S_{\text{grav}}}{\delta \lambda} = -(\Sigma - 1) \int d^d x \det f \left(\prod_{k=1}^d \tau_k^{1/p_k} \right)^{\frac{1}{\Sigma-1}}. \quad (23)$$

The operator on the right-hand side of the Eq. (23) is nonanalytic and not symmetric in terms of the stress-energy tensor eigenvalues τ_i . Particularly, when all p_k are equal to p , we obtain a $(\det T)^{[1/(d-p)]}$ deformation [4,7]. When $\Sigma = 2$, the deformation is of order $O(T^2)$. Let us now

consider the flow equation for the β parameters. We have

$$\beta_i \partial_{\beta_i} B - \beta_j \partial_{\beta_j} B = -(\tau_i - \tau_j) \prod_{k=1}^d \alpha_k. \quad (24)$$

Equation (24) suggests that the flow should be confined to the surface defined by $\prod_{k=1}^d \beta_k = 1$. Otherwise, the perturbing operator would explicitly depend on λ and β_k . The resulting flow equation is

$$\sum_{k=1}^d v^k \frac{\partial S_{\beta,\lambda}}{\partial \log \beta_k} = - \int d^d x \det f \left(\sum_{k=1}^d v^k \tau_k \right), \quad (25)$$

where v^k are constants satisfying $\sum_{k=1}^d v^k = 1$. Varying the β parameters on the surface $\prod_{k=1}^d \beta_k = 1$ leads to nonanalytic marginal deformations that commute with the $\prod_{k=1}^d \tau_k^{1/p_k}$ deformation. In two dimensions, the root- $T\bar{T}$ operator can be understood as the difference $\tau_1 - \tau_2$. However, explicitly expressing the difference between the τ_k in terms of $\text{tr}(T^j)$ is more difficult in higher dimensions. Let us now examine the initial conditions of the flow equations. When integrating out the auxiliary vielbein e_μ^a , one needs the equations of motion of e_μ^a ,

$$B(e^{-1}f)\delta_\nu^\mu - (e^{-1}f)_\alpha^\mu \frac{\partial B}{\partial (e^{-1}f)_\alpha^\nu} = - \frac{1}{\det e} \frac{\delta S_0}{\delta e_\mu^a} e_\nu^a. \quad (26)$$

The right-hand side is finite in the limit $\lambda \rightarrow 0$. Denoting the eigenvalues of $(T^{[0]})_\nu^\mu$ as $\tau_k^{[0]}$, the solution is

$$\alpha_j = \beta_j \left(\lambda (\tau_j^{[0]})^{-1} \left(\prod_{k=1}^d \tau_k^{[0]1/p_k} \right)^{\frac{1}{\Sigma-1}} + 1 \right)^{1/p_j}, \quad (27)$$

which implies that $\alpha_k = \beta_k + O(\lambda)$ when $\lambda \rightarrow 0$. When $\beta_k = 1$, we have $e_\mu^a \rightarrow f_\mu^a$ and the total action $S_{\beta,\lambda} = S_0[\phi, e_\mu^a] + S_{\text{grav}}[e_\mu^a, f_\mu^a]$ reduces to the original action $S_0[\phi, f_\mu^a]$. Equation (27) can be interpreted as the deformed boundary conditions in holography, formulated in terms of eigenvalue variables. In the Supplemental Material, we reproduce the root- $T\bar{T}$ deformed boundary conditions proposed in [41].

Examples—Several deformed field theories can be explored within this framework. A notable example is the ModMax theory [53] and its Born-Infeld-like (MMBI) extension [54]. The ModMax theory is a nonlinear conformal- and duality-invariant modification of Maxwell's theory. The MMBI extension maintains the duality invariance, and the action satisfies two commuting flow equations [5,55,56],

$$\frac{\partial \mathcal{L}_{\text{MMBI}}}{\partial \tilde{\lambda}} = \frac{1}{8} \left(T_{\mu\nu} T^{\mu\nu} - \frac{1}{2} T_\mu^\mu T_\nu^\nu \right), \quad (28)$$

$$\frac{\partial \mathcal{L}_{\text{MMBI}}}{\partial \tilde{\gamma}} = \frac{1}{2} \sqrt{T_{\mu\nu} T^{\mu\nu} - \frac{1}{4} T_\mu^\mu T_\nu^\nu}. \quad (29)$$

In the MMBI theory, the stress-energy tensor admits two degenerate eigenvalues τ_1 and τ_2 , each of multiplicity 2. Therefore, the following relations hold:

$$\text{tr}(T^2) - \frac{1}{2} (\text{tr}T)^2 = -4\sqrt{\det T} = -4\tau_1\tau_2, \quad (30)$$

$$\sqrt{\text{tr}(T^2) - \frac{1}{4} (\text{tr}T)^2} = \tau_1 - \tau_2. \quad (31)$$

Turning off the irrelevant deformation momentarily, one can notice that the flow (23) is satisfied in $d = 4$ by fixing $p_k = 2$ for each k , up to rescaling the irrelevant flow parameter. On the other hand, setting $\beta_1 = \beta_2 = e^\gamma = \beta_3^{-1} = \beta_4^{-1}$, the flow equation (25) can be identified with (29), up to a rescaling of γ . This shows that the MMBI flows (28) and (29) can be realized by coupling Maxwell's theory to the gravity action (20) with $d = 4$ and $p_k = 2$. In this case, (20) simply reduces to

$$S_{\text{grav}}[e_\mu^a, f_\mu^a] = \int d^4 x \det e \left[\frac{1}{\lambda} \prod_{k=1}^4 (\alpha_k^2 - \beta_k^2)^{1/2} \right]. \quad (32)$$

Note that the quantities α_k^2 represent the eigenvalues of $g^{\mu\rho} h_{\rho\nu}$: if we switch off the deformation induced by the β_k 's, the corresponding action can be expressed explicitly in terms of the metrics,

$$S_{\text{grav}}[h_{\mu\nu}, g_{\mu\nu}] = \frac{1}{\lambda} \int d^4 x \sqrt{\det(h_{\mu\nu} - g_{\mu\nu})}. \quad (33)$$

$T\bar{T}$ -like flows of six-dimensional two-form chiral theories were recently studied in [49]. In these models, T admits two degenerate eigenvalues of multiplicity 3 (throughout the flow), implying that our geometric construction can be straightforwardly implemented. Another example is the higher-dimensional generalized Nambu-Goto action of a self-interacting scalar field in d dimensions,

$$S_\lambda = \int d^d x \left[\frac{1 - \sqrt{1 - 2\lambda(1 - \lambda V) \partial^\mu \phi \partial_\mu \phi}}{\lambda(1 - \lambda V)} - \frac{2V}{1 - \lambda V} \right]. \quad (34)$$

The action (34) satisfies the flow equation (the $V = 0$ case has been proven in [49])

$$\frac{\partial S_\lambda}{\partial \lambda} = \int d^d x \left(\frac{1}{2d} \text{tr}(T^2) - \frac{1}{d^2} (\text{tr}T)^2 - \frac{d-2}{2\sqrt{d-1}d^{3/2}} \text{tr}(T) \sqrt{\text{tr}(T^2) - \frac{1}{d} (\text{tr}T)^2} \right). \quad (35)$$

The stress-energy tensor has a nondegenerate eigenvalue τ_1 and a degenerate eigenvalue τ_2 of multiplicity $d - 1$. In terms of eigenvalues, the deforming operator can be written as $-\frac{1}{2} \tau_1 \tau_2$. Therefore, such deformation can be achieved by setting $\beta_j = 1$, $p_1 = 1$ and $p_{k>2} = d - 1$ in (23). It was shown in [48] that the three-dimensional Born-Infeld theory also satisfies the flow equation (35), and one can show that T has a nondegenerate eigenvalue τ_1 and a

degenerate eigenvalue τ_2 of multiplicity 2, allowing for a similar description of the flow. Finally, alternative geometric formulations can be constructed when the theory's stress-energy tensor has two distinct degenerate eigenvalues, as described in the Supplemental Material.

Inclusion of dynamical gravity—In [50], it was pointed out that a $T\bar{T}$ -deformed matter action coupled to the Einstein-Hilbert action is equivalent to an undeformed matter theory coupled to a Ricci-based gravity. Continuing along the same line of thought, we now make the metric h dynamical and include the Einstein-Hilbert term within the first-order Palatini formalism. The total action is

$$S[h, g, \Gamma, \phi] = \frac{1}{2\kappa} \int d^d x \sqrt{\det h} h^{\mu\nu} R_{\mu\nu}(\Gamma) + \int d^d x \sqrt{\det g} B(g^{-1}h) + S_0[g, \phi], \quad (36)$$

where the Ricci curvature tensor is a functional of the connection

$$R_{\mu\nu}(\Gamma) = \partial_\alpha \Gamma_{\nu\mu}^\alpha - \partial_\nu \Gamma_{\alpha\mu}^\alpha + \Gamma_{\alpha\beta}^\alpha \Gamma_{\nu\mu}^\beta - \Gamma_{\nu\beta}^\alpha \Gamma_{\alpha\mu}^\beta. \quad (37)$$

The equations of motion for the connection $\Gamma_{\mu\nu}^\lambda$ lead to the compatibility conditions

$$\Gamma_{\mu\nu}^\lambda = \frac{1}{2} (h^{-1})^{\lambda\alpha} (\partial_\nu h_{\mu\alpha} + \partial_\mu h_{\alpha\nu} - \partial_\alpha h_{\mu\nu}). \quad (38)$$

Integrating out g in the action (36) we get

$$S[h, \Gamma, \phi] = \frac{1}{2\kappa} \int d^d x \sqrt{\det h} h^{\mu\nu} R_{\mu\nu}(\Gamma) + S_{\text{deformed}}[h, \phi], \quad (39)$$

which can be viewed as a deformed matter action S_{deformed} coupled to the standard Einstein-Hilbert action.

To obtain the Ricci-based gravity description, one can integrate out h in the action (36) and obtain

$$S[g, \Gamma, \phi] = \int d^d x \sqrt{\det g} \mathcal{L}(g^{-1}R) + S_0[g, \phi], \quad (40)$$

$$\mathcal{L} = \left[B(g^{-1}h) - \frac{1}{d-2} g^{\mu\alpha} h_{\alpha\nu} \frac{\partial B}{\partial (g^{\mu\beta} h_{\beta\nu})} \right] \Big|_{h=h^*(g)}, \quad (41)$$

which can be interpreted as an undeformed matter action coupled to a Ricci-based gravity theory $\mathcal{L}(g^{-1}R)$. This procedure yields a dynamical equivalence between an undeformed matter theory coupled to a Ricci-based gravity and a deformed theory coupled to standard general relativity.

It is, however, difficult to obtain an explicit expression for the Lagrangian $\mathcal{L}(g^{-1}R)$ associated with the B function given by (20) because the equations of motion of h are in general very complicated. In the Supplemental Material, we derive a flow equation for $\mathcal{L}(g^{-1}R)$,

$$\frac{\partial \mathcal{L}(\rho)}{\partial \lambda} = -(\Sigma - 1) \kappa^{-\frac{\Sigma}{\Sigma-1}} \left(\prod_{i=1}^d \alpha_i \right) \times \left(\prod_{k=1}^d \left(\alpha_k^{-2} \rho_k - \frac{1}{2} \sum_{j=1}^d \alpha_j^{-2} \rho_j \right)^{1/p_k} \right)^{\frac{1}{\Sigma-1}}, \quad (42)$$

where we express \mathcal{L} as a function of the eigenvalues ρ_k of $g^{-1}R$, and α_k can be determined through

$$\frac{\partial \mathcal{L}(\rho)}{\partial \rho_k} = \frac{1}{2\kappa \alpha_k^2} \prod_{i=1}^d \alpha_i. \quad (43)$$

The flow equation (42) allows computing the small λ expansion of $\mathcal{L}(g^{-1}R)$ (see the Supplemental Material).

Finally, in two dimensions, one can couple (5) to a Jackiw-Teitelboim-like gravity. We find that an undeformed matter theory coupled with a deformed Jackiw-Teitelboim-like gravity is dynamically equivalent to a deformed theory coupled to a Jackiw-Teitelboim-like gravity. The details are given in the Supplemental Material.

Including marginal flows in general deformations—One can consider Ricci-based gravity theories associated with more general deformations. For instance, the stress tensor deformation originating from Eddington-inspired Born-Infeld gravity [57] plays a role in $d = 4$ $T\bar{T}$ -like deformations of Abelian gauge theories [50]. For a stress tensor flow driven by an arbitrary operator $f(\tau_i)$ with parameter λ , the associated B function satisfies the flow equation

$$\frac{\partial B}{\partial \lambda} = f(\tau_i) \prod_{k=1}^d \alpha_k, \quad \tau_i = \alpha_i \partial_{\alpha_i} B \prod_{j=1}^d \alpha_j^{-1}. \quad (44)$$

One can also include marginal flows by replacing $\alpha_i \rightarrow \alpha_i/\beta_i$ with $\prod_{k=1}^d \beta_k = 1$ in the B function. The eigenvalues τ_i are modified as $\tau_i(\alpha_j) \rightarrow \tau_i(\alpha_j/\beta_j)$, and the form of the flow equation (44) remains unchanged. The flow equations associated with the β parameters are

$$\beta_i \frac{\partial B}{\partial \beta_i} - \beta_j \frac{\partial B}{\partial \beta_j} = \alpha_i \frac{\partial B}{\partial \alpha_i} + \alpha_j \frac{\partial B}{\partial \alpha_j} = -(\tau_i - \tau_j) \prod_{k=1}^d \alpha_k. \quad (45)$$

Therefore, it is possible to incorporate commutative marginal flows for any stress tensor deformation that admits a geometric realization. It follows from (42) that the associated Ricci-based gravity action should be modified as $\mathcal{L}(\rho_j) \rightarrow \mathcal{L}(\beta_j^{-2} \rho_j)$.

Conclusions—This Letter introduces a geometric formulation for the combination of $T\bar{T}$ and root- $T\bar{T}$ deformations in $d = 2$. We demonstrate that these deformations can be classically formulated by coupling the undeformed theory with a massive gravity action. Additionally, we extend the geometric framework to encompass various stress-energy tensor deformations in higher dimensions. These deformations are related to several well-known theories, including ModMax and its Born-Infeld-like extension. Furthermore, we study the Ricci-based gravities associated with such deformations. These findings might

have broad implications in key areas of string theory and holography, improving our understanding of the effects of stress tensor deformations. Note also that our approach appears suitable for studying various irrelevant and marginal deformations. However, challenges arise in finding exact solutions for more complex B functions in (15). Not all deformations will lead to explicit or unique solutions for the relevant constraints, which generalize (44) and (45).

There are several compelling avenues for future exploration stemming from our current work. A natural question is whether our formulation allows for the study of root- $T\bar{T}$ or more general deformations at the quantum level. The quantization of root- $T\bar{T}$ deformed theory poses a complex challenge, although some relevant progress has been made recently [58,59]. We anticipate that our formulation could offer insights into this intricate issue. Another avenue worth exploring is investigating the holographic dictionary of these deformations. Further, exploring the corresponding realization in celestial holography [60] would be valuable, as proposed in [61], which offers a potential avenue for constructing UV-complete gravity theories. The link between stress-energy flows and classical string or D-brane actions can provide insights into the UV completeness of deformed theories. Consequently, one can envisage constructing counterparts [61,62] in the framework of celestial holography to investigate their role in UV-complete theories.

Note added—After our work was submitted to arXiv, [63] appeared, also investigating the massive gravity description of the root- $T\bar{T}$ deformation and finding results consistent with ours. [63] also examined deformations with explicit λ dependence across various dimensions. In addition, an auxiliary field method to define integrable deformations of the principal chiral model was discussed in [64–66]. Exploring the potential connections between the two approaches remains an intriguing open problem.

Acknowledgments—The authors thank Dmitri Sorokin for his valuable discussions and comments. H. B.-A. expresses his gratitude to the Department of Physics and Astronomy “Galileo Galilei” at the University of Padova for their generous hospitality during the concluding phase of this project. S. H. acknowledges financial support from the Max Planck Partner Group, the Fundamental Research Funds for the Central Universities, and the Natural Science Foundation of China Grants No. 12075101, No. 12235016, No. 12347209, No. 12475053. H. O. is supported by the National Natural Science Foundation of China, Grant No. 12205115, and by the Science and Technology Development Plan Project of Jilin Province of China, Grant No. 20240101326JC. T. M. and R. T. received partial support from the INFN project SFT and the Prin (Progetti di rilevante interesse nazionale) Project No. 2022ABPBEY, with the title “Understanding quantum field theory through its deformations,” funded by the Italian Ministry of University and Research.

- [1] A. Cavaglià, S. Negro, I. M. Szécsényi, and R. Tateo, $T\bar{T}$ -deformed 2D quantum field theories, *J. High Energy Phys.* **10** (2016) 112.
- [2] F. A. Smirnov and A. B. Zamolodchikov, On space of integrable quantum field theories, *Nucl. Phys.* **B915**, 363 (2017).
- [3] A. B. Zamolodchikov, Expectation value of composite field $T\bar{T}$ in two-dimensional quantum field theory, [arXiv:hep-th/0401146](https://arxiv.org/abs/hep-th/0401146).
- [4] G. Bonelli, N. Doroud, and M. Zhu, $T\bar{T}$ -deformations in closed form, *J. High Energy Phys.* **06** (2018) 149.
- [5] R. Conti, L. Iannella, S. Negro, and R. Tateo, Generalised Born-Infeld models, Lax operators and the $T\bar{T}$ perturbation, *J. High Energy Phys.* **11** (2018) 007.
- [6] R. Conti, S. Negro, and R. Tateo, The $T\bar{T}$ perturbation and its geometric interpretation, *J. High Energy Phys.* **02** (2019) 085.
- [7] J. Cardy, The $T\bar{T}$ deformation of quantum field theory as random geometry, *J. High Energy Phys.* **10** (2018) 186.
- [8] S. Datta and Y. Jiang, $T\bar{T}$ deformed partition functions, *J. High Energy Phys.* **08** (2018) 106.
- [9] O. Aharony, S. Datta, A. Giveon, Y. Jiang, and D. Kutasov, Modular invariance and uniqueness of $T\bar{T}$ deformed CFT, *J. High Energy Phys.* **01** (2019) 086.
- [10] S. He, Y. Sun, and Y. X. Zhang, $T\bar{T}$ -flow effects on torus partition functions, *J. High Energy Phys.* **09** (2021) 061.
- [11] M. Baggio and A. Sfondrini, Strings on NS-NS backgrounds as integrable deformations, *Phys. Rev. D* **98**, 021902(R) (2018).
- [12] A. Dei and A. Sfondrini, Integrable S matrix, mirror TBA and spectrum for the stringy $\text{AdS}_3 \times S^3 \times S^3 \times S^1$ WZW model, *J. High Energy Phys.* **02** (2019) 072.
- [13] S. Chakraborty, A. Giveon, and D. Kutasov, $T\bar{T}$, $J\bar{T}$, $T\bar{J}$ and string theory, *J. Phys. A* **52**, 384003 (2019).
- [14] N. Callebaut, J. Kruthoff, and H. Verlinde, $T\bar{T}$ deformed CFT as a non-critical string, *J. High Energy Phys.* **04** (2020) 084.
- [15] L. McGough, M. Mezei, and H. Verlinde, Moving the CFT into the bulk with $T\bar{T}$, *J. High Energy Phys.* **04** (2018) 010.
- [16] P. Kraus, J. Liu, and D. Marolf, Cutoff AdS_3 versus the $T\bar{T}$ deformation, *J. High Energy Phys.* **07** (2018) 027.
- [17] W. Cottrell and A. Hashimoto, Comments on $T\bar{T}$ double trace deformations and boundary conditions, *Phys. Lett. B* **789**, 251 (2019).
- [18] M. Taylor, $T\bar{T}$ deformations in general dimensions, *Adv. Theor. Math. Phys.* **27**, 37 (2023).
- [19] T. Hartman, J. Kruthoff, E. Shaghoulian, and A. Tajdini, Holography at finite cutoff with a T^2 deformation, *J. High Energy Phys.* **03** (2019) 004.
- [20] P. Caputa, S. Datta, and V. Shyam, Sphere partition functions & cut-off AdS , *J. High Energy Phys.* **05** (2019) 112.
- [21] M. Guica and R. Monten, $T\bar{T}$ and the mirage of a bulk cutoff, *SciPost Phys.* **10**, 024 (2021).
- [22] D. J. Gross, J. Kruthoff, A. Rolph, and E. Shaghoulian, $T\bar{T}$ in AdS_2 and quantum mechanics, *Phys. Rev. D* **101**, 026011 (2020).
- [23] Y. Li and Y. Zhou, Cutoff AdS_3 versus $T\bar{T}$ CFT₂ in the large central charge sector: Correlators of energy-momentum tensor, *J. High Energy Phys.* **12** (2020) 168.
- [24] S. He, Y. Li, Y. Z. Li, and Y. Zhang, Holographic torus correlators of stress tensor in $\text{AdS}_3/\text{CFT}_2$, *J. High Energy Phys.* **06** (2023) 116.

- [25] S. He, Y. Z. Li, and Y. Zhang, Holographic torus correlators in AdS₃ gravity coupled to scalar field, *J. High Energy Phys.* **05** (2024) 254.
- [26] S. Ebert, Holographic renormalization group and stress tensor operators, [arXiv:2404.10190](https://arxiv.org/abs/2404.10190).
- [27] S. Dubovsky, V. Gorbenko, and M. Mirbabayi, Asymptotic fragility, near AdS₂ holography and $T\bar{T}$, *J. High Energy Phys.* **09** (2017) 136.
- [28] S. Dubovsky, V. Gorbenko, and G. Hernández-Chifflet, $T\bar{T}$ partition function from topological gravity, *J. High Energy Phys.* **09** (2018) 158.
- [29] A. J. Tolley, $T\bar{T}$ deformations, massive gravity and non-critical strings, *J. High Energy Phys.* **06** (2020) 050.
- [30] L. V. Iliesiu, J. Kruthoff, G. J. Turiaci, and H. Verlinde, JT gravity at finite cutoff, *SciPost Phys.* **9**, 023 (2020).
- [31] S. Okumura and K. Yoshida, $T\bar{T}$ -deformation and Liouville gravity, *Nucl. Phys.* **B957**, 115083 (2020).
- [32] S. Ebert, C. Ferko, H. Y. Sun, and Z. Sun, $T\bar{T}$ in JT gravity and BF gauge theory, *SciPost Phys.* **13**, 096 (2022).
- [33] Y. Jiang, A pedagogical review on solvable irrelevant deformations of 2D quantum field theory, *Commun. Theor. Phys.* **73**, 057201 (2021).
- [34] P. Caputa, P. Caputa, S. Datta, S. Datta, Y. Jiang, Y. Jiang, P. Kraus, and P. Kraus, Geometrizing $T\bar{T}$, *J. High Energy Phys.* **03** (2021) 140; **09** (2022) 110(E).
- [35] S. Dubovsky, S. Negro, and M. Porrati, Topological gauging and double current deformations, *J. High Energy Phys.* **05** (2023) 240.
- [36] R. Conti, J. Romano, and R. Tateo, Metric approach to a $T\bar{T}$ -like deformation in arbitrary dimensions, *J. High Energy Phys.* **09** (2022) 085.
- [37] H. Babaei-Aghbolagh, K. Babaei Velni, D. Mahdavian Yekta, and H. Mohammadzadeh, Marginal $T\bar{T}$ -like deformation and modified Maxwell theories in two dimensions, *Phys. Rev. D* **106**, 086022 (2022).
- [38] C. Ferko, A. Sfondrini, L. Smith, and G. Tartaglino-Mazzucchelli, Root- $T\bar{T}$ deformations in two-dimensional quantum field theories, *Phys. Rev. Lett.* **129**, 201604 (2022).
- [39] R. Borsato, C. Ferko, and A. Sfondrini, Classical integrability of root- $T\bar{T}$ flows, *Phys. Rev. D* **107**, 086011 (2023).
- [40] P. Rodríguez, D. Tempo, and R. Troncoso, Mapping relativistic to ultra/non-relativistic conformal symmetries in 2D and finite $\sqrt{T\bar{T}}$ deformations, *J. High Energy Phys.* **11** (2021) 133.
- [41] S. Ebert, C. Ferko, and Z. Sun, Root- $T\bar{T}$ deformed boundary conditions in holography, *Phys. Rev. D* **107**, 126022 (2023).
- [42] D. Tempo and R. Troncoso, Nonlinear automorphism of the conformal algebra in 2D and continuous $\sqrt{T\bar{T}}$ deformations, *J. High Energy Phys.* **12** (2022) 129.
- [43] J. Tian, T. Lai, and F. Omid, Modular transformations of on-shell actions of (root-)JT deformed holographic CFTs, [arXiv:2404.16354](https://arxiv.org/abs/2404.16354).
- [44] H. Babaei-Aghbolagh, K. Babaei Velni, D. M. Yekta, and H. Mohammadzadeh, $T\bar{T}$ -like flows in non-linear electrodynamic theories and S-duality, *J. High Energy Phys.* **04** (2021) 187.
- [45] H. Babaei-Aghbolagh, K. Babaei Velni, D. M. Yekta, and H. Mohammadzadeh, Manifestly SL(2, R) duality-symmetric forms in ModMax theory, *J. High Energy Phys.* **12** (2022) 147.
- [46] C. Ferko, S. M. Kuzenko, L. Smith, and G. Tartaglino-Mazzucchelli, Duality-invariant nonlinear electrodynamics and stress tensor flows, *Phys. Rev. D* **108**, 106021 (2023).
- [47] T. Flöss, D. Roest, and T. Westdijk, Non-linear electrodynamics from massive gravity, *J. High Energy Phys.* **02** (2024) 194.
- [48] C. Ferko, Y. Hu, Z. Huang, K. Koutrolikos, and G. Tartaglino-Mazzucchelli, $T\bar{T}$ -like flows and 3d nonlinear supersymmetry, *SciPost Phys.* **16**, 038 (2024).
- [49] C. Ferko, S. M. Kuzenko, K. Lechner, D. P. Sorokin, and G. Tartaglino-Mazzucchelli, Interacting chiral form field theories and $T\bar{T}$ -like flows in six and higher dimensions, *J. High Energy Phys.* **05** (2024) 320.
- [50] T. Morone, S. Negro, and R. Tateo, Gravity and $T\bar{T}$ flows in higher dimensions, *Nucl. Phys.* **B1005**, 116605 (2024).
- [51] G. J. Olmo and D. Rubiera-Garcia, Some recent results on Ricci-based gravity theories, *Int. J. Mod. Phys. D* **31**, 2240012 (2022).
- [52] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.133.111602> for the (1) details of the metric formulation, (2) details of the holographic boundary condition, (3) an alternative geometric formulation of the Born-Infeld-like extension of the ModMax theory and generalized Nambu-Goto, (4) the derivation of the Lagrangian of the Ricci-based gravity and (5) details of the deformed Jackiw-Teitelboim-like gravity.
- [53] I. Bandos, K. Lechner, D. Sorokin, and P. K. Townsend, A non-linear duality-invariant conformal extension of Maxwell's equations, *Phys. Rev. D* **102**, 121703(R) (2020).
- [54] I. Bandos, K. Lechner, D. Sorokin, and P. K. Townsend, On p-form gauge theories and their conformal limits, *J. High Energy Phys.* **03** (2021) 022.
- [55] H. Babaei-Aghbolagh, K. B. Velni, D. M. Yekta, and H. Mohammadzadeh, Emergence of non-linear electrodynamic theories from $T\bar{T}$ -like deformations, *Phys. Lett. B* **829**, 137079 (2022).
- [56] C. Ferko, L. Smith, and G. Tartaglino-Mazzucchelli, On current-squared flows and ModMax theories, *SciPost Phys.* **13**, 012 (2022).
- [57] J. Beltran Jimenez, L. Heisenberg, G. J. Olmo, and D. Rubiera-Garcia, Born-Infeld inspired modifications of gravity, *Phys. Rep.* **727**, 1 (2018).
- [58] S. Ebert, C. Ferko, C. L. Martin, and G. Tartaglino-Mazzucchelli, Flows in the space of interacting chiral boson theories, *Phys. Rev. D* **110**, 046005 (2024).
- [59] L. Hadasz and R. von Unge, Defining root- $T\bar{T}$, [arXiv:2405.17945](https://arxiv.org/abs/2405.17945).
- [60] S. Pasterski, S. H. Shao, and A. Strominger, Flat space amplitudes and conformal symmetry of the celestial sphere, *Phys. Rev. D* **96**, 065026 (2017).
- [61] S. He, P. Mao, and X. C. Mao, $T\bar{T}$ deformed soft theorem, *Phys. Rev. D* **107**, L101901 (2023).

- [62] S. He, P. Mao, and X. C. Mao, Loop corrections versus marginal deformation in celestial holography, [arXiv:2307.02743](#).
- [63] E. Tsolakidis, Massive gravity generalization of $T\bar{T}$ deformations, [arXiv:2405.07967](#).
- [64] C. Ferko and L. Smith, An infinite family of integrable sigma models using auxiliary fields, [arXiv:2405.05899](#).
- [65] D. Bielli, C. Ferko, L. Smith, and G. Tartaglino-Mazzucchelli, T-duality and $T\bar{T}$ -like deformations of sigma models, [arXiv:2407.11636](#).
- [66] D. Bielli, C. Ferko, L. Smith, and G. Tartaglino-Mazzucchelli, Integrable higher-spin deformations of sigma models from auxiliary fields, [arXiv:2407.16338](#).