



FIG. 4. The differential rho-production cross section  $d\sigma(1.7\text{-BeV}/c \pi^- + p \rightarrow \rho^0 + n)/dt$  versus four-momentum transfer between incoming proton and outgoing neutron. This curve is taken from data supplied by Fickinger, Robinson, and Salant.<sup>2</sup>

differential cross sections, offers useful parametrization of the data in terms of radii and volumes of interaction.

We are greatly indebted to Dr. E. O. Salant for lending the film, and to Serge Paul-Emile and to our technicians for their indispensable help.

\*Research supported by the U. S. Atomic Energy Commission Contract No (11-1)-1537.

<sup>1</sup>D. D. Allen, G. Godden, G. P. Fisher, J. B. Kopelman, L. Marshall, and R. Sears, *Phys. Letters* **21**, 468 (1966). (We estimate that at most 4% of the events in the  $\pi^-p\pi^0$  final state have been designated as elastics, and less than 1% of the  $\pi^-p\pi^0$  events used in this report are due to contamination from other final states.)

<sup>2</sup>W. J. Fickinger, D. K. Robinson, and E. O. Salant, *Phys. Rev. Letters* **10**, 457 (1963); and Proceedings of the Athens Conference on Resonant Particles, Ohio University, Athens, Ohio, 1965 (to be published).

<sup>3</sup>Aachen-Berlin-CERN Collaboration, CERN Report No. CERN/TC/P-65-23, 1965, to be published; K. J. Foley, S. J. Lindenbaum, W. A. Love, S. Ozaki, J. J. Russell, and L. C. L. Yuan, *Phys. Rev. Letters* **10**, 376 (1963); **11**, 425 (1963). S. Brandt, V. T. Cocconi, D. R. O. Morrison, A. Wroblewski, P. Fleury, G. Kayas, F. Muller, and C. Pelletier, *Phys. Rev. Letters* **10**, 413 (1963); I. Mannelli, A. Bigi, R. Carrara, M. Wahlig, and L. Sodickson, *Phys. Rev. Letters* **14**, 408 (1965); A. V. Stirling, P. Sonderegger, J. Kirz, P. Falk-Vairant, O. Guisan, C. Bruneton, P. Borgeaud, M. Yvert, J. P. Guillaud, C. Caverzasio, and B. Amblard, *Phys. Rev. Letters*, **14**, 763 (1965).

<sup>4</sup>N. Byers and C. N. Yang, *Phys. Rev.* **142**, 976 (1966), and see references quoted in this paper.

<sup>5</sup>C. Iddings and L. Marshall, to be published.

<sup>6</sup>J. D. Jackson, private communication to G. P. Fisher.

<sup>7</sup>K. Gottfried and J. D. Jackson, *Nuovo Cimento* **33**, 309 (1964); **34**, 735 (1964).

<sup>8</sup>I. Derado, V. P. Kenney, and W. D. Shephard, *Phys. Rev. Letters* **13**, 505 (1964).

<sup>9</sup>D. Barge, W. Chu, L. Leipuner, R. Crittenden, H. J. Martin, F. Ayer, L. Marshall, A. C. Li, W. Kernan, and M. L. Stevenson, *Phys. Rev. Letters* **13**, 69 (1964).

<sup>10</sup>A. Fridman, O. Renary, A. Michalon, B. Schiby, R. Strub, and G. Zech, "Elastic  $K^-$  Proton Interaction at 1.45 GeV/c and Comparison with Absorptive Models" (to be published).

## CURRENT COMMUTATORS, REPRESENTATION MIXING, AND MAGNETIC MOMENTS\*

Haim Harari†

Stanford Linear Accelerator Center, Stanford University, Stanford, California

(Received 22 April 1966)

A detailed analysis of the various sum rules which have been recently derived from the chiral  $U(3) \otimes U(3)$  algebra of currents<sup>1</sup> indicates that the exact sum rules may be approximated by sums over a few intermediate states which fall into a relatively simple reducible representation of the current algebra. In a previous paper<sup>2</sup> (hereafter denoted by I) we have shown

that the positive-helicity state of the nucleon can be properly described as having components in the  $\{(6, 3)_{L_z=0}\}$ ,  $\{(3^*, 3)_{L_z=0}\}$ , and  $\{(3, 3)_{L_z=1}\}$  representations of  $U(3) \otimes U(3)$ , and that by adjusting one free mixing angle one can then correctly predict the experimental values of  $G_A$ , the axial vector coupling constant in  $\beta$  decay;  $G^*$ , the strength of the axial-vector tran-

sition between the nucleon and the  $N^*(1238)$  resonance; and the  $d/f$  ratio for the axial-vector current between states of the baryon octet.<sup>3</sup> In the present paper we show that the same assumptions and the same mixing angle lead, in addition, to a prediction for the ratio between the anomalous magnetic moment of the neutron and the strength of the magnetic transition between the nucleon and  $N^*(1238)$ . The predicted value for this ratio is in excellent agreement with the experimental data. Furthermore, by adjusting one additional free parameter, we can obtain the correct ratio between the anomalous moments of the proton and the neutron.

Following the approach we used in I, we consider the algebra<sup>4</sup>  $U(3) \otimes U(3) \otimes U(3) \otimes U(3)$  generated by the equal-time commutators of the  $z$  and  $t$  components of the vector and axial-vector currents, evaluated between particle states moving with infinite momentum in the  $z$  direction. Since at infinite momentum the matrix elements of the  $z$  components are equal to those of the time components (both for the vector and the axial-vector current), it is sufficient to discuss an algebra  $U(3) \otimes U(3)$ , which can then be identified either as the chiral or as the collinear current algebra.<sup>4</sup> In view of the difficulties which follow from the assignment of the positive- and negative-helicity states of the nucleon to the  $(\underline{6}, \underline{3})$  and  $(\underline{3}, \underline{6})$  representations, respectively, it has been suggested that an appreciable amount of representation mixing is present. This was mainly motivated by the following observations:

(1) The analysis of the Adler-Weisberger sum rules for both the strangeness-conserving<sup>5</sup> and the strangeness-changing<sup>6</sup> currents clearly indicates that the decuplet states do not saturate the integrals of meson-nucleon cross sections and that the contributions of higher resonances (mostly with negative parities) cannot be neglected. Furthermore, the decuplet dominance assumption turns out to be inadequate in a few other cases,<sup>7-9</sup> although it is not clear, in these cases, whether the exact sum rules are verified.<sup>10</sup>

(2) If the nucleon belongs to any pure  $U(6)$  or  $U(3) \otimes U(3)$  representation having  $L_z = 0$ , its anomalous magnetic moment is predicted to vanish<sup>4</sup> ( $L_z$  is defined here as  $J_z - S_z$  where  $J_z$  and  $S_z$  are the  $z$  components of the total angular momentum and the "intrinsic quark spin," respectively). The prediction is easily obtained by observing that the anomalous-moment oper-

ator transforms under the current algebra like the  $z$  (or time) component of the electromagnetic current, but it changes  $L_z$  by one unit and therefore cannot connect two  $L_z = 0$  states. It is interesting to add, in this connection, that the sum rule derived by Fubini, Furlan, and Rossetti<sup>9</sup> for the anomalous moments of the baryons leads to  $\mu_A(B) = 0$  if we require  $SU(3)$  symmetry and decuplet dominance.<sup>11</sup> This result is, of course, intimately related to the observation that any pure  $L_z = 0$  representation for the baryon octet [such as the  $(\underline{6}, \underline{3})$  of  $U(3) \otimes U(3)$  or the  $\underline{56}$  of  $SU(6)$ ] implies a vanishing  $\mu_A$ .

According to the results of I, the positive-helicity state of the nucleon is given by

$$|N, J_z = \frac{1}{2}\rangle = \cos\theta |(\underline{6}, \underline{3})_{L_z = 0}\rangle + \sin\theta \left\{ \left(\frac{1}{3}\right)^{1/2} |(\underline{3}^*, \underline{3})_{L_z = 0}\rangle - \left(\frac{2}{3}\right)^{1/2} |(\underline{3}, \underline{3}^*)_{L_z = 1}\rangle \right\}, \quad (1)$$

whereas the  $J_z = +\frac{1}{2}$  component of  $N^*(1238)$  is purely in the  $\{(\underline{6}, \underline{3})_{L_z = 0}\}$  multiplet. The magnetic-transition operator transforms under the algebra like  $\{(8, \underline{1}) + (\underline{1}, 8); L_z = \pm 1\}$  and its matrix elements between two nucleons are given by two independent transition strengths:  $\{(\underline{6}, \underline{3})_{L_z = 0}\} \leftrightarrow \{(\underline{3}^*, \underline{3})_{L_z = -1}\}$  and  $\{(\underline{3}^*, \underline{3})_{L_z = 0}\} \leftrightarrow \{(\underline{3}^*, \underline{3})_{L_z = -1}\}$ . Since the  $(\underline{3}^*, \underline{3}) \leftrightarrow (\underline{3}^*, \underline{3})$  transition is a pure  $F$  transition, it does not contribute to the neutron moment. The magnetic  $N \rightarrow N^*$  transition is therefore simply related to  $\mu_A(n)$  and we find

$$\mu^*/\mu_A(n) = -\sqrt{2}/\cos\theta, \quad (2)$$

where  $\mu^*$  is the matrix element of the  $z$  component of the transition moment between the  $J_z = \frac{1}{2}$  states of  $N$  and  $N^*$ . In a recent analysis of photoproduction data in the neighborhood of the 3-3 resonance, Dalitz and Sutherland<sup>12</sup> have obtained (in nucleon magnetons)

$$\mu^* = (1.28 \pm 0.02) \times \frac{2}{3} \sqrt{2} \mu_{\text{tot}}(p) = 3.36 \pm 0.05.$$

By inserting the  $\cos\theta$  value obtained in I from the axial-vector transitions, we obtain from Eq. (2)

$$\mu^* - \mu_A(n) \times \sqrt{2}/\cos 37^\circ = 3.40.$$

The agreement is remarkably good. In fact it is better than what one should expect, in view of the approximations introduced in the model.

The ratio between the anomalous moments of the proton and neutron cannot be uniquely expressed in terms of the mixing angle  $\theta$  without additional assumptions. It depends on the ratio  $k$  between the reduced matrix elements of the magnetic transitions  $(\underline{3}^*, \underline{3}) \rightarrow (\underline{3}^*, \underline{3})$  and  $(\underline{6}, \underline{3}) \rightarrow (\underline{3}^*, \underline{3})$ . This ratio is a free adjustable parameter of the theory and it can always be fitted to the  $\mu_A(p)/\mu_A(n)$  ratio. We obtain

$$\mu_A(p)/\mu_A(n) = -1 + k \tan\theta, \quad (3)$$

and for  $k=0$ ,

$$\mu_A(p) = -\mu_A(n). \quad (4)$$

We find that the  $(\underline{3}^*, \underline{3}) \rightarrow (\underline{3}^*, \underline{3})$  transition should be smaller than the  $(\underline{6}, \underline{3}) \rightarrow (\underline{3}^*, \underline{3})$  transition, at least by an order of magnitude. In order to check this we can roughly estimate the photoproduction amplitude of the second nucleon resonance  $N^{**}(1512, J^P = \frac{3}{2}^-)$  assuming that its  $J_z = +\frac{1}{2}$  states are mostly in the  $(\underline{3}^*, \underline{3})$  and  $(\underline{3}, \underline{3}^*)$ . We find (for  $k=0$ )

$$\mu^{**} \approx \frac{1}{2} \mu^* \cot\theta \approx 2. \quad (5)$$

In the absence of a reliable detailed analysis of the photoproduction amplitudes in the  $N^{**}(1512)$  region, and in view of the difficulties in separating the background and the  $E1$ ,  $L1$ , and  $M2$  contributions, it is hard to compare this with the data, but the order of magnitude of the result is reasonable in the sense that it is smaller than  $\mu^*$  but not negligible. We cannot expect to do any better with our crude assumptions about the  $N^*$  and  $N^{**}$  classification.

A by-product of the present model of representation mixing is the predicted zero anoma-

lous moment for the decuplet resonances (including  $\Omega^-$ ). This is hard to test, of course, but we might add that SU(3) predicts that the total magnetic moment of any state in the decuplet is proportional to its charge, and that is, of course, consistent with our present result.<sup>13</sup>

We can now summarize the situation as follows: By assuming a simple model of  $U(3) \otimes U(3)$  representation mixing for the nucleon and using only one free mixing angle, we are able to calculate four transition-matrix elements which can be directly or indirectly [via partial conservation of axial-vector current (PCAC)] compared with experiment. The agreement, presented in Table I, is excellent. A fifth quantity,  $\mu_A(p)/\mu_A(n)$ , remains undetermined and is expressed in terms of an additional adjustable parameter. Our simple mixing scheme cannot be incorporated in any simple way into the larger  $U(6)_W$  current algebra, which includes, in addition to the usual vector and axial-vector currents, some components of tensor currents. The suggestion of Gatto, Maiani, and Preparata<sup>3</sup> that the nucleon has components in the  $\underline{56}$ ,  $L=0$  and  $\underline{20}$ ,  $L=1$  representations of SU(6) leads to  $\mu_A(p) = 0$ ,  $\mu^* = 0$ , and is clearly in contradiction with experiment. If we insist on having some SU(6)<sub>W</sub> interpretation, we should probably assign our  $(\underline{3}^*, \underline{3})$  and  $(\underline{3}, \underline{3}^*)$  to the  $\underline{70}$  and find ourselves with a nucleon having both  $W = \frac{1}{2}$  and  $W = \frac{3}{2}$  components. Another amusing possibility may be to assign the  $(\underline{3}^*, \underline{3})$  and  $(\underline{3}, \underline{3}^*)$  components of the  $J_z = \frac{1}{2}$  nucleon to different SU(6)<sub>W</sub> representations which cannot be connected by a  $\underline{35}$ , such as  $\underline{20}$  and  $\underline{700}$ . In this case we automatically get  $\mu(p) = -\mu(n)$  without

Table I. Comparison with experiment.

	Theoretical expression	Theoretical value for $\theta = 37^\circ$	Experimental value
$G_A$	$\frac{1}{3}(4 \cos^2\theta + 1)$	1.18	$1.18 \pm 0.02^a$
$G^*$	$\frac{4}{3} \cos\theta$	1.06	$1 \pm 0.2^b$
$\alpha = \frac{d}{d+f}$	$\frac{2 \cos^2\theta + 1}{4 \cos^2\theta + 1}$	0.64	$0.665 \pm 0.018^c$
$\mu^*/\mu_A(n)$	$-\sqrt{2}/\cos\theta$	1.78	$1.76 \pm 0.03^d$

<sup>a</sup>C. S. Wu, unpublished.

<sup>b</sup>The estimate  $G^* \approx 1$  is obtained by using PCAC and the contribution of the  $N^*$  region to the integral over  $\pi p$  cross sections which appears in the Adler-Weisberger formula. The error of 20% includes the 10-15% expected discrepancy for PCAC and the ambiguities of subtracting the resonance background.

<sup>c</sup>N. Brene, M. Roos, et al., private communication from M. Roos.

<sup>d</sup>R. H. Dalitz and D. G. Sutherland, to be published.

any additional assumptions. We prefer, however, both the simplicity and our better physical understanding of the  $U(3) \otimes U(3)$  current algebra, and we suggest that this is the appropriate framework for studying the problem.

All the results of this paper as well as of I, except the  $d/f$  ratio for the axial-vector transitions, can be derived by using the (chiral or collinear)  $U(2) \otimes U(2)$ . The  $J_z = \frac{1}{2}$  nucleon will then be in a combination of  $(1, \frac{1}{2})_{L_z=0}$ ,  $(0, \frac{1}{2})_{L_z=0}$ , and  $(\frac{1}{2}, 0)_{L_z=1}$ , and in order to obtain  $\mu_A(P) = -\mu_A(n)$  we have to assume that the transition  $\{(\frac{1}{2}, 0)_{L_z=0}\} \leftrightarrow \{(\frac{1}{2}, 0)_{L_z=1}\}$  can be neglected either because of a  $U(3) \otimes U(3)$  selection rule or because the reduced matrix element is negligible. We also observe that our specific mixture of  $(1, \frac{1}{2})$ ,  $(0, \frac{1}{2})$ , and  $(\frac{1}{2}, 0)$  is the only combination which fits the data and does not include  $|L_z| > 1$  or  $I > \frac{5}{2}$  states.

The author acknowledges helpful discussions with S. M. Berman, S. Fubini, M. Gell-Mann, H. J. Lipkin, and S. Meshkov. After completing this work the author was informed by B. W. Lee that he and I. S. Gerstein have obtained similar results using a slightly different mixing model.

---

\*Work supported by the U. S. Atomic Energy Commission.

†On leave of absence from the Weizmann Institute, Rehovoth, Israel.

<sup>1</sup>M. Gell-Mann, Phys. Rev. 125, 1067 (1962); Phys. Rev. 1, 63 (1964).

<sup>2</sup>H. Harari, Phys. Rev. Letters 16, 964 (1966).

<sup>3</sup>R. Gatto, L. Maiani, and G. Preparata, Phys. Rev. Letters 16, 377 (1966), have independently obtained results similar to those of Ref. 2. They used, however, the 20,  $L=1$  representation which leads to the disastrous predictions for the magnetic transitions discussed at the end of the present paper. Some other  $U(3) \otimes U(3)$  and  $SU(6)$  aspects of the problem were stud-

ied by H. J. Lipkin, H. Rubinstein, and S. Meshkov, to be published, and by N. Cabibbo and H. Ruegg, to be published.

<sup>4</sup>R. F. Dashen and M. Gell-Mann, in Proceedings of the Third Coral Gables Conference on Symmetry Principles at High Energies, University of Miami, 1966 (W. H. Freeman & Company, San Francisco, California, 1966).

<sup>5</sup>S. L. Adler, Phys. Rev. Letters 14, 1051 (1965); W. I. Weisberger, Phys. Rev. Letters 14, 1051 (1965).

<sup>6</sup>C. A. Levinson and I. J. Muzinich, Phys. Rev. Letters 15, 715 (1965); D. Amati, C. Bouchiat, and J. Nuyts, Phys. Letters 19, 59 (1965); L. Pandit and J. Schechter, Phys. Letters 19, 56 (1965); W. I. Weisberger, Phys. Rev. 143, 1302 (1966).

<sup>7</sup>In the magnetic-moment sum rule which was derived by S. L. Adler, Phys. Rev. 143, 1144 (1966); J. D. Bjorken, unpublished; and N. Cabibbo and L. A. Radicati, Phys. Letters 19, 697 (1966), the decuplet-dominance assumption leads to a wrong sign for the integral over intermediate states. The amount of success of the same assumption in the case of the photoproduction sum rule of S. Fubini, G. Furlan, and C. Rossetti, to be published, is still unsettled (S. L. Adler, private communication).

<sup>8</sup>Adler, Bjorken, and Cabibbo and Radicati, Ref. 7; Dashen and Gell-Mann, Ref. 4.

<sup>9</sup>Fubini, Furlan, and Rossetti, Ref. 7.

<sup>10</sup>Since the sum rules derived in Refs. 8 and 9 involve current densities and not only integrated currents, one cannot rule out the possibility of nontrivial divergence terms in the commutator.

<sup>11</sup>This can be proved by evaluating the sum rule between  $\Lambda$  states and obtaining from decuplet dominance  $g_{\Lambda\Lambda\eta}\mu(\Lambda)=0$ . Since the  $d/f$  ratio for the  $BBM$  coupling (or the axial-vector current) does not vanish,  $g_{\Lambda\Lambda\eta} \neq 0$  and  $\mu(\Lambda)=0$ . This implies  $\mu(n)=0$ . Decuplet dominance also leads to  $\mu_A^S(N)=0$  and, combined with  $\mu(n)=0$ , to  $\mu(p)=0$ . We also obtain  $\mu^*=0$ , for all octet-decuplet transitions.

<sup>12</sup>R. H. Dalitz and D. G. Sutherland, to be published.

<sup>13</sup>As long as there is no experimental determination of the decuplet moments, the theoretical ambiguities in defining an "anomalous moment" for a spin- $\frac{3}{2}$  object are not very important for our purposes.