

¹⁰T. L. Trueman, Nucl. Phys. 26, 57 (1961).

¹¹This result follows from the fact that the scattering lengths can be expressed directly in terms of T_0 (which is determined) and the effective-range relation from Ref. 10.

¹²T. B. Day, Nuovo Cimento 18, 381 (1960). The $2P$ - $1S$ capture rate is $0.8 \times 10^{13} \text{ Sec}^{-1}$.

¹³G. R. Burleson, D. Cohen, R. C. Lamb, D. N. Mi-

chael, R. A. Schluter, and T. O. White, Jr., Phys. Rev. Letters 15, 70 (1965).

¹⁴D. N. Michael, Phys. Rev. 158, 1343 (1967). We use the experimental results from this latter report since it is subsequent to Ref. 13.

¹⁵F. von Hippel and J. H. Douglas, Phys. Rev. 146, 1042 (1966).

¹⁶J. L. Uretsky, Phys. Rev. 147, 906 (1966).

ERRATA

SPIN POLARIZATION NEAR LOCALIZED MOMENTS IN METALS. C. P. Flynn, G. W. Stupian and D. Lazarus [Phys. Rev. Letters 19, 572 (1967)].

In the table caption, "1000°C" should read "1000°K". The value of Γ_{ex} calculated from the Knight-shift change caused by Gd impurities should read -0.8 eV , or $\Gamma_{\text{ex}}V \approx -25 \text{ eV } \text{\AA}^3$, in place of the values -1.0 eV or $-30 \text{ eV } \text{\AA}^3$ quoted. These corrections do not modify the conclusions drawn in the Letter.

EVIDENCE FOR THE EXISTENCE OF THE HYPERNUCLEUS ΛLi^6 . Dirk-Michael Harmsen [Phys. Rev. Letters 19, 1186 (1967)].

Page 1186, column 2, line 17 should read, "... free-particle system $\Lambda + p + \text{He}^4 \dots$."

Page 1187, column 2, line 37 should read, "... the odds against the ΛBe^7 interpretation for the present event are at least 10:1. This estimate takes into account that only about 35 % of all ΛLi have a range longer than $25 \mu\text{m}$ (see Ref. 4)." I am indebted to Dr. D. H. Davis and Dr. J. Sacton for pointing this out.