

PHYSICAL REVIEW LETTERS

VOLUME 19

31 JULY 1967

NUMBER 5

NUCLEAR ALIGNMENT OF THE 1S_0 GROUND STATE OF ^{131}Xe BY ELECTRON PUMPING AND METASTABILITY-EXCHANGE COLLISIONS

Tetsuo Hadeishi and Chung-Heng Liu

Lawrence Radiation Laboratory, University of California, Berkeley, California

(Received 5 June 1967)

An alignment of a nucleus in the 1S_0 ground state of ^{131}Xe was achieved through exchange collisions with atoms in the metastable state formed and aligned by electron impact. It was observed by monitoring the change of linearly polarized resonance radiation by the $\text{Xe}(^3P_2)$ metastable state as we caused a nuclear magnetic resonance.

In this Letter, we report on a new method of nuclear alignment in the 1S_0 ground state of a noble-gas atom. The nucleus was aligned by means of a transfer of alignment due to "strong" collisions between (a) the initially randomly oriented nucleus of ^{131}Xe with nuclear spin $I = \frac{3}{2}$ in the 1S_0 ground state and (b) 3P_2 metastable-state atoms aligned by electron pumping by electron impact; the collisions result in nuclear alignment of ^{131}Xe in the ground state. There are two major processes of metastable-state alignment by electron impact: (a) by direct electron-impact excitation from the ground state to the metastable state, resulting in the alignment, and (b) electron-impact excitation and alignment of those energy levels lying higher than the metastable state, which then cascade to the metastable state, resulting in transfer of alignments to the metastable state. Although the second process (cascading) corresponds to electron pumping, similar to optical pumping, we shall for simplicity refer to processes (a) and (b) together as electron pumping.

A method of alignment of the metastable-state atoms of Xe was described in our previous Letter,¹ dealing with exchange collisions between

the $\text{Xe}^+(^2P_{3/2})$ ionic ground state and the $\text{Xe}(^3P_2)$ metastable state. We used a mechanism of nuclear alignment very similar to the exchange collisions between the ionic ground state and metastable state of Xe.¹

The main purposes of this experiment were (a) to give additional experimental evidence to support the processes involved in our observation of exchange collisions between the ionic ground state $\text{Xe}^+(^2P_{3/2})$ and the metastable state $\text{Xe}(^3P_2)$, both aligned by electron impact, since considerable controversy has arisen in regard to the processes involved; (b) to offer a new method that may work with short-lived radioactive gases that it is not possible to observe by conventional nmr techniques; and (c) to investigate an electron-pumping cycle.

Definite advantages of our method over the conventional nmr and optical-pumping method are (a) a much smaller sample required for this experiment than for the conventional nmr method and (b) faster alignment of the metastable state by electron pumping than by optical pumping, thus extending the nmr investigation to a great variety of atoms.

The method of detection is based on the spin-exchange experiment first performed in the

brilliant work by Dehmelt,² using an optical-pumping method. In the Dehmelt experiment, a mixture of vapor is illuminated by circularly polarized resonance radiation of one of two species of atoms, thus orienting the system of atoms in the ground state. The orientation so generated in the first system is communicated to the second system by collisions. The magnetic resonance of the second system is monitored by change in the transparency of the first system to resonance radiation.

In our experiment, instead of one of the systems being oriented by resonance radiation, the system is aligned by electron pumping from the ground state to the metastable state (first system). The alignment of the metastable state is communicated to the ground state of ^{131}Xe (second system) through metastability exchange collisions.³

A "strong" collision, in which one atom is in the ground state and the other in the metastable state, results in an exchange of electron angular momentum and excitation energy, so that a nucleus which enters the collision in a ground state may leave it in a metastable state.^{3,4} It is assumed that angular momentum is conserved in the collisions.

Our paramagnetic resonance experiment on Xe metastable-state atoms $\text{Xe}(^3P_2)$, formed and aligned by electron impact,¹ showed that the magnetic sublevels with $M_J = 0$ and ± 1 are more selectively excited than those states with $M_J = \pm 2$, as one would expect from the theory of inelastic collisions for low-energy electronic impact.

Since the nuclear spin of ^{131}Xe is $\frac{3}{2}$, it is evident that because of conservation of angular momentum in collisions, the $M_I = \frac{1}{2}$ and $-\frac{1}{2}$ in the 1S_0 ground state are more populated in equilibrium,⁵ considering the hyperfine structure of ^{131}Xe . The magnetic resonance of the ground state can be monitored by the change in the transparency of the system of metastable-state atoms to resonance radiation.⁵

Figure 1 shows the experimental results. The magnetic field was calibrated by noting the paramagnetic resonance of even isotopes of the metastable state of $\text{Xe}(^3P_2)$.¹ Full magnetic-resonance bandwidth was 15 Hz. We believe that this surprisingly wide bandwidth is due to the inhomogeneous magnetic field due to the indirectly heated cathode heater wire. The observed nuclear magnetic moment is $\mu = 0.6874 \pm 0.0023 \mu_B$. It is in agreement with




FIG. 1. The first trace (a) shows the paramagnetic resonance of $\text{Xe}(^3P_2)$ even isotopes. The second trace (b) shows the nuclear magnetic resonance of ^{131}Xe in the 1S_0 ground state with a linearly oscillating magnetic field at $\nu = 2.5680$ Hz. The third trace (c) shows a typical nmr signal by a digitalized signal-averaging system for four sweeps over the resonance at different H_0 and ν values from those of trace (a).

published values found by means of conventional nmr methods.⁶ Since the purpose of this experiment was not to measure the magnetic moment but instead to study the phenomena, no attempts were made to improve the accuracy. However, it is evident, even with $\Delta\nu = 15$ Hz, that at higher value of magnetic field, extremely precise nuclear-magnetic-moment determination is possible.

Figure 2 shows the experimental arrangement. It is identical to that used in our previous experiment,¹ except that it has no resonance lamp for observing the resonance absorption. Although it may seem surprising, we found that it is possible to use the resonance radiation emitted from the plasma as a light source to monitor the change in the resonance absorption by




FIG. 2. Block diagram of experimental setup. Electronic equipment used throughout the experiment was all commercially available except for a home-made electron tube.

the metastable state. The electron tube was operated under the space-charge neutralization condition, with a gas pressure of about 0.55×10^{-3} Torr and an electron current density of 70 mA/cm^2 , at slightly above the ionization potential energy.

We believe that our observation of the nmr signal of the 1S_0 ground state with $I = \frac{3}{2}$ demonstrated the transfer of alignment of the $\text{Xe}(^3P_2)$ metastable state to the nucleus in the 1S_0 ground state through collisions. We believe that our nmr observation of the nucleus gives a strong support for our observation of exchange collisions between the $\text{Xe}+(^2P_{3/2})$ and $\text{Xe}(^3P_2)$ states in the case of alignment (not orientation), since the only difference in this experiment is that the angular momentum involved in the 1S_0 ground state is the nuclear angular momentum with $I = \frac{3}{2}$, while in our previous experiment the angular momentum involved is the electronic angular momentum with $J = \frac{3}{2}$. The theory describing how the alignment of the metastable state is expected to be changed as one resonates the aligned ground state is to be published elsewhere. This experiment may perhaps be thought of as an "electron-pumping" experiment, similar to an optical pumping of ground-state atoms

first demonstrated by Kastler,⁷ although the processes involved in this experiment are much more complex. Also, electron-pumping phenomena of the $\text{Xe}(^3P_2)$ metastable state due to cascading transitions from the higher states, resulting in the alignment of the metastable state, in addition to the alignment of the metastable state by direct electron impact, was experimentally observed. This phenomenon will be reported elsewhere.

Another interesting feature of this experiment may perhaps be that, by means of an extremely simple device (just a simple diode!), it was possible to observe nmr of ^{131}Xe quite readily. It is somewhat obvious that by use of this technique, radioactive samples such as ^{133}Xe , ^{135}Xe , and many other noble gas atoms having $I > \frac{1}{2}$ should readily be observed.

It is our great pleasure to express our appreciation to Dr. G. W. Series of The Clarendon Laboratory for stimulating conversation and private communication. The first author also appreciates the interesting and stimulating comments of Professor H. G. Dehmelt of the University of Washington during his visit to our Laboratory.

*Work supported by the U. S. Atomic Energy Commission.

¹T. Hadeishi and C.-H. Liu, Phys. Rev. Letters **17**, 513 (1966). A more satisfactory theory of this experiment is to be published.

²H. G. Dehmelt, Phys. Rev. **105**, 1924 (1957).

³F. D. Colegrove, L. D. Scheerer, and G. K. Walters, Phys. Rev. **132**, 2561 (1963).

⁴R. B. Partridge and G. W. Series, Proc. Phys. Soc. (London) **88**, 983 (1966).

⁵A theory describing this phenomenon is to appear in a forthcoming paper.

⁶D. Brinkmann, Helv. Phys. Acta **36**, 413 (1963).

⁷P. A. Kastler, J. Phys. Rad. **11**, 255 (1950).




FIG. 1. The first trace (a) shows the paramagnetic resonance of $\text{Xe}(^3P_2)$ even isotopes. The second trace (b) shows the nuclear magnetic resonance of ^{131}Xe in the 1S_0 ground state with a linearly oscillating magnetic field at $\nu = 2.5680 \text{ Hz}$. The third trace (c) shows a typical nmr signal by a digitalized signal-averaging system for four sweeps over the resonance at different H_0 and ν values from those of trace (a).