

ABSENCE OF MOTT TRANSITION IN AN EXACT SOLUTION
OF THE SHORT-RANGE, ONE-BAND MODEL IN ONE DIMENSION

Elliott H. Lieb* and F. Y. Wu

Department of Physics, Northeastern University, Boston, Massachusetts

(Received 22 April 1968)

The short-range, one-band model for electron correlations in a narrow energy band is solved exactly in the one-dimensional case. The ground-state energy, wave function, and the chemical potentials are obtained, and it is found that the ground state exhibits no conductor-insulator transition as the correlation strength is increased.

The correlation effect of electrons in a partially filled energy band has been a subject of interest for many years.¹⁻⁴ A realistic model which takes this correlation into consideration, and which is hopefully amenable to mathematical treatment, is the short-range, one-band model considered by a number of authors.²⁻⁵ In this model, one pictures the electrons in a narrow energy band hopping between the Wannier states of neighboring lattice sites, with a repulsive interaction energy between two electrons of opposite spins occupying the same lattice site. The central problems of interest have been (a) the possible existence of a "Mott transition" between conducting and insulating states as the strength of the interaction is increased, and (b) the magnetic nature (ferromagnetic or antiferromagnetic) of the ground state. While previous treatments of this model have always been approximate, we have succeeded in solving the model exactly in the one-dimensional case. Our exact result shows that the Mott transition does occur in the ground state of the one-dimensional model. Furthermore, a general theorem of Lieb and Mattis⁶ on one-dimensional systems tells us that the ground state is necessarily antiferromagnetic.

It may be argued that the absence of a Mott transition in one dimension is irrelevant for the study of real three-dimensional systems because of the folkloristic dictum that there are never any phase transitions in one dimension with short-range interactions. In actual fact, the dictum is only true for nonzero temperature; the ground state is another matter. Generally speaking, when a Hamiltonian is considered to be a function of some parameter, U (which in our case is the electron-electron repulsion), singularities with respect to U usually do appear in the ground-state wave function, energy, polarizability, etc., even in one dimension. A good example of this is the one-dimensional Heisenberg chain (to which the present model is very close) which, when considered as a function of the anisotropy parameter, does have two singularities

in the ground state and, presumably, no singularities for nonzero temperatures.^{7,8}

Consider a crystal (one-, two-, or three-dimensional) of N_a lattice sites with a total of $N \leq 2N_a$ electrons. We suppose that the electrons can hop between the Wannier states of neighboring lattice sites, and that each site is capable of accommodating two electrons of opposite spins, with an interaction energy $U > 0$. The Hamiltonian to consider is then²⁻⁵

$$H = T \sum_{\langle ij \rangle} \sum_{\sigma} c_{i\sigma}^{\dagger} c_{j\sigma} + U \sum_i c_{i\uparrow}^{\dagger} c_{i\uparrow} c_{i\downarrow}^{\dagger} c_{i\downarrow}, \quad (1)$$

where $c_{i\sigma}^{\dagger}, c_{i\sigma}$ are, respectively, the creation and annihilation operators for an electron of spin σ in the Wannier state at the i th lattice site, and the sum

$$\sum_{\langle ij \rangle}$$

is restricted to nearest-neighbor sites.

First of all, it can be shown that the energy spectrum of H is invariant under the replacement of T by $-T$.⁹ Therefore, for simplicity we shall take, in appropriate units, $T = -1$. Since the numbers M of down-spin electrons and M' of up-spin electrons are good quantum numbers ($M + M' = N$), we may designate the ground-state energy of H by $E(M, M'; U)$. It is then easy to derive the following relations [by considering holes instead of particles in (1)]:

$$\begin{aligned} E(M, M'; U) &= -(N_a - M - M')U \\ &\quad + E(N_a - M, N_a - M'; U) \\ &= MU + E(M, N_a - M'; -U) \\ &= M'U + E(N_a - M, M'; -U). \end{aligned} \quad (2)$$

Without loss of generality, therefore, we may take

$$S_z \equiv \frac{1}{2}(N - 2M) \geq 0 \text{ and } N \leq N_a$$

(less than half-filled band).

It can similarly be shown that the maximum energy $G(M, M'; U)$ is related to the ground-state energy by

$$G(M, M'; U) = M'U - E(N_a - M, M'; U).$$

Therefore, a knowledge of the ground-state energies also tells us about the maximum energies.

For a one-dimensional system, the lattice sites can be numbered consecutively from 1 to N_a . Let $f(x_1, x_2, \dots, x_M, x_{M+1}, \dots, x_N)$ represent the amplitude in ψ for which the down spins are at the sites x_1, \dots, x_M , and the up spins at x_{M+1}, \dots, x_N . Then the eigenvalue equation $H\psi = E\psi$ leads to

$$\begin{aligned} & - \sum_{i=1}^N \sum_{s=\pm 1} f(x_1, \dots, x_i + s, \dots, x_N) \\ & + U \sum_{i < j} \delta(x_i - x_j) f(x_1 \dots x_N) \\ & = E f(x_1 \dots x_N), \end{aligned} \tag{3}$$

where it is understood that we require a solution of the form

$$f(x_1, x_2, \dots, x_M | x_{M+1}, x_{M+2}, \dots, x_N) \tag{4}$$

which is antisymmetric in the first M and the last $N-M$ variables.

In each region defined by $1 \leq x_{Q1} \leq x_{Q2} \leq \dots \leq x_{QN} \leq N$, we make the following Ansatz for f :

$$\begin{aligned} & f(x_1, \dots, x_M | x_{M+1}, \dots, x_N) \\ & = \sum_P [Q, P] \exp(i \sum_{j=1}^N k_{P_j} x_{Q_j}), \end{aligned} \tag{5}$$

where $P = (P_1, P_2, \dots, P_N)$ and $Q = (Q_1, Q_2, \dots, Q_N)$ are two permutations of the numbers $(1, 2, \dots, N)$, $\{k_1, k_2, \dots, k_N\}$ is a set of N unequal real numbers, and $[Q, P]$ is a set of $N! \times N!$ coefficients to be determined.

The coefficients $[Q, P]$ are not all independent. The condition of single valuedness (or continuity) of f and the requirement that (5) be a solution of (3) lead to the following:

$$E = -2 \sum_{j=1}^N \cos k_j \tag{6}$$

and, for all Q and P , the coefficients $[Q, P]$ must

be chosen to satisfy the relations

$$[Q, P] = Y_{nm}^{ab} [Q, P']. \tag{7}$$

In (7), Y_{nm}^{ab} is an operator defined by

$$\begin{aligned} Y_{nm}^{ab} &= \frac{-\frac{1}{2}iU}{\sin k_n - \sin k_m + \frac{1}{2}iU} \\ &+ \frac{\sin k_n - \sin k_m}{\sin k_n - \sin k_m + \frac{1}{2}iU} P^{ab}, \end{aligned} \tag{8}$$

where, for $j = i + 1$,

$$Q_i = a = Q'_j, \quad Q_j = b = Q'_i,$$

$$Q_k = Q'_k \text{ for all } k \neq i, j;$$

$$P_i = m = P'_j, \quad P_j = n = P'_i,$$

$$P_k = P'_k \text{ for all } k \neq i, j;$$

and P^{ab} is an operator which exchanges $Q_i = a$ and $Q_j = b$.

It is fortunate that the Ansatz (5) and the algebraic consistency conditions (7) and (8) have, in essence, appeared before in the study of the one-dimensional delta-function gas for particles in a continuum. The first solution of that problem was for bosons (symmetric f) by Lieb and Luttinger¹⁰ but this case is not relevant here, besides which the consistency conditions there are trivial to solve. The two-component fermion case was solved by McGuire¹¹ for $M = 1$, but again (7) is trivial because of translational invariance. The next development was the solution of the case $M = 2$ by Flicker and Lieb¹² by an inelegant algebraic method which could not be easily generalized. However, the case $M = 2$ is the simplest one which displays the full difficulty of the problem. Shortly thereafter, Gaudin¹³ published the solution of the general- M problem. The method of his brilliant solution did not appear for some time and is now available as his thesis.¹⁴ In the meantime, Yang¹⁵ also discovered the method of solution (essentially the same as Gaudin's) and published it with considerable detail. Here, we have followed Yang's notation with slight modification.

The important point is that our Eqs. (7) and (8) are the same as for the continuum gas except for the replacement of k by $\sin k$ in the latter. This has no effect on the beautiful algebraic analysis which finally leads to the following condi-

tions which determine the set $\{k_1, 2, \dots, k_N\}$:

$$N_a k_j = 2\pi I_j + \sum_{\beta=1}^M \theta(2 \sin k_j - 2\Lambda_\beta), \quad j=1, 2, \dots, N, \tag{9}$$

where the Λ 's are a set of real numbers related to the k 's through

$$-\sum_{j=1}^N \theta(2\Lambda_\alpha - 2 \sin k_j) = 2\pi J_\alpha - \sum_{\beta=1}^M \theta(\Lambda_\alpha - \Lambda_\beta), \quad \alpha=1, 2, \dots, M, \tag{10}$$

$$\theta(p) \equiv -2 \tan^{-1}(2p/U), \quad -\pi \leq \theta < \pi, \tag{11}$$

and I_j = integers (or half-odd integers) for M = even (or odd), J_α = integers (or half-odd integers) for M' = odd (or even). An immediate consequence is

$$\sum_{j=1}^N k_j = \frac{1}{N_a} (\sum_j I_j + \sum_\alpha J_\alpha). \tag{12}$$

For the ground state, J_α and I_j are consecutive integers (or half-odd integers) centered around the origin and satisfying $\sum_j k_j = 0$.

In the limit of $N \rightarrow \infty$, $N_a \rightarrow \infty$, $M \rightarrow \infty$ with the ratios N/N_a , M/N_a kept finite, the real numbers k and Λ are distributed continuously between $-Q$ and $Q \leq \pi$ and $-B$ and $B \leq \infty$, with density functions $\rho(k)$ and $\sigma(\Lambda)$, respectively. Equations (9) and (10) then lead to the coupled integral equations for the distribution function $\rho(k)$ and $\sigma(\Lambda)$:

$$2\pi\rho(k) = 1 + \cos k \int_{-B}^B \frac{8U\sigma(\Lambda)d\Lambda}{U^2 + 16(\sin k - \Lambda)^2}, \tag{13}$$

$$\int_{-Q}^Q \frac{8U\rho(k)dk}{U^2 + 16(\Lambda - \sin k)^2} = 2\pi\sigma(\Lambda) + \int_{-B}^B \frac{4U\sigma(\Lambda')d\Lambda'}{U^2 + 4(\Lambda - \Lambda')^2}, \tag{14}$$

where Q and B are determined by the conditions

$$\int_{-Q}^Q \rho(k)dk = N/N_a, \tag{15}$$

$$\int_{-B}^B \sigma(\Lambda)d\Lambda = M/N_a. \tag{16}$$

The ground-state energy (6) now becomes

$$E = -2N_a \int_{-Q}^Q \rho(k) \cos k dk. \tag{17}$$

We have established the following:

(a) Equations (13)-(16) have a unique solution which is positive for all allowed B and Q .

(b) M/N is a monotonically increasing function of B reaching a maximum of $\frac{1}{2}$ at $B = \infty$. This is the antiferromagnetic case, $S_z = 0$, and corresponds to the absolute ground state.

(c) N/N_a is a monotonically increasing function of Q , reaching a maximum of 1 (half-filled band) at $Q = \pi$.

For $B = \infty$ and $Q = \pi$, (13)-(16) can be solved in closed form by Fourier transforms with the re-

sult

$$\sigma(\Lambda) = (2\pi)^{-1} \int_0^\infty \text{sech}(\frac{1}{4}\omega U) \times \cos(\omega\Lambda) J_0(\omega) d\omega, \tag{18}$$

$$\rho(k) = (2\pi)^{-1} + \pi^{-1} \cos k$$

$$\times \int_0^\infty \frac{\cos(\omega \sin k) J_0(\omega) d\omega}{1 + \exp(\frac{1}{2}\omega U)}, \tag{19}$$

$$E \equiv E(\frac{1}{2}N_a, \frac{1}{2}N_a; U)$$

$$= -4N_a \int_0^\infty \frac{J_0(\omega) J_1(\omega) d\omega}{\omega [1 + \exp(\frac{1}{2}\omega U)]}, \tag{20}$$

where J_0 and J_1 are Bessel functions.

To investigate whether the ground state is conducting or insulating, we compute the chemical

potentials μ_+ and μ_- as defined in a forthcoming paper by Mattis¹⁶:

$$\begin{aligned}\mu_+ &\equiv E(M+1, M; U) - E(M, M; U), \\ \mu_- &\equiv E(M, M; U) - E(M-1, M; U).\end{aligned}\quad (21)$$

If μ_+ and μ_- are equal, the system has the property of a conductor. If, on the other hand, we find $\mu_+ > \mu_-$, then the system shares the property of an insulator. We can compute μ_- directly from (9) and (10) by replacing $M \rightarrow M-1$ and $N \rightarrow N-1$, while letting all the k 's, Λ 's, and their distribution functions change slightly. The procedure is quite similar to the calculation of the excitation spectrum of the continuum gas.¹⁰ If $N < \frac{1}{2}N_0$, we can compute μ_+ in the same way and thereby find that $\mu_+ = \mu_-$ for all U . If, however, N is exactly $\frac{1}{2}N_0$, then we must compute μ_+ by using the first line of (2) which tells us that

$$\mu_+ = U - \mu_- \text{ (half-filled band).} \quad (22)$$

The calculation of μ_- can be done in closed form for a half-filled band with the result

$$\begin{aligned}\mu_- - 2 &= -4 \int_0^\infty \frac{J_1(\omega) d\omega}{\omega [1 + \exp(\frac{1}{2}\omega U)]} \\ &= -4 \sum_{n=1}^\infty (-1)^n n \left[\left(1 + \frac{1}{4}n^2 U^2\right)^{\frac{1}{2}} - \frac{1}{2}nU \right].\end{aligned}\quad (23)$$

It can be established from (22) and (23) that, indeed, $\mu_+ > \mu_-$ for $U > 0$, and

$$\lim_{U \rightarrow 0} \mu_\pm = 0.$$

Therefore, we conclude that the ground state for a half-filled band is insulating for any nonzero U , and conducting for $U=0$. That is, there is no Mott transition for nonzero U . This absence of a Mott transition is also reflected by the fact that the ground-state energy and the ground-state wave function are analytic in U on the real axis (except at the origin).

We have also investigated the excitation spectrum $E(p)$ for a given total momentum $\sum_j k_j = p$ and a given value of S_z . Just as in the case of a continuum gas for which the spectrum can be regarded as consisting of several elementary excitations,^{10,15} we find three types of excitations: (I) a "hole" state in the Λ distribution, (II) a "hole" state in the k distribution, and (III) a

"particle" state in the k distribution. While the $S_z = 0$ spin-wave state may have any of these three types of spectra, the $S_z = 1$ spin-wave state is always associated with the type-I spectrum. The type-I excitation has the lowest energy and is characterized by a double periodicity similar to that of an antiferromagnetic chain.⁷ In the limit $U \rightarrow 0$, it goes over to $E(p) = |\sin p|$, while the type-II and -III spectra have the identical limiting form $E(p) = |2 \sin(\frac{1}{2}p)|$. Detailed discussions of these matters will be given elsewhere.

We are grateful to Dr. D. C. Mattis for helpful advice and suggestions and E. L. would like to thank Dr. J. Zittartz for interesting him in the problem.

*Work partially supported by National Science Foundation Grant No. GP-6851.

¹J. H. Van Vleck, Rev. Mod. Phys. 25, 220 (1953).

²M. C. Gutzwiller, Phys. Rev. Letters 10, 159 (1963), and Phys. Rev. 134, A923 (1964), and 137, A1726 (1965).

³J. Hubbard, Proc. Roy. Soc. (London), Ser. A 276, 238 (1963), and 277, 237 (1964).

⁴G. Kemeny, Ann. Phys. (N.Y.) 32, 69, 404 (1965).

⁵See also the review article by C. Herring, in *Magnetism*, edited by G. T. Rado and H. Suhl (Academic Press, Inc., New York, 1966), Vol. IV, Chap. 10.

⁶E. Lieb and D. C. Mattis, Phys. Rev. 125, 164 (1962). The proof of the theorem given in this reference can be adopted to our Hamiltonian (1).

⁷J. des Cloizeaux and M. Gaudin, J. Math. Phys. 7, 1384 (1966).

⁸C. N. Yang and C. P. Yang, Phys. Rev. 150, 321, 327 (1966).

⁹The proof assumes A and B sublattices and uses the unitary transformation

$$\exp[i\pi \sum_{\sigma} \sum_{i \in A} c_{i\sigma}^\dagger c_{i\sigma}]$$

which changes $c_{i\sigma}$ to $-c_{i\sigma}$ and $c_{i\sigma}^\dagger$ to $-c_{i\sigma}^\dagger$ for $i \in A$. This transformation does not change the number operator.

¹⁰E. Lieb and W. Liniger, Phys. Rev. 130, 1605 (1963); E. Lieb, Phys. Rev. 130, 1616 (1963).

¹¹J. B. McGuire, J. Math. Phys. 6, 432 (1965).

¹²M. Flicker, thesis, Yeshiva University, 1966 (unpublished); M. Flicker and E. Lieb, Phys. Rev. 161, 179 (1967).

¹³M. Gaudin, Phys. Letters 24A, 55 (1967).

¹⁴M. Gaudin, thesis, University of Paris, 1967 (unpublished).

¹⁵C. N. Yang, Phys. Rev. Letters 19, 1312 (1967).

¹⁶D. Mattis, to be published.