

two cases: There can be a single Lorentz pole at $s=0$ or an infinite number of poles. Any finite number $m > 1$ leads to nonfactorizable Regge residues β_n for $n > m$.

¹⁰The little group of the Poincare group for unequal-mass scattering at $s=0$ is $E(2)$. Since this is a symmetry group of the scattering amplitude, it might be supposed that the parent-daughter sequence of Regge poles would classify in this case as a single irreducible representation of $E(2)$. This is not so. The rep-

resentation coefficients of $E(2)$ are Bessel functions (cf. J. F. Boyce, R. Delbourgo, A. Salam, and J. Strathdee, to be published). No single irreducible representation can yield the expected t^α behavior of $A(s, t, u)$ for $s=0, t \rightarrow \infty$. The result obtained from a single irreducible representation of $E(2)$ is also singular in the limits $m_1 - m_2 \rightarrow 0, m_3 - m_4 \rightarrow 0$.

¹¹J. B. Bronzan and C. E. Jones, Phys. Rev. Letters 21, 564 (1968).

ERRATA

SUPERCONDUCTIVITY IN MULTIPLE PAIR-BREAKING REGIMES. R. P. Guertin, W. E. Masker, T. W. Mihalisin, R. P. Groff, and R. D. Parks [Phys. Rev. Letters 20, 387 (1968)].

In line 38 of the second column on page 388, the factor 0.006 should be changed to 0.003. The same change should be made in Fig. 1 and the caption for Fig. 1.

TRAJECTORIES IN VENEZIANO'S MODEL. Ralph Roskies [Phys. Rev. Letters 21, 1851 (1968)].

In condition (A) of the theorem, replace $\ln S$ by $\ln s$. The first term of the right side of Eq. (22) should be $2\nu Ks / (\ln s)^{\nu+1}$, and the right-hand side of (25) should read

$$\exp\left[\frac{1}{2}\pi s / (\ln s)\right]^\nu \times \text{lower order terms.}$$

These changes do not affect the argument at all. I wish to thank Professor B. Desai for pointing

out the error in (25).

COULOMB DISTORTION EFFECTS IN LARGE-ANGLE M1 ELECTROEXCITATION. B. T. Chertok and W. T. K. Johnson [Phys. Rev. Letters 22, 67 (1969)].

A paper by Dieter Dreschel [Nucl. Phys. A113, 665 (1968)] was brought to our attention after submission of this Letter for publication. Dreschel has given an excellent quantitative evaluation of the model independency of Coulomb corrections in $M1$ electroexcitation and has demonstrated the inadequacy of the transition radius, R_{tr} , as a unique (experimentally determined) quantity for transverse transitions. Preliminary calculations of the model dependency of the $M1$ radiative widths in our work indicate, e.g., a possible additional uncertainty of 5-7% in Γ_0 for the 11.42-MeV, 1^+ level of ^{28}Si .

The end of the last sentence of paragraph 6 should read, "... versus an experimental value for the ratio of the two shape factors, $A = a^-(^{12}\text{B}) / a^+(^{12}\text{N})$, $A = (1.07 \pm 0.24)\%$ per MeV."