

μb at 150 Mev; this value is consistent with the S -wave total cross section predicted by the value of S determined from the angular distribution. The theory of Drell, Friedman, and Zachariasen,³ taking into account the internal rescattering of a photoproduced π^+ in an S state, predicts a total cross section of $1 \mu\text{b}$ at 150 Mev.

The authors wish to express their appreciation to Gilbert Davidson and Zvi Westreich for their help in taking data.

* This work is supported in part through funds provided by the U. S. Atomic Energy Commission, the Office of Naval Research, and the Air Force Office of Scientific Research.

¹H. L. Anderson and M. Glicksman, Phys. Rev. 100, 268 (1955).

²Chew, Low, Goldberger, and Nambu, Phys. Rev. 106, 1345 (1957).

³Drell, Friedman, and Zachariasen, Phys. Rev. 104, 236 (1956).

NOTES ON ANTIBARYON INTERACTIONS*

A. Pais†

Lawrence Radiation Laboratory, University of California, Berkeley, California
(Received August 3, 1959)

This note presents some simple consequences of (or checks for) invariance considerations as applied to interactions of antibaryons.

A. P conservation in strong interactions.— Consider the reactions $\bar{p} + p \rightarrow n$ particles. If the reaction occurs at rest for $n \geq 4$ or in flight for $n \geq 3$, there are enough independent momentum vectors to form, in any coordinate system, a nonvanishing quantity $\vec{p}_1 \cdot (\vec{p}_2 \times \vec{p}_3)$. The condition of symmetry in the up-down distribution of \vec{p}_1 relative to the (p_2, p_3) plane is then a consequence of P conservation, unless other inherent symmetries require this distribution to be symmetrical anyway. Thus, consider the reaction in flight

$$\bar{p} + p \rightarrow 1 + 2 + R, \quad (1)$$

where 1 and 2 are some specific particles and where the "rest" R may be any assembly of particles. We always work in the (\bar{p}, p) -c.m. system and consider exclusively unpolarized beams and targets. The initial state is in general not an eigenstate of P . But it is an eigenstate of PR , where R is a 180° rotation around any axis perpendicular to (\bar{p}, p) and which we may take perpendicular to $(\bar{p}, 1)$. Let $W(1, E_1, \theta_1; 2, E_2, \theta_2, \phi)$ denote the probability of finding particle 1 (2) with an energy E_1 (E_2) at an angle θ_1 (θ_2) relative to the direction of \bar{p} , where ϕ is the azimuth of 2 relative to the $(\bar{p}, 1)$ plane. Then PR implies

$$\begin{aligned} W(1, E_1, \theta_1; 2, E_2, \theta_2, \phi) \\ = W(1, E_1, \theta_1; 2, E_2, \theta_2, -\phi). \end{aligned} \quad (2)$$

As an example of other symmetries which would imply Eq. (2), we note that if 1 and 2 are both π^+ , Eq. (2) is valid as a consequence of Bose sta-

tistics. If one considers the triple (\bar{p}, π^+, π^-) , however, there seems to be no other known symmetry than PR which leads to Eq. (2).

B. C invariance of strong interactions.— Consider again Reaction (1) and also

$$\bar{p} + p \rightarrow \bar{1} + \bar{2} + \bar{R} \quad (3)$$

[The products in Reaction (3) may or may not be identical with those of Reaction (1).] The initial state ψ_S ($S=0, 1$ is the total spin) is in general not an eigenstate of C . However, we have

$$CR\psi_S = (-1)^{S+S_z}\psi_S, \quad (4)$$

which makes CR useful for an unpolarized beam and target, as here the cross section for any reaction does not involve interference between initial spin states so that CR invariance may be applied to the final states. Let the probabilities referring to Eq. (2) be denoted by \bar{W} . Then CR implies

$$\begin{aligned} W(1, E_1, \theta_1; 2, E_2, \theta_2, \phi) \\ = \bar{W}(\bar{1}, E_1, \pi - \theta_1; \bar{2}, E_2, \pi - \theta_2, \pi - \phi). \end{aligned} \quad (5)$$

In the case of pure pion annihilation the symbols \bar{W} and W refer to the same reaction. In other instances, such as $\bar{p} + p \rightarrow \bar{p} + \Lambda + K^+$, $\bar{p} + p \rightarrow \bar{p} + \bar{\Lambda} + K^-$, they refer to different reactions.

We may also apply CP to the final states, as

$$CP\psi_S = (-1)^{S+1}\psi_S, \quad (6)$$

and obtain

$$\begin{aligned} W(1, E_1, \theta_1; 2, E_2, \theta_2, \phi) \\ = \bar{W}(\bar{1}, E_1, \pi - \theta_1; \bar{2}, E_2, \pi - \theta_2, \pi + \phi). \end{aligned} \quad (7)$$

Equations (2), (5), and (7) are derived channel by

channel. They may therefore also be applied to sums over channels so that there are no complications because of incomplete knowledge of the neutral particles produced.

Whenever P conservation is established, one may check C by using a relation that, in itself, follows either from CR or from CP , such as

$$W(1, E, \theta) = \bar{W}(\bar{1}, E, \pi - \theta). \quad (8)$$

From this relation it follows that the π^0 distribution in any pure pion annihilation channel is symmetric around 90° .

Because antinucleons play a virtual role in low-energy nuclear phenomena, the high degree to which P conservation is known to hold in the latter domain has implications also for these antiparticles. In this sense, relations like Eqs. (2), (5), and (7) may be considered as a useful complement to the low-energy information. Their applicability to very-high-energy phenomena (regardless of the complexity of the events) makes it possible to verify the validity of these conservation laws at frequencies that are perhaps not as sensitively explored in the low-energy nuclear effects.

For two-body reactions like

$$\bar{p} + p \rightarrow \bar{\Lambda} + \Lambda, \quad (9)$$

Eqs. (5) and (7) are trivial. However, here CP also has a useful application. If we denote by $\vec{q}(\Lambda, \theta)$ and $\vec{q}(\bar{\Lambda}, \theta)$, respectively, the polarizations of Λ and $\bar{\Lambda}$, if any, where θ is the angle between the particle in question and the \bar{p} direction, then CP implies

$$\vec{q}(\Lambda, \theta) = \vec{q}(\bar{\Lambda}, \pi - \theta). \quad (10)$$

(Of course P implies more stringently that the \vec{q} vectors are perpendicular to the production plane.) Equation (6) is here also valid in the final state, so that the differential cross section of Eq. (9) is the sum of a triplet-triplet and a singlet-singlet differential cross section. Furthermore, it is readily shown that Eq. (10) also applies generally if $\vec{q}(\Lambda, \theta)$ refers to Reaction (1) and $\vec{q}(\bar{\Lambda}, \theta)$ to (3), again as a consequence of CP . Thus Eq. (10) holds for $(\bar{\Lambda}, \Sigma^0)$ as compared to $(\Lambda, \bar{\Sigma}^0)$ production; the partial cross sections of these reactions should be each other's mirror around 90° .

C. Charge symmetry (CS) and charge independence (CI).—Antihyperon annihilations via strong interactions provide in principle various means to verify CS and CI. In practice, large numbers of antihyperon events are needed for this.¹

For $(\bar{\Lambda}, p)$ annihilation, we have² from CI

$$(\bar{\Lambda}, p \rightarrow K^0, \pi^+) = 2(\bar{\Lambda}, p \rightarrow K^+, \pi^0). \quad (11)$$

This is the only relation that CI imposes.³ A much more interesting situation obtains if the $\bar{\Lambda}$'s are annihilated in deuterium. The reason is that the $(\bar{\Lambda}, d)$ system is self-charge symmetric. [It shares this property with (Λ, d) , but it has considerable phase-space advantage.] Thus let $W(K^+)$ and $W(K^0)$ be the probabilities, respectively, of producing a K^+ (K^0) in $(\bar{\Lambda}, d)$ annihilation. Then CS tells us²

$$W(K^+) = W(K^0), \quad (12)$$

regardless of the complexity of the various annihilation modes possible.⁴ Thus $(\bar{\Lambda}, d)$ interactions may be of particular interest at very high energies. In addition, we have⁵ from CI

$$(\bar{\Lambda}, d \rightarrow n, K^+, \pi^0) = \frac{1}{4}(\bar{\Lambda}, d \rightarrow n, K^0, \pi^+). \quad (13)$$

The annihilation in flight of $\bar{\Sigma}^\pm$ in d also tests CS. Call $W_\pm(K)$ the probability of producing a K in $\bar{\Sigma}^\pm - d$ reactions. Then we have

$$W_\pm(K^+) = W_\mp(K^0). \quad (14)$$

Finally, production reactions in d yield a few CI relations, namely:

$$(\bar{p}, d \rightarrow \bar{\Lambda}, \Lambda, p, \pi^-) = 2(\bar{p}, d \rightarrow \bar{\Lambda}, \Lambda, n, \pi^0), \quad (15)$$

$$(\bar{p}, d \rightarrow \bar{\Sigma}^+, \Lambda, p) = 2(\bar{p}, d \rightarrow \bar{\Sigma}^0, \Lambda, n), \quad (16)$$

$$(\bar{p}, d \rightarrow \bar{\Lambda}, \Sigma^-, p) = 2(\bar{p}, d \rightarrow \bar{\Lambda}, \Sigma^0, n). \quad (17)$$

D. $\bar{\Lambda}$ decay and T invariance.—It has been noted by Okubo⁶ that CPT invariance by itself does not imply the equality of the partial lifetimes of hyperon decay into a given channel and of anti-hyperon decay into the corresponding charge-conjugate channel. In general there are three independent sufficient grounds for two such quantities to be equal, namely (a) absence of final-state interactions, (b) C invariance, and (c) T invariance. In the case of the $\bar{\Lambda}$, there is a fourth independent sufficient ground, namely the $\Delta I = \frac{1}{2}$ rule.⁷ Thus the validity of the latter rule would obviate the possibility of testing T invariance by means of a partial lifetime comparison. This is in principle not the case when one compares the up-down asymmetries α_{ch} and $\bar{\alpha}_{\text{ch}}$ of the decays $\Lambda \rightarrow p + \pi^-$ and $\bar{\Lambda} \rightarrow \bar{p} + \pi^+$. In fact, if one assumes the $\Delta I = \frac{1}{2}$ rule to be valid, one has

$$\rho \equiv \frac{\alpha_{\text{ch}}}{\bar{\alpha}_{\text{ch}}} = \frac{\sin(\delta_{11} - \delta_1 - \Delta)}{\sin(\delta_{11} - \delta_1 + \Delta)}, \quad (18)$$

where δ_{11} and δ_1 are the $P_{1/2}$ and $S_{1/2}$ π -nucleon phase shifts in the $I=\frac{1}{2}$ state. Here Δ has the following properties: If C invariance holds, we have $\Delta=0$, so $\rho=1$; if T invariance holds, we have $\Delta=\pi/2$, so $\rho=-1$. From the magnitude of α_{ch} we know already⁸ that $|\Delta| \gtrsim \pi/4$. We also see that $\rho=-1$ if $\delta_{11}-\delta_1$ is neglected relative to Δ . As this neglect is justified to a good approximation, it follows that a 20% deviation from $\rho=-1$ is the most that can be anticipated.⁹

I am indebted to many physicists at the Lawrence Radiation Laboratory for stimulating discussions.

*This work done under the auspices of the U. S. Atomic Energy Commission.

[†]Permanent address: Institute for Advanced Study,

Princeton, New Jersey.

¹Thus for $\bar{\Lambda}$ production in hydrogen by 1-Bev anti-protons, one would need to have an annihilation cross section as large as ~ 50 mb to have an effect $\sim 1\%$.

²The relations (11) to (17) refer to relative rates and are valid for all energies and all angles.

³In this note, we do not consider inequalities following from CI.

⁴Of course Eq. (12) may also be applied to an individual channel and its charge-symmetric one, such as (n, K^0, π^+) versus (p, K^+, π^-) , etc.

⁵Here the pure $I=1$ state of the K -nucleon system is involved. In $(\bar{\Lambda}, d \rightarrow p, K^0) = (\bar{\Lambda}, d \rightarrow n, K^+)$ we deal with the corresponding pure $I=0$ state.

⁶S. Okubo, Phys. Rev. 109, 984 (1958).

⁷Electromagnetic effects are ignored here.

⁸R. Gatto, Phys. Rev. 108, 1103 (1957); see also S. Weinberg, Phys. Rev. 110, 782 (1958).

⁹Putting $\Delta = \pi/2 - \epsilon$, $\rho \approx -(1 + 0, 2 \tan \epsilon)$.

ERRATA

VALLEY-ORBIT SPLITTING OF ARSENIC DONOR GROUND STATE IN GERMANIUM. G. Weinreich, W. S. Boyle, H. G. White, and K. F. Rodgers [Phys. Rev. Letters 2, 96 (1959)].

In the previous Letter we reported measurements which indicated that in the arsenic donor ground state in germanium the triplet component has a lower energy than the singlet. Since very strong evidence to the contrary^{1,2} has now appeared, we were led to re-examine our experimental procedure in order to attempt to locate the origin of the discrepancy.

We have modified our equipment so as to improve greatly both the homogeneity of the applied strain and the certainty of its orientation. Data taken with the new arrangement indeed reveal a line splitting opposite to the one we previously reported, establishing that the singlet is the lower state in agreement with the other evidence

quoted. Apparently we had been dealing with a rather peculiar pattern of inhomogeneity which, instead of simply broadening the absorption, produced the deceptive line shape shown in our first report.

These new experiments were performed on samples with about ten times less arsenic content than the original ones, since at this concentration the line is much better defined. We do not believe, however, that the different results are connected with the different concentration, since the new samples under the old strain conditions show the same spurious behavior as the more strongly doped ones.

A full report of the experiment will be submitted to the Physical Review.

¹H. Fritzsche, Bull. Am. Phys. Soc. 4, 185 (1959).

²G. Feher, D. K. Wilson, and E. A. Gere, Phys. Rev. Letters 3, 25 (1959).

ELECTROMAGNETIC TRANSITIONS BETWEEN μ MESON AND ELECTRON. S. Weinberg and G. Feinberg [Phys. Rev. Letters 3, 111 (1959)].

In comparing the rate $\omega_{N \rightarrow N'}$ for $\mu^- + N \rightarrow e^- + N'$ via virtual photon absorption with the rate ω_{abs}

for ordinary muon absorption, an important numerical factor was omitted from ω_{abs} . Equation (11) should read¹

$$\omega_{abs} \approx (1/2\pi^2) Z_{eff}^4 \alpha^3 \xi_\beta^2 m_\mu C, \quad (11)'$$