
ERRATA

STRONG POLARIZATION OF THE $1p$ -SHELL CORE IN THE LOWEST 0^+ STATES OF ^{24}Mg AND ^{28}Si . H. P. Morsch, D. Dehnhard, and T. K. Li [Phys. Rev. Lett. 34, 1527 (1975)].

The change made by the editorial office in our original notation for the particle-hole (p-h) components has generated a very misleading and unphysical picture of the process. Our notation, e.g., $(1p_{1/2} - 1d_{5/2})$ was changed to $(1p_{1/2})(1d_{5/2})^{-1}$ which can never couple to $J^\pi = 0^+$. The notation $(1p_{1/2} - 1d_{5/2})$ was meant to include all possible $(1p_{1/2})^{-n}(1d_{5/2})^n$ components. Dominant contributions are from 2p-2h and 4p-4h components. 1p-1h excitations which involve two major shells, e.g. $(1p_{1/2})^{-1}(2p_{1/2})$, were not included.

PREFERRED SITES OF IMPURITIES IMPLANTED IN Be: LATTICE LOCATION AND QUADRUPOLE INTERACTIONS. E. N. Kaufmann, P. Raghavan, R. S. Raghavan, E. J. Ansaldo, and R. A. Naumann [Phys. Rev. Lett. 34, 1558 (1975)].

The agency support acknowledgement for two of the authors was omitted. The research of E. J. A. and R. A. N. was supported by the U. S. Energy Research and Development Administration.