

Dynamical Calculation of the  $A_1$  \*

D. D. Brayshaw†

Stanford Linear Accelerator Center, Stanford University, Stanford, California 94305

(Received 7 July 1975)

A relativistic three-body theory previously used to generate the  $\omega$  has been applied to the  $1^+$  state of three pions. An  $A_1$  resonance pole is produced with  $M_{A_1} = 1160$  MeV,  $\Gamma/2 = 90$  MeV, but there is no associated phase variation of the amplitude.

There has been growing concern over the embarrassments suffered by the quark model in predicting the meson spectroscopy, and in particular the experimental absence of  $1^+$  states such as the  $Q$  and  $A_1$ . Thus, the diffractively produced enhancement observed in  $\pi p \rightarrow (3\pi)p$  at 1.1 GeV has recently been subjected to a number of independent and rather sophisticated analyses, and all agree on the absence of a resonant  $A_1$  signal.<sup>1</sup> As developed by Ascoli and collaborators, the procedure is a variant of the isobar model in which one writes the amplitude for a three-body "decay" in the form  $T = \sum_{\alpha} f_{\alpha} t_{\alpha}$ , where  $t_{\alpha}$  is a two-body scattering amplitude and depends explicitly on the pair subenergy  $s_{\alpha}$ . The  $f_{\alpha}$  are treated as complex fitting parameters, and one subsequently studies their phase variation with respect to some (nonresonant) reference amplitude. This works quite nicely for the  $A_2$ , which exhibits Breit-Wigner behavior with the phase varying through  $90^\circ$ . In contrast, the  $A_1$  phase variation is quite flat, and one is apparently forced to a nonresonant interpretation. A recent attempt by Ascoli and Wyld to answer previous criticism of the methodology<sup>2</sup> has led to the same negative conclusion.<sup>3</sup>

In this Letter I show that an absence of distinctive phase variation is not only compatible with an  $A_1$  resonance pole, but (1) is an automatic consequence of a simple dynamical model. Moreover, we argue that such behavior (2) is a general feature of a properly unitarized amplitude, *provided the effect is dynamical*, and (3) has a simple physical interpretation. Specifically, we observe that a resonant amplitude need not exhibit a large change of phase, as assumed in these analyses. In fact, this is a familiar phenomenon associated with very *inelastic* resonances. For example, consider a system in which two orthogonal channels are dynamically coupled (e.g.,  $\pi\pi$  and  $K\bar{K}$ ); in such a system the elastic amplitudes have the form  $t_{\alpha} = [\eta \exp(2i\delta_{\alpha}) - 1]/2i$ . If the amplitudes are resonant and  $\eta < \frac{1}{2}$ , the phase shifts  $\delta_{\alpha}$  are roughly sinusoidal, pass-

ing through zero (rather than  $\pi/2$ ) when  $\text{Re}(t_{\alpha}) = 0$ . Typically, the magnitude of this oscillation is quite small ( $< 30^\circ$ ), in which case one may easily show that the phase of the amplitude itself exhibits a similar small oscillation about  $\pi/2$ . One would not expect to detect this in an Ascoli-type analysis, particularly in view of simplifying approximations (e.g., neglect of the subenergy dependence of the parameters  $f_{\alpha}$ ). We then observe that in the isobar approximation, the  $A_1$  state of three pions is precisely such a system, containing two strong competing channels ( $\rho\pi$  and  $\epsilon\pi$ ). Therefore, it would not be at all surprising if the relevant phase behavior were of the second, weaker type.

This conjecture is supported by explicit calculations based on the author's covariant boundary-condition formalism (BCF), which has recently been applied to a number of relativistic three-particle systems.<sup>4,5</sup> In the present context, we have used it to study the amplitude  $T_3 = \sum_{\alpha} \tau_{\alpha}$  describing  $3\pi$  scattering in a  $1^+$  ( $I=1$ ) state. We thus consider  $T(N\pi \rightarrow N3\pi)$  to be of the form  $T = T_p * T_3 + \dots$ , where  $T_p$  is an appropriate production amplitude. Provided that the  $A_1$  is indeed a dynamical effect, we would expect  $T_3$  to contain the corresponding resonance pole (rather than  $T_p$ ). For this purpose the BCF may be employed in two complementary ways. In its most general form, it provides a general solution of the three-particle unitarity relations, and hence any physical amplitude can be constructed given suitable input. Conversely, the class of allowable input exhausts the possible physical amplitudes, and thus one can determine whether *any* input which produces an  $A_1$  peak in  $T_3$  will also produce a large phase variation. The answer turns out to be "no," which is a model-independent result. Furthermore, the formalism permits an explicit analytic continuation onto the second sheet of the total ( $3\pi$ ) energy; in this way it has been verified that each such peak corresponds to an associated pole. Secondly, input to the BCF has a straightforward dynamical inter-

pretation, and can be estimated for a simple model which has previously been applied to calculate the  $\pi$  and  $\omega$  as dynamical  $3\pi$  effects.<sup>4</sup> It is therefore interesting that the same model "predicts" an  $A_1$  state of approximately the right mass and width (and with negligible phase variation). Together, these results indicate that the  $0^-$ ,  $1^-$ ,  $1^+$  states can be understood in terms of potentiallike forces generated by particle exchange.

As noted by Amado,<sup>6</sup> a minimal scheme for unitarizing three-particle amplitudes must take the form of a one-dimensional integral equation. The key ingredients of such an equation are well known; in order to produce the primary (model-independent) singularity structure, the kernel must contain a pole corresponding to the free

propagation of all three particles, and two-particle propagators characterized by elastic phase shifts. The minimal form of the BCF corresponds to a trivial dynamical model possessing all of these features; additional parameters in the general form correct the dynamical details via the addition of nonsingular terms. Although the BCF is particularly efficient for this type of analysis, it should be emphasized that our more general results [(2) and (3) above] will be a feature of any properly unitarized amplitude.

The BCF builds on a trivial model in which the pairwise interaction of  $\beta$ - $\gamma$  is compressed to the surface of an impenetrable boundary at  $|\vec{r}_\beta - \vec{r}_\gamma| = a_\alpha$ . The scattering is described by an energy-dependent logarithmic derivative of the wave function at that radius,  $\lambda_{\alpha l}(\kappa_\alpha^2)$ . The two-body amplitudes are then  $t_{\alpha l}(\kappa_\alpha) = N_{\alpha l}/D_{\alpha l}$ , with

$$\begin{aligned} N_{\alpha l}(\kappa_\alpha) &= [a_\alpha \lambda_{\alpha l}(\kappa_\alpha^2) - l] j_l(a_\alpha \kappa_\alpha) + a_\alpha \kappa_\alpha j_{l+1}(a_\alpha \kappa_\alpha), \\ D_{\alpha l}(\kappa_\alpha) &= i \kappa_\alpha \{ [a_\alpha \lambda_{\alpha l}(\kappa_\alpha^2) - l] h_l(a_\alpha \kappa_\alpha) + a_\alpha \kappa_\alpha h_{l+1}(a_\alpha \kappa_\alpha) \}. \end{aligned} \quad (1)$$

Here  $\kappa_\alpha$  is the c.m. momentum of the  $\beta$ - $\gamma$  pair;  $\lambda_{\alpha l}(\kappa_\alpha^2)$  and  $a_\alpha$  are fitted to scattering data in the physical region  $\kappa_\alpha^2 \geq 0$ . Since  $\lambda_{\alpha l}$  must be meromorphic in  $\kappa_\alpha^2$  in order to produce unitary (elastic) amplitudes, this fit permits analytic continuation of  $N_{\alpha l}, D_{\alpha l}$  for  $\kappa_\alpha^2 < 0$ . Below I use the notation  $N_{\alpha l}^c, D_{\alpha l}^c$  to denote  $N_{\alpha l}, D_{\alpha l}$  evaluated with  $\lambda_{\alpha l}(\kappa_\alpha^2)$  replaced by the constant value  $\lambda_{\alpha l}(-\mu^2)$ . If one considers the three-particle system with  $\beta$ - $\gamma$  separated by  $a_\alpha$  in their c.m. frame, it is obvious that there is a characteristic distance  $b_\alpha$  of particle  $\alpha$  from the  $\beta$ - $\gamma$  c.m. such that  $|\vec{r}_\alpha - \vec{r}_\beta| \leq a_\gamma$  and/or  $|\vec{r}_\alpha - \vec{r}_\gamma| \leq a_\beta$ ;  $b_\alpha$  characterizes the interior region in which the cores may overlap.

In the present application we take the  $1^+$  state to be composed of two components corresponding to  $l=1, \lambda=0$ , and  $l=0, \lambda=1$ , where  $\lambda$  is the angular momentum of the spectator pion. For convenience we label these by  $\rho$  and  $\epsilon$ , respectively, although an  $\epsilon$  pole was not explicitly assumed. The amplitudes  $\tau_\alpha(s, s_\alpha)$  may be computed from the solution of

$$X_i(s, s_i) = \Omega_i(s, s_i) + \sum_j \int_0^\infty dq_j q_j^2 K_{ij}(q_i', q_j; s) X_j(s, s_j), \quad (2)$$

written in terms of the three-momentum  $q_i$  of the spectator in the  $i$ th-pair c.m. frame (equivalent to  $s_i$  for fixed  $s$ ). Here  $i, j$  take on the values  $\rho, \epsilon$ . The relation of  $\tau$  to  $X$  is such that  $X_\rho(s, s_\rho)$  is to be compared with the isobar amplitude  $f_\rho$ , and  $\Omega_i(s, s_i)$  is an appropriate projection onto an initial  $3\pi$  state. However, these details are not required for our present purposes. A pole in  $\tau_\alpha$  (and hence  $T_3$ ) can only arise via a pole in the operator  $(1 - K)^{-1}$ ; this corresponds to a complex zero of the determinant  $D(s) \equiv |1 - K|$ . One can thus study the resonant properties by constructing  $D(s)$ ; the rapidly varying factor of an appropriate cross section is proportional to  $|D|^{-2}$ .

Consequently, we need only consider the kernel of Eq. (2). For the minimal equation this may be expressed as<sup>7</sup>

$$\begin{aligned} K_{ij}(q_i', q_j; s) &= \Lambda_{ij} \frac{N_{ij}^s(q_i', q_j)}{D_j(\kappa_j)} \frac{N_j(\kappa_j)}{N_j^c(\kappa_j)}, \\ N_{ij}^s(q_i', q_j) &= -\frac{2\kappa_j}{\pi} \int_{-1}^1 dz G_{ij}(z, \hat{K}_{ij} \cdot \hat{q}_j, \hat{Q}_{ij} \cdot \hat{q}_j) \frac{g_i(b_i q_i', b_i Q_{ij})}{q_i'^2 - Q_{ij}^2 - i\epsilon} \frac{N_i^c(K_{ij})}{Q_{ij}}. \end{aligned} \quad (3)$$

Here  $\Lambda_{ij}$  is an isospin recoupling coefficient ( $\Lambda_{\rho\rho} = \frac{1}{2}$ ,  $\Lambda_{\epsilon\rho} = -\Lambda_{\rho\epsilon} = \frac{1}{\sqrt{3}}$ ,  $\Lambda_{\epsilon\epsilon} = \frac{1}{3}$ ), and  $G_{ij}$  is a geometrical recoupling coefficient which would be unity if all particles were in relative  $s$  waves. The three-vectors  $\vec{K}_{ij}, \vec{Q}_{ij}$  are the values of  $\vec{k}_i, \vec{q}_i$  in the  $i$  c.m. system expressed in terms of  $\vec{k}_j, \vec{q}_j$  in the  $j$  frame, and  $z = \hat{k}_j \cdot \hat{q}_j$ . The function  $g_i$  arises from excluding the inner region;

$$g_i(\kappa, y) = iy [y j_\lambda(x) h_{\lambda+1}(y) - x j_{\lambda+1}(x) h_\lambda(y)]. \quad (4)$$

We note that  $g_i(x, x) = 1$ , and hence the residue of the integrand at the Green's function pole  $q_i' = Q_{ij}$  is given by the two-body amplitude  $t_j(\kappa_j)$ .

The general form of the BCF is obtained by replacing  $N_{ij}^s \rightarrow N_{ij}^s + A_{ij}$ , where  $A_{ij}(q_i', q_j) \cdot \kappa_j^{ij}$  is an arbitrary  $L_2$  function which must be real valued to describe elastic three-body scattering. As noted in Ref. 4, a rough estimate of  $A_{ij}$  can be derived if one assumes it to be dominated by off-shell corrections to the  $\rho\pi\pi$  vertex. This leads to the specific model

$$A_{\rho\rho}(q', q) = \frac{\gamma_{\rho\rho} \kappa_\rho^2 + 4\mu^2}{\kappa_\rho} g_\rho(q') g_\rho(q), \quad (5)$$

with  $g_\rho(q) = (q^2 + 4\mu^2)^{-1}$ ,  $\gamma_{\rho\rho} \approx \frac{1}{2}$  (similar estimates give  $A_{\epsilon\epsilon}, A_{\rho\epsilon}, A_{\epsilon\rho}$ , which are numerically unimportant). More generally, if potentiallike mechanisms dominate, one expects  $A_{ij}$  to have a relatively weak dependence on  $s$ , and to be a smooth function of  $q', q$ .<sup>8</sup> It can then be expanded in a complete set  $A = \sum_\lambda c_\lambda |\lambda\rangle \langle \lambda|$ , and the  $c_\lambda$  treated as real fitting parameters. A number of such forms were employed to test the model dependence of our result.

The required numerical procedures are straightforward: One distorts the  $q_j$ -integration contour to avoid the singularity at  $q_i' = Q_{ij}$ , and employs Gaussian quadrature to reduce the equation to finite matrix form in order to calculate  $D(s)$ . To simplify the numerics a cutoff was employed at  $q_j^{\text{max}} = 30 \text{ fm}^{-1}$ ; the calculation was quite insensitive to this choice (a 1% effect for  $25 \text{ fm}^{-1} < q_j^{\text{max}} < 35 \text{ fm}^{-1}$ ). Several choices of  $s$ - and  $p$ -wave  $\pi$ - $\pi$  phases were employed corresponding to the range of models reported by Basdevant, Froggatt, and Petersen,<sup>9</sup> as well as a simple  $s$  wave which does not exhibit the rapid change of phase at the  $K\bar{K}$  threshold (no  $S^*$ ). In practice, the  $S^*$  region turns out to be relatively unimportant since it requires a very small spectator momentum ( $q_j \approx 0$ ), and this is suppressed both by the  $\lambda = 1$  character and the  $q_j^2 dq_j$  integration weight.

Given this input and the simple model of Eq. (5), a  $1^+$  resonance is indeed generated in the vicinity of 1100 MeV for  $\gamma_{\rho\rho}$  in the estimated range. A typical example is illustrated in Fig. 1 (solid curve), corresponding to  $\gamma_{\rho\rho} = 0.57$ . Writing  $T = N/D(s)$ , this result would imply a width of 220 MeV if the  $s$  dependence of  $N$  were negligible. It is clear that the phase  $\varphi(D)$ , which would normally signal the presence of simple Breit-Wigner behavior, exhibits no noticeable variation




FIG. 1. Dependence of  $|D|^{-2}$  (upper figure) and  $\varphi(D)$  (lower figure) on the three-pion mass. The curves correspond to complex energies  $\sqrt{s} = M_{3\pi} - i\Delta$  with  $\Delta = 0, 30, 60$  MeV for the solid, dashed, and dash-dotted curves, respectively.

associated with the enhancement. Since the fitting parameters ( $\gamma_{\rho\rho}$ , or more generally the  $c_\lambda$ ) carry only weak  $s$  dependence (in the absence of inelastic thresholds), one can hold them fixed and perform an explicit analytic continuation. In this way we confirm the existence of a pole 90 MeV below the real axis on the second sheet, with a mass of 1160 MeV. Although it was possible to vary  $A$  in such a way as to produce no peak, it was found that peak, pole, and minor phase variation were always correlated.

With regard to the proposed "inelastic" mechanism, it is very suggestive that as we take  $\sqrt{s}$  deeper onto the second sheet,  $\varphi(D)$  increasingly takes on the characteristic appearance of such a resonance (dashed curves of Fig. 1). We note that by doing so we also approach closer to the  $\rho$  and  $\epsilon$  poles which occur in the factor  $D_j^{-1}(\kappa_j)$ , and hence more closely approach a coupled-channel problem involving stable "particles." Numerical studies confirm that the interplay between the  $\rho$  and  $\epsilon$  channels is vital in producing the effect (whereas an  $S^*$  pole is not required).

On the other hand, the limitations of the isobar model are apparent in the damping of the effect for real  $\sqrt{s}$ . Thus, one cannot escape the fact that we are dealing with *three* particles, with an associated three-particle cut as well as  $\rho$  and  $\epsilon$  thresholds. The net  $s$  dependence is an integrated product of these factors, and is necessarily quite complicated; this shows up both in the non-descript phase behavior and in the shape of the bump in  $|D|^{-2}$  (solid curves). We conclude that such an  $A_1$  cannot be established by an Ascoli-type analysis, and may well be present in the data.

\*Work supported in part by the U. S. Energy Research and Development Administration.

†Alfred P. Sloan Foundation Fellow.

<sup>1</sup>Yu. M. Antipov *et al.*, Nucl. Phys. **B63**, 153 (1973); G. Ascoli *et al.*, Phys. Rev. D **7**, 669 (1973); M. Tabak

*et al.*, in *Experimental Meson Spectroscopy—1974*, AIP Conference Proceedings No. 21, edited by D. A. Garelick (American Institute of Physics, New York, 1974).

<sup>2</sup>R. Aaron and R. D. Amado, Phys. Rev. Lett. **31**, 1157 (1973).

<sup>3</sup>G. Ascoli and H. W. Wyld, Phys. Rev. D **12**, 43 (1975).

<sup>4</sup>D. D. Brayshaw, Phys. Rev. D **11**, 2583 (1975).

<sup>5</sup>D. D. Brayshaw, Phys. Rev. C **11**, 1196 (1975); D. D. Brayshaw and H. Pierre Noyes, Phys. Rev. Lett. **34**, 1582 (1975).

<sup>6</sup>R. D. Amado, Phys. Rev. Lett. **33**, 333 (1974).

<sup>7</sup>The general form of this equation is given in operator notation in Ref. 4, as well as its derivation. It is shown that the explicit form differs from the nonrelativistic version only by the kinematics, and hence one may compare the expressions given in D. D. Brayshaw, Phys. Rev. D **8**, 952 (1973).

<sup>8</sup>D. D. Brayshaw, SLAC Report No. SLAC-PUB-1648, September 1975 (to be published).

<sup>9</sup>J. L. Basdevant, C. D. Froggatt, and J. L. Petersen, Phys. Lett. **41B**, 178 (1972).

## Measurement of the Reaction $e^+e^- \rightarrow \gamma\gamma$ in the Region of the $\psi(3684)^*$

E. B. Hughes, B. L. Beron, R. L. Carrington, R. L. Ford, E. Hilger,† R. Hofstadter, A. D. Liberman, T. W. Martin, L. H. O'Neill, and J. W. Simpson  
*High Energy Physics Laboratory and Department of Physics, Stanford University, Stanford, California 94305*

and

L. K. Resvanis  
*Department of Physics, University of Pennsylvania, Philadelphia, Pennsylvania 19104*  
(Received 20 October 1975)

In a study of the reaction  $e^+e^- \rightarrow \gamma\gamma$  in the region of the  $\psi(3684)$  resonance, no evidence is found for a deviation from quantum electrodynamics. Limits are also set on the decays  $\psi(3684) \rightarrow \pi^0\gamma$ ,  $\psi(3684) \rightarrow \eta^0\gamma$ , and  $\psi(3684) \rightarrow X^0\gamma$ , where  $X^0$  is a pseudoscalar state in the mass range 2500–3000 MeV which subsequently decays into two  $\gamma$  rays.

The purpose of this Letter is to report the result of a measurement of the reaction  $e^+e^- \rightarrow \gamma\gamma$  at center-of-mass energies close to the  $\psi(3684)$  resonance. If this resonance has the quantum numbers of the photon,  $J^{PC} = 1^{--}$ , as indicated by the destructive interference and decay angular distribution observed in the channel  $e^+e^- \rightarrow \mu^+\mu^-$ ,<sup>1</sup> then the event rate for the reaction  $e^+e^- \rightarrow \gamma\gamma$  should be solely that expected from quantum electrodynamics (QED). Apparent violations of this expectation could be observed in the present experiment if the decays  $\psi(3684) \rightarrow \pi^0\gamma$  or  $\psi(3684) \rightarrow \eta^0\gamma$  were to occur, or if the decay  $\psi(3684) \rightarrow X^0\gamma$ ,  $X^0 \rightarrow \gamma\gamma$  were to exist,<sup>2</sup> where  $X^0$  is a pseudoscalar

state with a mass in the range 2500–3000 MeV.

The experiment was done at the electron-positron storage ring (SPEAR) at the Stanford Linear Accelerator Center using an apparatus which has been described before,<sup>3</sup> consisting of two identical spectrometers mounted in a collinear configuration about the beam interaction region. For the detection of  $\gamma$  rays each spectrometer contains three multiwire proportional chambers (MWPC's) with space for a lead converter between the first two, and a 20-radiation-length-thick NaI(Tl) total-absorption detector 30 in. in diameter. Throughout the experiment the spectrometers were set to accept  $\gamma$  rays produced at