

depends on both the magnitude and sign of G_E/G_M . Unpolarized elastic scattering experiments determine G_E^2 and G_M^2 , but not the sign of G_E/G_M . For $Q^2 = 0.765$ (GeV/c)² these experiments¹⁰ give $|\mu G_E/G_M| = 0.98 \pm 0.04$ in which $\mu = 2.79$. If G_E and G_M have the same sign, Eq. (2) yields $A = +0.112 \pm 0.001$, while if G_E and G_M have the opposite sign Eq. (2) gives $A = -0.017 \pm 0.002$. From our measured value of A we conclude that the theoretical and experimental values are in good agreement provided the signs of G_E and G_M are the same. The effect of proton structure on the hyperfine-structure interval in hydrogen involves an integral of the product of the proton structure functions and also gives the sign of G_E/G_M to be positive.¹¹

The experimental method described in this Letter could in principle^{2,12} be applied to determine G_E in the region $Q^2 \lesssim 2$ (GeV/c)², where G_E is not well known, but its practical usefulness is limited by low counting rates.

We are happy to acknowledge the important contributions to this experiment by M. Browne, S. Dhawan, R. Eisele, Z. Farkas, R. Fong-Tom, H. Hogg, E. Garwin, R. Koontz, J. Sodja, S. St. Lorant, J. Wesley, and M. Zeller.

*Research supported in part by the U. S. Energy Research and Development Administration under Contract No. E(11-1)-3075 (Yale) and Contract No. E(04-3)-515,

(Stanford Linear Accelerator Center), the German Federal Ministry of Research and Technology, and the University of Bielefeld, the Japan Society for the Promotion of Science, and the National Science Foundation.

¹M. J. Alguard *et al.*, following Letter [Phys. Rev. Lett. **37**, 1261 (1976)].

²N. Dombey, Rev. Mod. Phys. **41**, 236 (1969). The methodology of this paper was used to derive Eq. (2).

³V. W. Hughes *et al.*, Phys. Rev. A **5**, 195 (1972); M. J. Alguard *et al.*, in *Proceedings of the Ninth International Conference on High Energy Accelerators, Stanford Linear Accelerator Center, Stanford, California, 1974*, CONF-740 522 (National Technical Information Service, Springfield, Va., 1974), p. 309.

⁴P. S. Cooper *et al.*, Phys. Rev. Lett. **34**, 1589 (1975).

⁵We thank the members of SLAC Group A for their invaluable help in making the electron polarization measurement by Møller scattering in March 1976.

⁶M. J. Alguard *et al.*, Bull. Am. Phys. Soc. **21**, 98 (1976).

⁷W. W. Ash, in Proceedings of the Brookhaven National Laboratory Workshop in Physics with Polarized Targets, June 1974, BNL Report No. 20415 (unpublished), p. 309; W. W. Ash *et al.*, to be published.

⁸M. Borghini *et al.*, Nucl. Instrum. Methods **84**, 168 (1970).

⁹Isolite is the name for a plastic formed by adding a wavelength-shifter to Lucite.

¹⁰Ch. Berger *et al.*, Phys. Lett. **35B**, 87 (1971);

W. Bartel *et al.*, Nucl. Phys. B **58**, 429 (1973).

¹¹H. Grotch and D. R. Yennie, Rev. Mod. Phys. **41**, 350 (1969); C. Zemach, Phys. Rev. **104**, 1771 (1956).

¹²A. I. Akhiezer *et al.*, Zh. Eksp. Teor. Fiz. **33**, 765 (1957) [Sov. Phys. JETP **6**, 588 (1958)].

Deep Inelastic Scattering of Polarized Electrons by Polarized Protons*

M. J. Alguard, W. W. Ash, G. Baum, J. E. Clendenin, P. S. Cooper, D. H. Coward, R. D. Ehrlich, A. Etkin, V. W. Hughes, H. Kobayakawa, K. Kondo, M. S. Lubell, R. H. Miller, D. A. Palmer, W. Raith, N. Sasao, K. P. Schüler, D. J. Sherden, C. K. Sinclair, and P. A. Souder
University of Bielefeld, Bielefeld, West Germany, and City University of New York, New York, New York 10031, and Nagoya University, Nagoya, Japan, and Stanford Linear Accelerator Center, Stanford, California 94305, and University of Tsukuba, Ibaraki, Japan, and Yale University, New Haven, Connecticut 06520

(Received 5 August 1976)

We report measurements of the asymmetry in deep inelastic scattering of longitudinally polarized electrons by longitudinally polarized protons. The antiparallel-parallel asymmetries are positive and large in agreement with predictions of quark-parton models of the proton. A limit is obtained on parity nonconservation in the scattering of longitudinally polarized electrons by unpolarized nucleons.

Experimental and theoretical studies of deep inelastic electron scattering from protons and neutrons have led in the past eight years to the important discovery of scaling and to the quark-par-

ton model of nucleon structure.¹ Deep inelastic muon² and neutrino³ scattering have confirmed these general ideas.⁴

For deep inelastic electron-proton scattering,

accurate data have been obtained on the differential cross section $d^2\sigma/d\Omega dE'$ over a wide range of the energy loss, ν , of the electron and the square of the four-momentum transfer, q^2 , to the proton. The two spin-averaged proton structure functions $W_1(\nu, q^2)$ and $W_2(\nu, q^2)$ have been determined from these data. Important, independent information is contained in two additional spin-dependent proton structure functions whose determination requires the measurement of spin correlation asymmetries.⁵

In this Letter we report the first results of an experiment done at the Stanford Linear Accelerator Center (SLAC) to measure the asymmetry, A , in the deep inelastic scattering of longitudinally polarized electrons by longitudinally polarized

protons, where A is given by

$$A = [d\sigma(\uparrow\uparrow) - d\sigma(\uparrow\downarrow)] / [d\sigma(\uparrow\uparrow) + d\sigma(\uparrow\downarrow)], \quad (1)$$

with $d\sigma$ denoting the differential cross section $d^2\sigma(E, E', \theta) / d\Omega dE'$ for electrons of incident (scattered) energy E (E') and laboratory scattering angle θ , and the arrows denoting the antiparallel and parallel spin configurations.

If the scattering is described by the one-photon-exchange approximation, then for unpolarized electrons the virtual photons are linearly polarized, whereas for polarized electrons the photons are elliptically polarized. The differential cross section for the scattering of longitudinally polarized electrons by longitudinally polarized protons is

$$\frac{d^2\sigma}{d\Omega dE'} = \left(\frac{d\sigma}{d\Omega} \right)_M \left(\frac{1}{\epsilon(1 + \nu^2/Q^2)} \right) W_1 \{ 1 + \epsilon R \pm (1 - \epsilon^2)^{1/2} \cos\psi A_1 \pm [2\epsilon(1 - \epsilon)]^{1/2} \sin\psi A_2 \}, \quad (2)$$

in which $(d\sigma/d\Omega)_M$ is the Mott differential cross section, $\epsilon = [1 + 2(1 + \nu^2/Q^2) \tan^2 \frac{1}{2}\theta]^{-1}$, $Q^2 = -q^2$, $R = \sigma_L/\sigma_T$ is the ratio of the cross sections for absorption of longitudinal and transverse virtual photons, and ψ is the angle between the directions of the virtual photon momentum and the proton spin. The + (-) signs in Eq. (2) refer to the antiparallel (parallel) spin configurations.

The spin-dependent terms A_1 and A_2 are two new measurable quantities which can be expressed in terms of two spin-dependent structure functions.^{5,6} Equivalently, they can be expressed in terms of the total absorption cross sections of circularly polarized photons on polarized protons as

$$\begin{aligned} A_1 &= (\sigma_{1/2} - \sigma_{3/2}) / (\sigma_{1/2} + \sigma_{3/2}), \\ A_2 &= 2\sigma_{TL} / (\sigma_{1/2} + \sigma_{3/2}), \end{aligned} \quad (3)$$

where $\sigma_{1/2}$ ($\sigma_{3/2}$) is the total absorption cross section when the z component (z is the direction of the virtual photon momentum) of angular momentum of the virtual photon plus proton is $\frac{1}{2}$ ($\frac{3}{2}$), and σ_{TL} , which may be negative, is a term which arises from the interference between transverse and longitudinal photon-nucleon amplitudes. It should be noted that $\sigma_{1/2}$ and $\sigma_{3/2}$ are related to σ_T by $\sigma_{1/2} + \sigma_{3/2} = 2\sigma_T$.

For the case of protons polarized along the incident beam direction, the asymmetry A of Eq. (1) is

$$A = D(A_1 + \eta A_2), \quad (4)$$

where

$$\begin{aligned} D &= (E - E'\epsilon) / E(1 + \epsilon R) \\ &= (1 - \epsilon^2)^{1/2} \cos\psi / (1 + \epsilon R), \end{aligned} \quad (5)$$

and

$$\begin{aligned} \eta &= \epsilon(Q^2)^{1/2} / (E - E'\epsilon) \\ &= [2\epsilon / (1 + \epsilon)]^{1/2} \tan\psi \simeq \tan\psi. \end{aligned} \quad (6)$$

The quantity D can be regarded as a kinematic depolarization factor of the virtual photon and is ~ 0.3 for our kinematic points. Positivity limits imposed on A_1 and A_2 are⁷

$$|A_1| \leq 1, \quad |A_2| \leq \sqrt{R}. \quad (7)$$

In this experiment we determine the combination $A_1 + \eta A_2$ by dividing the measured electron-proton asymmetry A by the depolarization factor D . Although we do not separately determine A_1 and A_2 , our result is dominated by A_1 because the kinematic factor η is small.

On the basis of a high-energy sum rule derived with the algebra of currents for a quark model, it has been predicted⁸ that A_1 has a positive value greater than 0.2 over a large region of the deep inelastic continuum. Scaling relations are predicted for the spin-dependent proton structure functions, and hence also for A_1 ⁹:

$$A_1(\nu, Q^2) \rightarrow A_1(\omega) \text{ as } \nu, Q^2 \rightarrow \infty, \text{ with } \omega \text{ held constant} \quad (8)$$

($\omega = 2M\nu/Q^2$, M is the proton mass). Specific models of proton structure make widely varying predictions for A_1 . The simplest quark-parton

TABLE I. Results of asymmetry measurements.

E (GeV)	θ (deg)	Q^2 [(GeV/c) ²]	W^a (GeV)	ω	Δ (%)	A^b	D^c	$A_1 + \eta A_2^b$	$ \eta A_2 $
9.711	9.000	1.680	2.059	3	0.44 ± 0.11	0.191 ± 0.057 (0.044)	0.284	0.67 ± 0.20 (0.16)	< 0.146
12.948	9.000	2.735	2.519	3	0.50 ± 0.17	0.215 ± 0.089 (0.080)	0.352	0.61 ± 0.25 (0.23)	< 0.109
9.711	9.000	1.418	2.560	3	0.28 ± 0.11	0.141 ± 0.058 (0.051)	0.412	0.34 ± 0.14 (0.12)	< 0.087

^a W is the missing mass of undetected hadron system.

^bThe total errors are the statistical counting errors added in quadrature to the systematic errors in P_e , P_p , and F ; the numbers in parentheses are the 1-standard-deviation counting errors.

^c D is obtained from Eq. (5) using $R=0.14$.

model predicts that $A_1 = \frac{5}{9}$, and more elaborate models also predict large positive values for $A_1(\omega)$.^{5,10}

The method of measuring the experimental asymmetry, Δ , for deep inelastic electron-proton scattering was the same as that described for elastic scattering in the preceding Letter.¹¹ For the inelastic case, the scattered electron counting rate was lower (0.02 to 0.06 electrons per pulse).

TABLE II. False asymmetries.^a

Combination of miniruns	Average asymmetry ^b (%)	$\chi^2(0)$ per degree of freedom
	$\omega = 3, Q^2 = 1.680$	
<u>(1234) - (5678)</u> (1234) + (5678)	0.04 ± 0.11	18/34
<u>(1357) - (2468)</u> (1357) + (2468)	-0.04 ± 0.11	38/34
<u>(2367) - (1458)</u> (2367) + (1458)	+0.14 ± 0.11	27/34
	$\omega = 3, Q^2 = 2.735$	
<u>(1234) - (5678)</u> (1234) + (5678)	-0.30 ± 0.17	33/30
<u>(1357) - (2468)</u> (1357) + (2468)	-0.03 ± 0.17	26/30
<u>(2367) - (1458)</u> (2367) + (1458)	+0.24 ± 0.017	40/30
	$\omega^2 = 5, Q^2 = 1.418$	
<u>(1234) - (5678)</u> (1234) + (5678)	-0.12 ± 0.11	34/35
<u>(1357) - (2468)</u> (1357) + (2468)	-0.10 ± 0.11	34/35
<u>(2367) - (1458)</u> (2367) + (1458)	-0.03 ± 0.11	30/35

^aSee preceding Letter (Ref. 11) for definitions of false asymmetries.

^bIrrespective of sign of target polarization.

Background due to misidentified pions was again negligible.

The antiparallel-parallel asymmetry Δ was measured for three deep inelastic kinematic points and the results are given in Table I. Several false asymmetries were also measured and are listed in Table II, together with the χ^2 values for the agreement with zero of the measured false asymmetries for the indicated degrees of freedom (number of individual runs). No statistically significant false asymmetries were found.

The asymmetry A of Eq. (1) is related to Δ by

$$\Delta = P_e P_p F A. \quad (9)$$

The electron polarization, P_e , was 0.51 ± 0.06 , and the average target polarization, P_p , measured for each kinematic point, was ≈ 0.40 with 10% uncertainty. The quantity F is the fraction of detected electrons scattered from free protons. This is taken as the ratio of the number of free protons to the total number of nucleons in the target, including measured contributions from helium and other background sources. A small correction for the difference in scattering cross sections of neutrons and protons was also included. The value for F , determined for each point, was ≈ 0.11 with a 10% uncertainty.

The measured values of A are listed in Table I. The uncertainties are dominated by counting statistics. No radiative corrections have yet been made. Also listed are the quantities D (evaluated using $R=0.14$),¹ $A/D = A_1 + \eta A_2$, and upper limits for $|\eta A_2|$ (taking $A_2 = \sqrt{R}$). From Table I it is seen that A/D is dominated by A_1 . Furthermore, parton theories predict¹² that the interference term A_2 will be considerably smaller than its positivity limit \sqrt{R} . It is therefore valid to compare our measured value of A/D to theoretical predictions for A_1 as shown in Fig. 1.

With the explicit assumption that $A/D = A_1$, our




FIG. 1. Experimental values of $A/D \approx A_1$ and theoretical predictions of the virtual-photon-proton asymmetry A_1 versus ω . Theoretical curves a , b , c , and d are obtained from Refs. 5, 10, 13, and 14, respectively. For curve c the quark model with symmetry breaking is used: The model does not give values for A_1 in the range $1 < \omega < 2$, but rather gives $A_1(1) = 1$. For curve d the quantity μ^2/m_p^2 in the theory is taken equal to 0.12.

values of A_1 are indeed positive and large in accord with early theoretical expectations from sum rules.⁸ The two values for $\omega = 3$ agree within their errors, which is consistent with the expectation that A_1 satisfies the scaling relation, given by Eq. (8). Our data are consistent with the predictions of the quark-parton models shown as curves a ⁵ and b ¹⁰ in Fig. 1, but disagree strongly with the resonance model¹³ (curve c) and the bare-nucleon-bare-meson model¹⁴ (curve d). We note that the theoretical curves are all given for the scaling limit.

Data from this experiment can also be used to place a limit on parity nonconservation in the scattering of longitudinally polarized electrons from unpolarized nucleons, i.e., an interaction term of the form $\vec{\sigma}_e \cdot \vec{p}_e$ in which $\vec{\sigma}_e$ is the electron spin and \vec{p}_e is the electron incident momentum. If we define Δ^+ (Δ^-) as the asymmetry for protons polarized along (against) the beam direction and if the magnitude of P_p is the same for both cases, then we can define an asymmetry, Δ_{PNC} , associated with parity nonconservation by¹⁵

$$\Delta_{\text{PNC}} = (\Delta^+ - \Delta^-)/2 \equiv rP_e, \quad (10)$$

in which $r = (d\sigma^- - d\sigma^+)/ (d\sigma^- + d\sigma^+)$ is the asymmetry for electron polarization $P_e = 1$, and the minus

and plus superscripts refer to the electron beam helicity. From the deep inelastic scattering data summarized in Table I for Q^2 between 1.4 and 2.7 (GeV/c)², we find that r is consistent with zero. For the combined data we have an upper limit of $r < 5 \times 10^{-3}$ with a 95% confidence level. For the elastic scattering data reported in the preceding Letter,¹¹ again r is consistent with zero and its upper limit is less than 3×10^{-3} with a 95% confidence level. The gauge theories of weak and electromagnetic interactions, which contain parity nonconservation, predict^{16,17} considerably smaller values of $r \approx (10^{-5} \text{ to } 10^{-4})Q^2/M^2$.

We are happy to acknowledge helpful and stimulating discussions with J. D. Bjorken, F. Gilman, and J. Kuti.

*Research supported in part by the U. S. Energy Research and Development Administration under Contract No. E(11-1)-3075 (Yale) and Contract No. E(04-3)-515 (Stanford Linear Accelerator Center), the German Federal Ministry of Research and Technology and the University of Bielefeld, the Japan Society for the Promotion of Science, and the National Science Foundation.

¹R. E. Taylor, in *Proceedings of the International Symposium on Lepton and Photon Interactions at High Energies, Stanford, California, 1975*, edited by W. T. Kirk (Stanford Linear Accelerator Center, Stanford, Calif., 1975), p. 679. See also references therein.

²L. Mo, in *Proceedings of the International Symposium on Lepton and Photon Interactions at High Energies, Stanford, California, 1975*, edited by W. T. Kirk (Stanford Linear Accelerator Center, Stanford, Calif., 1975), p. 651.

³D. H. Perkins, in *Proceedings of the International Symposium on Lepton and Photon Interactions at High Energies, Stanford, California, 1975*, edited by W. T. Kirk (Stanford Linear Accelerator Center, Stanford, Calif., 1975), p. 571.

⁴C. H. Llewellyn-Smith, in *Proceedings of the International Symposium on Lepton and Photon Interactions at High Energies, Stanford, California, 1975*, edited by W. T. Kirk (Stanford Linear Accelerator Center, Stanford, Calif., 1975), p. 709. See also references therein.

⁵J. Kuti and V. F. Weisskopf, *Phys. Rev. D* **4**, 3418 (1971).

⁶F. Gilman, *Phys. Rep.* **4**, 95 (1972); F. Gilman, SLAC Report No. SLAC-167, 1973 (unpublished), Vol. I, p. 71.

⁷M. G. Doncel and E. de Rafael, *Nuovo Cimento* **4A**, 363 (1971).

⁸J. D. Bjorken, *Phys. Rev. D* **1**, 1376 (1970).

⁹L. Galfi *et al.*, *Phys. Lett.* **31B**, 465 (1970).

¹⁰F. Close, *Nucl. Phys.* **B80**, 269 (1974), and references therein.

¹¹M. J. Alguard *et al.*, preceding Letter [*Phys. Rev.*

Lett. **37**, 1258 (1976)].

¹²J. D. Bjorken and F. Gilman, private communication.

¹³G. Domokos *et al.*, Phys. Rev. D **3**, 1191 (1971).

¹⁴S. D. Drell and T. D. Lee, Phys. Rev. D **5**, 1738 (1972).

¹⁵In the actual analysis, target polarization differences were included. Since $P_p FA \approx 0.01$ is small, these differences have little effect.

¹⁶S. M. Berman and J. R. Primack, Phys. Rev. D **9**, 2171 (1974).

¹⁷G. Feinberg, Phys. Rev. D **12**, 3575 (1975).

Observation of a Population Inversion in a Possible Extreme Ultraviolet Lasing System

R. J. Dewhurst,* D. Jacoby, G. J. Pert, and S. A. Ramsden

Department of Applied Physics, University of Hull, Hull HU6 7RX, England

(Received 14 June 1976)

Spectrographic observation of a population inversion between the levels $n=3$ and $n=2$ in a rapidly expanding C VI plasma is reported. These and other results are shown to be in good agreement with a computer model, which is used to predict the conditions required to obtain laser action at 182 Å.

For some time now there has been considerable interest in obtaining laser action at extreme ultraviolet and x-ray wavelengths and a large number of schemes by which this might be achieved have been proposed.^{1,2} Although Jaeglé *et al.*³ have reported gain at 114 Å on the $2p^5 4d^3 P_1 - 2p^6 S_0$ intercombination line in Al^{3+} , the mechanism producing the inversion is unclear and the interpretation of the results has been disputed by other workers.⁴ A population inversion in the extreme ultraviolet region has been observed in a laser produced carbon plasma by Irons and Peacock⁵ but at densities too low to obtain useful gain.

In this work we have achieved a population inversion of the $n=3$ to $n=2$ transition of C VI at 182 Å, using the recombination scheme,⁶ which is sufficiently high to allow a laser to be constructed. The basic difficulty in this scheme is the need to obtain extremely rapid expansion and cooling of a fully ionized plasma⁷; recombination then leads to a preferential population of the upper states and hence to an inversion. In this work we have used very thin cylindrical carbon fibers irradiated in vacuum by a Nd glass laser; such targets allow a limited number of particles to be uniformly heated which then rapidly expand in all directions.

The carbon fiber, 5.3 μm in diameter, was supported vertically at the focus of the laser beam 13 cm in front of the horizontal entrance slit of a 2-m grazing incidence spectrograph (Hilger and Watts type E580). The spectrograph was carefully calibrated for absolute intensity measurements over the wavelength range 10–100 Å using the calibrated x-ray source and proportional counter

system described by Morgan, Gabriel and Barton⁸ and it is of interest to note that the calibration was practically identical to that reported for similar instruments by both Morgan, Gabriel, and Barton,⁸ and Hobby and Peacock.⁹ Irons and Peacock¹⁰ have also demonstrated that in this wavelength range, there is no reciprocity failure for the photographic emulsion used (Ilford Q2). An auxiliary vertical slit between the fiber and the spectrograph gave a spatial resolution of 100 μm. Without this slit, spectra could be obtained with a single laser shot, whereas with it in position thirty shots were needed to obtain a satisfactory exposure.

The laser was a conventional Nd-glass system giving up to 0.5 J in a pulse of 140-ps duration although energies of only 150 mJ were used in these experiments. Streak-camera measurements¹¹ showed that the pulse width was reproducible to ± 10 ps and that there were no spurious or satellite pulses. The single-pulse extinction ratio was at least 10^{-4} and the fiber was further protected from spontaneous emission and stray flash-lamp light by a saturable absorber. The laser energy was reproducible to ± 30 mJ for about 75% of the shots; on the remaining 25% there was an oscillator malfunction leading to a very weak output pulse. The contribution of these weak pulses in multishot experiments was less than the 10% of the total plate intensity and can be neglected; this is confirmed by the good agreement of single-shot and multishot intensity ratios of the Lyman lines.

We have found it necessary to use a small prepulse to break the fiber and form a dense cold plasma with which the main pulse interacts. Since