

in the $K^{\mp}\pi^{\pm}\pi^{\pm}$ mass spectrum does not contain 1.865-GeV/ c^2 $K^{\mp}\pi^{\pm}$ combinations and its recoil mass spectrum does not contain any clear structure. We do not consider it significant.

⁶J. D. Bjorken and S. L. Glashow, *Phys. Lett.* **11**, 255 (1964); S. L. Glashow, J. Iliopoulos, and L. Maiani,

Phys. Rev. D **2**, 1285 (1970); M. K. Gaillard, B. W. Lee, and J. L. Rosner, *Rev. Mod. Phys.* **47**, 277 (1975); A. De Rújula, H. Georgi, and S. L. Glashow, *Phys. Rev. Lett.* **37**, 398 (1976); K. Lane and E. Eichten, Cornell University Report No. COO-2220-78, 1976 (to be published).

Dimuon Production on Nuclear Targets

M. Binkley, I. Gaines, J. Peoples, B. Knapp, W. Lee, P. Leung, S. D. Smith, A. Wijangco, J. Knauer, J. Bronstein, R. Coleman, G. Gladding, M. Goodman, M. Gormley, R. Messner, T. O'Halloran, J. Sarracino, and A. Wattenberg
Fermi National Accelerator Laboratory, Batavia, Illinois 60510, and
 Columbia University, Irvington-on-Hudson, New York 10533,† and
 University of Hawaii, Honolulu, Hawaii 96822,‡ and
 University of Illinois, Urbana, Illinois 61801**
 (Received 17 May 1976)

We have measured the production of $\rho + \omega$ and J vector mesons, through their dimuon decay modes, by high-energy neutrons on nuclear targets at Fermilab. We determined the A dependence to be $A^{0.62 \pm 0.03}$ for $\rho + \omega$, $A^{0.93 \pm 0.04}$ for J , and $A^{0.85 \pm 0.05}$ for the continuum of dimuons between the $\rho + \omega$ and J masses.

Several recent experiments at Fermilab, the CERN intersecting storage rings, and Serpukhov have extended the study of dilepton production in high-energy hadron-hadron collisions.¹⁻³ These experiments showed the presence of two strong resonances above a continuum, one due to the $\rho + \omega$ and the other due to the J .⁴ Between these two masses the cross section decreases monotonically as the mass increases. Because of the relevance of dilepton production to several open experimental questions, such as the nature of J production in hadronic collisions, the contribution of dileptons to single-lepton production, and the possible existence of the Drell-Yan process⁵ for dilepton production, it is important to measure the features of dilepton production in greater detail than in previous experiments.

The primary purpose of this experiment was to measure the A dependence of the total inclusive cross section for $\rho + \omega$, J , and the continuum between those resonances in neutron-nucleus collisions. The second objective, which is discussed in a subsequent paper, was to determine the p_{\perp} and X_f dependence [$X_f = (p/p_{\max})_{nA \text{ c.m.}}$] of the invariant cross section for this process. By making measurements on four elements it was possible to extrapolate the yields to the cross section for production on a single nucleon. The details of the beam and the apparatus were reported in a previous publication.¹ The neutron beam was produced by the interaction of 400-GeV protons,

rather than 300-GeV protons as in our previously published work.

We detected muon pairs only after they emerged from an absorber as shown in Fig. 1. The target was placed close to the absorber in order to limit the number of muon pairs from meson decays. During this run the absorber consisted of 144 g/cm² of Be, a steel-scintillator hadron calorimeter of mass 530 g/cm², and an additional 760 g/cm² of uninstrumented steel. The absorber was located 78 cm downstream from the target. The bulk of the data which were used to determine the A dependence were obtained using Be, Al, Cu, and Pb targets which were 15 cm long and which had a mass per unit area of 28.5 g/cm². They were segmented in order to obtain equal length and density. Data were also taken with a solid Pb




FIG. 1. Schematic layout of experimental apparatus. A, T, and D are scintillation counters; H, V, μH , and μV are scintillation counter hodoscopes. P1, P2, P3, and P4 are multiwire proportional chambers.

target of 86.5 g/cm² and a solid Cu target of 54 g/cm².

A neutron interaction in either the target or the absorber would generate an event trigger if there were a coincidence between the D counter and either two H counters and one V counter or two V counters and one H counter. The information on the event was recorded on tape if either two μ_H counters and one μ_V counter or two μ_V counters and one μ_H counter had counts in coincidence with the event trigger. The A counter was required to be off. Two or more hits were required in two of the three planes of proportional chambers P_2 , P_3 , and P_4 . The target counter T was not used in the event trigger, although its status and the status of all other counters were recorded with the multiwire-proportional-chamber information.

Events which had two or more tracks extending from P_1 through P_4 were candidates for muon pairs. The distance of closest approach of the tracks was required to be less than 2.2 cm. The interaction point, the midpoint on the line segment connecting the two tracks at their distance of closest approach, was required to be within 2.5 cm of the axis of the neutral beam. The event was classified as a dimuon if three of the four muon counters which were intercepted by the tracks had counts.

The invariant mass of the pair was calculated in two ways. In both calculations the magnitudes of the momenta were determined from the deflection of the tracks in the magnet and then corrected for the energy loss in the absorber. The methods differed in their calculation of the opening angle of the muon pair. In the first method the mass, M_V , was calculated using the opening angle obtained from the directions of the tracks emerging from the absorber. This calculation did not depend on whether the dimuon was produced in the target or the dump. In the second method the mass, M_C , was calculated by assuming that the interaction point occurred in the target and exploiting the correlation between the deviation in the direction of the muon and the deviation of the muon position from the unscattered trajectory due to multiple scattering. We calculated the opening angle θ_{12} as the ratio of the separation of the tracks at a plane midway through the absorber to the distance from the midpoint of the target to this plane. The mass M_C is

$$M_C^2 = M_\mu^2 (2 + P_1/P_2 + P_2/P_1) + P_1 P_2 (\theta_{12})^2.$$

The error in the mass due to multiple scattering

in the second calculation was half the error of the first calculation. If the dimuon was actually produced in the absorber the second calculation yielded a mass value which was a factor of 2 too small due to the mistaken assumption that the dimuon was produced in the target. The value of M_C was taken to be the dilepton mass.

The difference between these two calculations, $|M_C - M_V|$, was very sensitive to the actual production point. For that reason it was used to determine whether the event came from the target or from the absorber. If $M_C > 1.4$ GeV/ c^2 , then $|M_C - M_V|$ was required to be < 0.48 GeV/ c^2 . If $M_C < 1.4$ GeV/ c^2 , then $|M_C - M_V|$ was required to be < 0.32 GeV/ c^2 . On the basis of our Monte Carlo calculation, if $M_C = 780$ MeV ($M_C = 3095$ MeV) this procedure excluded 19% (4%) of the events which came from the target and included 23% ($< 1\%$) of the events which came from the absorber. In order to reduce the background of events from the absorber further it was required that T have a count if $M_C < 1.4$ GeV/ c^2 .

The yield of dimuons with charge zero and total momentum greater than 75 GeV/ c , subject to the preceding restrictions, is shown in Fig. 2. If an event had three muons, both zero-charge combinations were included. The fraction of events with more than two muons was less than 1%.

The A dependence of the cross section was determined solely from the yield of dimuons from each target, normalized to the relative neutron intensity. In order to arrive at a statistically significant sample of events at a given mass, we have defined four mass regions as follows: $\rho + \omega$, $0.60 < M < 0.90$; C_1 (continuum), $1.10 < M < 1.40$;




FIG. 2. Yield from target of dimuons of momentum > 75 GeV/ c , with events from all targets added. (a) Mass < 1.4 GeV/ c^2 ; (b) mass > 1.4 GeV/ c^2 . Flux in (b) = $2.5 \times$ flux in (a).

C_2 (continuum), $1.40 < M < 2.6$; and J , $2.6 < M < 3.6$. Although the mass spectra are already quite clean, there remains a small background correction in the first two mass regions, which is performed as follows.

The first source of background is events produced in the dump with the T counter on, due to splashback or secondary production. This was evaluated using the events with the T counter off, which provided a nearly pure sample of dimuons produced in the dump. The invariant mass, M_C , was calculated for these events using the incorrect hypothesis that the dimuon originated in the target. The background was calculated by normalizing the T -off spectrum to be equal to the T -on spectrum in the region 0.4 to 0.55 GeV/c^2 where a clear peak due to ρ production in the dump is present. This resulted in a background subtraction of 10% at the $\rho + \omega$ mass and 10% in the C_1 mass region. (No subtraction is made at higher mass since the background is measured to be $< 2\%$ using the T -off spectrum.)

Next, we attempted to evaluate the nonresonant background under the $\rho + \omega$ peak. For this we used only the T -on events, and fitted the mass spectrum to a sum of ρ , ω , and ϕ resonances and a background. The latter consisted of a continuum term which decreased linearly as the mass increased and a term which was due to ρ production in the absorber. The fits were insensitive to the relative admixture of ρ and ω , because of our mass resolution, but did require a ϕ contribution to fit the shoulder in the mass

spectrum near 1 GeV/c^2 . This method gave identical results, within statistical variations, for the $\rho + \omega$ yield as method 1.

A final source of background came from pion decays. The number of events which can be attributed to pion decay was approximately equal to the number of events which have charge 2, if it was assumed that these events were due entirely to pion decay. No correction was made for this background since it was only $\frac{1}{2}\%$ at the ρ mass.

The relative intensity of the neutral beam was determined from the counting rate in the D counter. The relationship between the counting rate in D and the neutron flux was established by comparing the counting rate of the hadron calorimeter, HC , and D during a special low-intensity run. The D rate and the D rate gated by the live time of the electronics were recorded for each pulse. The latter was used to normalize the yield of dimuons.

During the high-intensity running it was necessary to insert a $3.8 \times 3.8 \times 30.5\text{-cm}^3$ core of tungsten into the absorber in front of HC . This reduced the counting rate in D per incident neutron by a factor of 1.66. The relation between the neutron flux and D for this configuration was made by normalizing the yield of $\rho + \omega$ from the high-intensity data to the moderate-intensity data for similar targets.

No correction for electronic dead time was necessary since the gated D counting rate was proportional to the neutron flux during the live time. The yield for $M_C < 1.4 \text{ GeV}/c^2$ was correct-




FIG. 3. (a) A dependence of yields of $\rho + \omega$ and J . Straight lines are least-squares fits to the data. Error bars are statistical only. (b) A dependence of the dipole yield as a function of mass; cross-section per nu-

cleus proportional to $A^{\gamma(m)}$. Vertical bars are statistical errors from least-squares fits. (c) A dependence of the dipole yield in the $\rho + \omega$ mass region as a function of p_{\parallel} and p_{\perp} of the pair.

ed for the 5.5% inefficiency of the T counter. This efficiency was determined from the ratio of the number of J events from the target which had the T counter on to all J which came from the target. In the J mass region the separation of target and dump is unambiguous.

We have fitted the intensity-normalized dilepton yields per nucleus for $p > 75$ GeV/ c with the power law A^γ . We find values of γ of 0.62 ± 0.03 for the $\rho + \omega$ mass region, 0.85 ± 0.05 for C_1 , 0.85 ± 0.04 for C_2 , and 0.93 ± 0.04 for the J region. The yields for the $\rho + \omega$ and J as functions of A are shown in Fig. 3(a). Most of the increase in γ occurs in the mass interval between the ρ and 1.2 GeV/ c^2 as can be seen from Fig. 3(b). The variation in γ for larger masses is consistent with either a constant value of γ near 0.9 or a slow increase from 0.8 toward 1.0. In the $\rho + \omega$ region γ depends on p_\perp as can be seen from Fig. 3(c). This behavior has been observed for other hadrons by Cronin *et al.*⁶ An examination of the J mass region showed that γ did not depend on p_\perp . We note that the dependence of γ on p_\perp is very similar to its dependence on mass.

The dependence of γ on p_\parallel in the $\rho + \omega$ mass region is also shown in Fig. 3(c). γ did not depend on p_\parallel in the J region to within statistics. Finally, we note that the A dependence in the continuum region is three standard deviations away from an A^1 behavior, which might be expected if the only source of dileptons in this mass range were the Drell-Yan process.⁷

It is a pleasure to acknowledge the assistance

of the staff of the Fermi National Accelerator Laboratory, especially the Proton Department and the Computing Department.

*Supported by the U. S. Energy Research and Development Administration.

†Supported by the National Science Foundation.

¹B. Knapp *et al.*, Phys. Rev. Lett. **34**, 1044 (1975); T. O'Halloran, in *Proceedings of the International Symposium on Lepton and Photon Interactions at High Energies, Stanford, California, 1975*, edited by W. T. Kirk (Stanford Linear Accelerator Center, Stanford, Calif., 1975).

²G. J. Blunar *et al.*, Phys. Rev. Lett. **35**, 346 (1975); F. W. Busser *et al.*, Phys. Lett. **56B**, 482 (1975); K. J. Anderson *et al.*, Phys. Rev. Lett. **36**, 237 (1976).

³Y. M. Antipov *et al.*, Phys. Lett. **60B**, 309 (1976). Earlier observations of dileptons at lower energies were made at Brookhaven National Laboratory by J. H. Christenson *et al.*, Phys. Rev. Lett. **25**, 1523 (1970), and Phys. Rev. D **8**, 2016 (1973).

⁴The J was first observed in hadron-hadron collisions by J. J. Aubert *et al.*, Phys. Rev. Lett. **33**, 1404 (1974), and subsequently by the authors of Refs. 1-3, as well as by H. D. Snyder *et al.*, Phys. Rev. Lett. **36**, 1415 (1976). This last experiment has also made observations on the dilepton mass spectrum between 3 and 10 GeV/ c^2 . D. C. Hom *et al.*, Phys. Rev. Lett. **36**, 1236 (1976).

⁵S. D. Drell and D. M. Yan, Phys. Rev. Lett. **25**, 316 (1970).

⁶J. W. Cronin *et al.*, Phys. Rev. D **11**, 3105.

⁷For example, G. R. Farrar, Phys. Lett. **56B**, 185 (1975).

Dimuon Production in the ρ , J , and Continuum Regions

M. Binkley, I. Gaines, J. Peoples, B. Knapp, W. Lee, P. Leung, S. D. Smith, A. Wijangco, J. Knauer, J. Bronstein, R. Coleman, G. Gladding, M. Goodman, M. Gormley, R. Messner, T. O'Halloran, J. Sarracino, and A. Wattenberg
Fermi National Accelerator Laboratory, Batavia, Illinois 60510, and Columbia University, Irvington-on-Hudson, New York 10533,† and University of Hawaii, Honolulu, Hawaii 96822,‡ and University of Illinois, Urbana, Illinois 61801**

(Received 17 May 1976)

We have measured the p_\perp and p_\parallel dependence of J production by high-energy neutrons in comparison with the production of the conventional vector mesons. We also report values for total inclusive cross sections for resonance and continuum dimuons, and give an upper limit for ψ' production.

Although it has been a year and a half since the discovery of $J^{1,2}$ and of the $\psi'(3684)$ particles,³ very little is known about the mechanism for their production by hadrons. The increase of the inclusive cross section for J production by a factor of

100 between ~ 30 and ~ 300 GeV does not have an experimentally verified explanation.⁴ We have made a measurement of the dynamical dependence of this inclusive cross section. At the same time we have measured the same properties of the in-