

and the U. S. Energy Research and Development Administration.

[†]On leave from Ecole Polytechnique, Paris, France.

¹B. Alper *et al.*, Phys. Lett. **47B**, 275 (1973), and, for example, Nucl. Phys. **B100**, 237 (1975).

²J. W. Cronin *et al.*, Phys. Rev. Lett. **31**, 1426 (1973), and Phys. Rev. D **11**, 3105 (1975).

³R. Blankenbecler, S. J. Brodsky, and J. Gunion, Phys. Rev. **D 12**, 3469 (1975).

⁴S. Matsuda, Phys. Rev. D **12**, 1940 (1975); P. V. Landshoff, J. C. Polkinghorne, and D. M. Scott, Phys.

Rev. D **12**, 3738 (1975); M. Teper, Phys. Rev. D **13**, 629 (1976); R. D. Field and R. P. Feynman, California Institute of Technology Report No. CALT-68-565 (to be published).

⁵At the lower values of $p_{\perp}$, protons and antiprotons are not yet highly relativistic. The p and $\bar{p}$ production angles for $p_{\perp} = 0.77, 1.16, 2.31$, and 3.08 GeV are $125^{\circ}, 106^{\circ}, 99^{\circ}$, and 93° , respectively, at 300 GeV.

⁶S. J. Brodsky and G. R. Farrar, Phys. Rev. Lett. **31**, 1153 (1973); V. A. Matveev, R. M. Muradyan, and A. N. Tavkhelidze, Lett. Nuovo Cimento **7**, 719 (1973).

Inclusive Anomalous Muon Production in e^+e^- Annihilation*

G. J. Feldman, F. Bulos, D. Lüke,[†] G. S. Abrams, M. S. Alam, A. M. Boyarski, M. Breidenbach, J. Dorfan, C. E. Friedberg, D. Fryberger, G. Goldhaber, G. Hanson, F. B. Heile, J. A. Jaros, J. A. Kadyk, R. R. Larsen, A. M. Litke, V. Lüth, R. J. Madaras, C. C. Morehouse, H. K. Nguyen,[‡] J. M. Paterson, M. L. Perl, I. Peruzzi,[§] M. Piccolo,[§] F. M. Pierre,^{||} T. P. Pun, P. Rapidis, B. Richter, B. Sadoulet, R. F. Schwitters, W. Tanenbaum, G. H. Trilling, F. Vannucci,[¶] J. S. Whitaker, and J. E. Wiss

Stanford Linear Accelerator Center, Stanford University, Stanford, California 94305, and Lawrence Berkeley Laboratory and Department of Physics, University of California, Berkeley, California 94720

(Received 8 November 1976)

We present measurements of inclusive anomalous muon production in e^+e^- annihilations in three energy ranges. In all three ranges we observe a large anomalous muon production rate in two-prong events which is compatible with the expected decays of pairs of heavy leptons. In the highest energy range there is also appreciable anomalous muon production in multiprong events which, due to its magnitude and momentum dependence, must come in part from a source other than a heavy lepton.

In previous Letters we presented evidence suggesting that a new lepton^{1,2} and charmed mesons³ are produced in e^+e^- annihilations at center-of-mass energies ($E_{c.m.}$) above 4 GeV. By measuring inclusive muon production we can investigate the leptonic and semileptonic decays of these and possibly other new weakly decaying particles.⁴

The data presented here come from the Stanford Linear Accelerator Center-Lawrence Berkeley Laboratory magnetic detector⁵ at SPEAR and, in particular, from a muon detector (the muon tower) which is located above the main detector. A schematic drawing of the muon-detection system is shown in Fig. 1. To reach the first layer of muon-detection spark chambers (level one), a particle must pass through a 24-g/cm²-thick aluminum coil, a 42-g/cm²-thick lead-scintillator sandwich shower counter, and a 160-g/cm²-thick iron magnetic flux return. To reach the second or third layers of spark chambers (level two or three), a particle must, in addition, pass through one or two 222-g/cm²-thick slabs of bar-

ite-loaded concrete. By taking the average angle of incidence into account, the total material before levels two and three is equivalent to 65 and 92 cm of iron, respectively. Thus the minimum average momentum required for a muon to reach level two is 910 MeV/c. The solid angle subtended by level two is 1.1 sr.

Tracks are momentum-analyzed by the cylindrical spark chambers and projected into the muon tower. For the purpose of this Letter, a *candidate* is a particle which, because of its direction and momentum, would be detected by a muon spark chamber at level two or three if it were a muon. A *muon* is a candidate which is detected in every muon spark chamber through which it is projected within 3 standard deviations of the expected multiple Coulomb scattering. *Anomalous muons* are those remaining after background muons from well-known sources have been subtracted.

Two important sources of background muons are hadron penetration and decay. We calculate




FIG. 1. The magnetic detector as seen looking in the direction of the incident beams. Proportional chambers around the beam pipe and trigger counters around the beam pipe and just inside the coil are not shown.

the former by doing Monte Carlo simulations with the high-energy nucleon-meson transport code HETC⁶ and the latter analytically. In these calculations the candidates are assumed to be 80% π 's and 20% K 's, in rough agreement with measurements.⁷

There are two experimental checks to these calculations. First, we can calculate the number

of muons expected in multiprong ψ decays, where we expect no anomalous muon production. From a sample of ψ decays containing 1879 candidates, we calculate that there should be 52.7 muons from hadron penetration or decays and we observe 57 muons. Second, we can calculate the number of candidates at level three which are detected in spark chambers at levels one and two but are not detected at level three. These events are almost entirely due to hadrons which penetrate or decay to level two. From high-energy multiprong data containing 3491 candidates at level three, we calculate that there should be 161 events of this type and we observe 156 events. From these checks and our estimates of possible systematic errors, we consider the penetration and decay calculations to be accurate to 25%.

Other than π 's and K 's, the ψ 's are the only particles whose decays could conceivably contribute a substantial number of muons. We have looked for inclusive ψ production by plotting the invariant-mass distribution of pairs of particles which include a muon. There is no enhancement at the ψ mass, implying that the number of muons from ψ decay is negligible for these data.

The other sources of background muons we have considered are radiative μ -pair production, $e^+e^- \rightarrow \mu^+\mu^-\gamma$ ⁸ and $\mu^+\mu^-\gamma\gamma$,⁹ and the two-photon process $e^+e^- \rightarrow e^+e^-\mu^+\mu^-$.¹⁰ These reactions primarily contribute muons to events in which only two charged particles are detected. To remove most of the muons from these reactions we require two-prong events to be acoplanar with the incident beams by at least 20° and for the square

TABLE I. Anomalous muon production results.

$E_{c.m.}$ range (GeV)	3.9 to 4.3	4.3 to 4.8	5.8 to 7.8
Average $E_{c.m.}$ (GeV)	4.05	4.4	6.9
Integrated luminosity (pb^{-1})	2.44	2.35	16.2
<u>Two-prong events</u>			
Candidates ^a	181	224	902
Muons	24	29	177
Radiative μ pairs	2.3	2.2	17
$ee\mu\mu$ events	1.4	1.8	29
Hadron penetration or decay	5.0	6.4	28
Anomalous muons	15.3 ± 5.1	18.6 ± 5.7	103 ± 18
Anomalous cross section (pb)	194 ± 71	253 ± 86	212 ± 49
Expected heavy lepton contribution (pb) ^b	252 to 57	290 to 197	195 to 218
<u>Three-or-more-prong events</u>			
Candidates	1217	1392	10738
Muons	47	52	571
Hadron penetration or decay	33.2	38.4	325
Anomalous muons	13.8 ± 10.8	13.6 ± 12.0	246 ± 85
Anomalous cross section (pb)	103 ± 82	105 ± 94	248 ± 93
Expected heavy lepton contribution (pb) ^b	51 to 10	58 to 36	47 to 46

^a Events with both prongs identified as electrons are excluded.

^b For a mass range from 1.6 to 2.0 GeV/c^2 . See text for other parameters.




FIG. 2. Differential cross section for anomalous muon production vs momentum for (a) two-prong events and (b) multiprongs in the $E_{c.m.}$ range from 5.8 to 7.8 GeV. The solid curve represents the expected cross section from the decays of heavy leptons with parameters as specified in the text.

of the missing mass (m_x^2) recoiling against the two observed prongs to be greater than $1.5(\text{GeV}/c^2)^2$. As a check on our calculation of radiative μ -pair production and our ability to measure muon production rates, we have also examined two-prong events which are acoplanar by at least 20° , but which have $m_x^2 < 1.5(\text{GeV}/c^2)^2$. Muons found in these events should result almost entirely from radiative μ -pair production. In the $E_{c.m.}$ region from 3.9 to 4.8 GeV, we expect 37.9 events and we observe 41 events; and in the $E_{c.m.}$ region from 5.8 to 7.8 GeV, we expect 55.8 events and we observe 53 events. We consider the calculation of backgrounds from leptonic sources to be accurate to 20%.

In events with three or more observed charged prongs, we eliminated pairs of oppositely charged particles the cosine of whose opening angle was greater than 0.99, because these pairs were most likely from photon conversions in the 0.052 radiation lengths of material in the vicinity of the beam pipe. In addition, all events containing a muon were examined individually and obvious examples of μ -pair production with photon conversions were eliminated.

The results for anomalous muon production are given in Table I. The cross sections given are corrected for all geometrical, detection, and




FIG. 3. (a) Anomalous muon-production cross section and (b) ratio of anomalous muons to candidates vs the number of observed charged prongs in the $E_{c.m.}$ range from 5.8 to 7.8 GeV. Note that the two-prong cross section is not corrected for a coplanarity cut and is thus artificially suppressed relative to multiprongs cross sections. In calculating the two-prong μ fraction, the number of candidates has been corrected to eliminate leptonic reactions.

triggering efficiencies, but not for the average minimum momentum required to be detected in the tower (910 MeV/c) or, for two prongs, the coplanarity and missing-mass cuts.

In the two-prong case, there is a large anomalous muon signal in all three energy regions.¹¹ Theoretically, one expects that about 87% of the time the decay products of a heavy lepton will contain only one charged particle.¹² Thus, muons from the decay of a pair of heavy leptons should appear primarily in two-prong events. The expected two-prong cross sections for a heavy lepton with a $V-A$ decay, a branching fraction to $\mu\nu\bar{\nu}$ of 0.17, and a mass between 1.6 and 2.0 GeV/c^2 are given in Table I.¹³ In all three energy ranges, the measured anomalous muon cross sections in two-prong events are consistent with coming almost entirely from heavy lepton pairs.

The momentum spectrum for the anomalous muons in two-prong events in the highest $E_{c.m.}$ range is shown in Fig. 2(a), along with the spectrum expected from a heavy lepton in the mass range from 1.6 to 2.0 GeV/c^2 and other parameters as given above. The two spectra are in reasonable agreement.

In the multiprongs data, there is a suggestion of

anomalous muon production in the two lower energy ranges, but it is not conclusive. In the highest energy range, the signal is 5 ± 2 times that expected from heavy-lepton pairs with branching fractions as given above. Furthermore, the muon-momentum spectrum from heavy-lepton pairs should be independent of the event multiplicity, whereas the spectrum observed in multiprong events, shown in Fig. 2(b), is much steeper than that observed for two-prong events or expected from heavy-lepton decays. Figure 3 shows the anomalous cross section and the anomalous-muon-to-candidate ratio in the highest $E_{c.m.}$ range. In both cases, the contrast between two prongs and more than two prongs is striking.

In summary, we have observed a large rate of anomalous muon production in two-prong events consistent with what is expected from heavy-lepton decays. In the 5.8- to 7.8-GeV energy range, we have also observed an anomalous muon-production rate in multiprong events which is several times that expected from heavy-lepton decays and which decreases with muon momentum at a much faster rate than expected from such decays. Thus, these muons must come in part from another source, presumably the weak decays of new hadrons.

We are indebted to T. A. Gabriel, K. J. F. Gaemers, G. Grammer, Jr., and P. Lepage for their assistance with the background calculations.

*Work supported by the U. S. Energy Research and Development Administration.

†Fellow of Deutsche Forschungsgemeinschaft.

‡Permanent address: Laboratoire de Physique Nucléaire et Hautes Energies, Université de Paris VI, Paris, France.

§Permanent address: Laboratori Nazionali, Frascati, Rome, Italy.

||Permanent address: Centre d'Etudes Nucléaires de Saclay, France.

¶Permanent address: Institut de Physique Nucléaire, Orsay, France.

¹M. L. Perl *et al.*, Phys. Rev. Lett. **35**, 1489 (1975).

²M. L. Perl *et al.*, Phys. Lett. **63B**, 466 (1976).

³G. Goldhaber *et al.*, Phys. Rev. Lett. **37**, 255 (1976); I. Peruzzi *et al.*, Phys. Rev. Lett. **37**, 569 (1976).

⁴Other experiments which have reported evidence of inclusive anomalous lepton production in e^+e^- annihilations are the following: M. Cavalli-Sforza *et al.*, Phys. Rev. Lett. **36**, 558 (1976); W. Braunschweig *et al.*, Phys. Lett. **63B**, 471 (1976); and J. Burnmester *et al.*, Deutsches Elektron-Synchrotron Report No. 76/50, 1976 (unpublished).

⁵J.-E. Augustin *et al.*, Phys. Rev. Lett. **34**, 233 (1975).

⁶Neutron Physics Division, Oak Ridge National Laboratory Report No. CCC-178 (unpublished).

⁷T. L. Atwood *et al.*, Phys. Rev. Lett. **35**, 704 (1975).

⁸F. A. Berends, K. J. F. Gaemers, and R. Gastmans, Nucl. Phys. **B57**, 381 (1973).

⁹K. J. F. Gaemers and F. B. Heile, unpublished calculation. The calculation adds a second hard photon in the peaking approximation to the exact α^3 calculation of Ref. 8.

¹⁰G. Grammer, Jr., and T. Kinoshita, Nucl. Phys. **B80**, 461 (1974).

¹¹In 59 of the two-prong muon events with $5.8 \leq E_{c.m.} \leq 7.8$ GeV, the second prong can be identified as a μ , e , or hadron, using the criteria of Ref. 1. After correcting for all backgrounds and misidentifications, there are $6.8 \pm 6.3 \mu\mu$ events, $14.2 \pm 4.8 \mu e$ events, and $19.5 \pm 6.5 \mu$ -hadron events. From the data of Ref. 2, we expect $6.2^{+5.1}_{-2.0} \mu e$ events.

¹²Y. S. Tsai, Phys. Rev. D **4**, 2821 (1971). Details of this calculation are given by G. J. Feldman in the Proceedings of the Summer Institute on Particle Physics, Stanford, California, 2-13 August (to be published).

¹³In Ref. 2, if the observed anomalous $e\mu$ events are interpreted as coming from a sequential heavy lepton with a $V-A$ decay, then the branching fraction to $\mu\nu\bar{\nu}$ is $0.17^{+0.06}_{-0.03}$ and the mass lies between 1.6 and 2.0 GeV/ c^2 .