

<sup>4</sup>H. J. Leamy, G. H. Gilmer, and K. A. Jackson, in *Surface Physics of Crystalline Materials*, edited by J. M. Blakely (Academic, New York, 1976).

<sup>5</sup>R. I. Dobrushin, *Theory Probab. Its Appl. (USSR)* **17**, 619 (1972).

<sup>6</sup>H. van Beijeren, *Commun. Math. Phys.* **40**, 1 (1975).

<sup>7</sup>A. M. Lapunov, *Bull. Acad. Sci. St. Petersburg* **13**, 359 (1900).

<sup>8</sup>M. S. Fisz, *Probability Theory and Mathematical Statistics* (Wiley, New York, 1963), 3rd ed., p. 76.

<sup>9</sup>D. L. Hunter and G. A. Baker, Jr., *Phys. Rev. B* **7**, 3346, 3377 (1973).

<sup>10</sup>The exponents recently calculated in a mean-field (MF) theory of roughening satisfy inequalities (6) and (10) as equalities. ( $\theta_{\mu}^{MF} = \frac{1}{2}$ ,  $\theta_2^{MF} = 1$ ,  $\theta_4^{MF} = 2$ .) R. H. Swendsen, *Phys. Rev. B* **15**, 689 (1976).

---

## ERRATA

---

PERHAPS A STABLE DIHYPERON. R. L. Jaffe [*Phys. Rev. Lett.* **38**, 195 (1977)].

The flavor-octet dihyperon with  $Y=1=0$  and  $J^P = 1^+, H^*$ , does not couple to  $\Lambda\Lambda$  or  $\Sigma\Sigma$  because of statistics. It may be seen as a bump in  $N\Xi$  invariant-mass plots or in the missing mass in  $p\bar{p} \rightarrow K^+K^+X$ . (We thank Dr. L. Littenburg for calling this to our attention.)

The masses of the  $J=2$  8 and 27 were inadvertently omitted from Table I; they are 2066 and 2357 MeV, respectively, in the limit  $m_s=0$ .

SEARCH WITH SYNCHROTRON RADIATION FOR SUPERHEAVY ELEMENTS IN GIANT-HALO INCLUSIONS. C. J. Sparks, Jr., S. Raman, H. L. Yakel, R. V. Gentry, and M. O. Krause [*Phys. Rev. Lett.* **38**, 205 (1977)].

On page 206, column 2, the sixteenth line from the top, the sentence should read, "The numerical values used for  $\sigma_i$  in units of  $10^{-21}$  cm<sup>2</sup>/atom are 2.22 for Cd  $K\alpha$ , 3.66 for Cs  $K\alpha$ , 0.24 for Th  $L\gamma_{1,2,3}$ , and 4.0 for  $L\alpha_1$  of element 126 at 37 keV."