

---

**ERRATA**

---

DETERMINATION OF RELAXATION TIME IN BULK SUPERCONDUCTORS. J. R. Leibowitz and M. C. Wilt [Phys. Rev. Lett. 38, 1167 (1977)].

On page 1167, column 2, line 10, there is a factor-of-2 numerical error in  $\tau_{e-ph}$ . It should read "... we obtain  $\tau_{e-ph} = 0.63 \times 10^{-10}$  (sec)." Also, omit the next  $3\frac{1}{2}$  lines, i.e., lines 11-13, and line 14 up to but not including the word "from." Thus, line 14 should now read: "From (N state) ..."

We are grateful to Dr. Henrik Smith for bringing to our attention the factor-of-2 error.

COSMOLOGICAL UPPER BOUND ON HEAVY-NEUTRINO LIFETIMES. Duane A. Dicus, Edward W. Kolb, and Vigdor L. Teplitz [Phys. Rev. Lett. 39, 168 (1977)].

A distinction must be drawn between two (exhaustive) classes of decay modes of  $\nu_H$ : (1)  $\nu_H \rightarrow 3\nu_L$  ( $\nu_L = \nu_e, \bar{\nu}_e, \nu_\mu,$  and  $\bar{\nu}_\mu$  with mass zero), and (2)  $\nu_H \rightarrow \nu_L + \gamma$  and/or  $\nu_H \rightarrow \nu_L +$  charged particles  $\rightarrow n\nu_L + \gamma$ 's. The results of the paper for lifetime bounds and for cosmological missing mass are valid for the first case. For the second case, there cannot be any contribution to missing cosmological energy but more restrictive bounds (on the order of one year) on the  $\nu_H$  lifetime can be derived from the condition that the  $\gamma$ 's thermalize into the current, 2.7°K, blackbody background.