is the first observation of optically pumped magnetic resonance transitions observed in a flame. This opens the possibility of using spin-exchange spectroscopic techniques¹ to study the molecular structure of free radicals in flames or measuring the concentrations of radicals and other flame constituents.

This research was supported by the U. S. Air Force Office of Scientific Research under Grant No. AFOSR-74-2685 with supplementary support by the Joint Services Electronics Program under Contract No. DAAG29-77-C-0019. One of us (S.M.C.) acknowledges receipt of an Alfred P. Sloan Foundation Fellowship (1975-1979).

(a)On leave from the University of Texas at Dallas, P. O. Box 688, Richardson, Tex. 75080.

- (b)On leave from the Ionosphere Research Laboratory, Kyoto University, Kyoto, Japan.
- ¹W. Happer, Rev. Mod. Phys. 44, 216 (1972).
- ²A. G. Gaydon and H. G. Wolfhard, *Flames, Their Structure*, *Radiation and Temperature* (Chapman and Hall, London, 1953).
- ³C. P. Fenimore, *Chemistry in Premixed Flames* (Macmillan, New York, 1964), p. 7.
- ⁴A. A. Westenberg and R. M. Fristrom, in *Proceedings of the Tenth Symposium on Combustion, Camburidge, Massachusetts*, 1964 (The Combustion Institute, Pittsburgh, 1965), p. 473.
- ⁵P. Regnier and J. P. E. Taran, Appl. Phys. Lett. <u>23</u>, 240 (1973).
- ⁶H. P. Hooymayers and C. Th. J. Alkemade, J. Quant. Spectrosc. Radiat. Transfer <u>6</u>, 847 (1966).
- ⁷P. L. Lijnse, J. Quant. Spectrosc. Radiat. Transfer 14, 1195 (1974).
- ⁸See, for example, R. L. DeZafra and W. Kirk, Am. J. Phys. 35, 573 (1967).
- ⁹W. Franzen and A. G. Emslie, Phys. Rev. <u>108</u>, 1453 (1957).
- ¹⁰R. J. McNeal, J. Chem. Phys. 37, 2726 (1962).

ERRATUM

EXACT SOLUTION FOR THE DIFFUSION OF A PARTICLE IN A MEDIUM WITH SITE DIAGONAL AND OFF-DIAGONAL DYNAMIC DISORDER.

A. Madhukar and W. Post [Phys. Rev. Lett. 39, 1424 (1977)].

In line 7 following Eq. (2), $\beta_{m,n>m\pm 1}(t)=0$ should read $\beta_{m,|n-m|>1}(t)=0$.

In line 3 following Eq. (7), $\beta_{nm}(t) = 0$ for all |m| > |n+1| should read $\beta_{nm}(t) = 0$ for all |m-n| > 1.

In line 5 following Eq. (7), capital K and K' should read k and k'.

On page 1426, in the second line above Eq. (14), the exponential factor should read $e^{-\Gamma t}$.