## ERRATA

DIRECT OBSERVATION OF ELASTIC AND IN-ELASTIC PHOTON SCATTERING BY THE GIANT DIPOLE RESONANCE IN <sup>60</sup>Ni. T. J. Bowles, R. J. Holt, H. E. Jackson, R. M. Laszewski, A. M. Nathan, J. R. Specht, and R. Starr [Phys. Rev. Lett. 41, 1095 (1978)].

The following acknowledgment should be added: "A part of this research, including the development and operation of MUSL-II and its experimental areas, was supported by the National Science Foundation."

<sup>3</sup>He AND <sup>4</sup>He FORM FACTORS AND THE DIMEN-SIONAL-SCALING QUARK MODEL. Benson T. Chertok [Phys. Rev. Lett. 41, 1155 (1978)].

On page 1158, the second and third lines below Eq. (6) should read "... (see Eq. 28 of Ref. 4) ...."

The sentence after Eq. (10) should read "... soft-core potentials.<sup>4</sup>"

In the third paragraph from the end of the Letter on page 1158, the references should be, "parton-model analyses.<sup>4,12</sup>"

DISAPPEARING CENTRAL PEAK IN PARAELEC-TRIC POTASSIUM DIHYDROGEN PHOSPHATE (KDP). Eric Courtens [Phys. Rev. Lett. <u>41</u>, 1171 (1978)].

The correct caption of Fig. 2 should read as follows:

FIG. 2. A typical deviation plot. This plot applies to the experimental point indicated by the arrow in Fig. 1. For this spectrum the integrated number of photocounts is  $2 \times 10^5$ . The abscissa covers one-half free-spectral range, with the laser frequency at channel 0, and the Brillouin peak around channel 13 with  $1.2 \times 10^4$  counts. The ordinate at channel *i* is  $\sigma_i = (n_i - N_i) / \sqrt{n_i}$ , where  $n_i$ is the measured and  $N_i$  the calculated photocount numbers. The  $\chi^2$  is  $\sum \sigma_i^2/46 = 0.69$  for this particular fit.

 $K_s$  REGENERATION ON ELECTRONS FROM 30 TO 100 GeV/c: A MEASUREMENT OF THE  $K^0$ CHARGE RADIUS. W. R. Molzon, J. Hoffnagle, J. Roehrig, V. L. Telegdi, B. Winstein, S. H. Aronson, G. J. Bock, D. Hedin, G. B. Thomson, and A. Gsponer [Phys. Rev. Lett. <u>41</u>, 1213 (1978)].

The sentence beginning on line 5, column 1, p. 1213 should be delected and replaced with: " $\overline{K}^{0}$ "s would be scattered with an amplitude  $\overline{f}^{e}$ =  $-f^{e}$  since  $K^{0}$  and  $\overline{K}^{0}$  have conjugate charge distributions." Also, the sentence beginning on the third line, column 2, page 1214 should begin, "About  $2 \times 10^{6} K_{\pi 2}$  candidates...."