

The partial width is then given by<sup>9</sup>

$$\begin{aligned}\Gamma(\eta' \rightarrow \eta\pi\pi) &= 3|M|^2[1 + 0.24\alpha + 0.27\alpha^2] \\ &= 210 \text{ keV.}\end{aligned}\quad (10)$$

We thus find a width enhanced by almost a factor of 70 from the naive current-algebra result. The dominant contribution to the decay rate arises from  $\delta$  while the  $\epsilon$  term yields the correct slope. To summarize our results, the inclusion of  $\delta$  and  $\epsilon$  resonances satisfactorily accounts for the slope parameter and the decay rate for  $\eta' \rightarrow \eta\pi\pi$ . The reason for no deviation from background in the  $\eta\pi$  distribution and the absence of Adler zero are understood from the accidental cancellation of the zero with the  $\delta$  pole term.

The authors would like to acknowledge the hospitality accorded them by the Theory Group at Los Alamos Scientific Laboratory. This work was supported by the U. S. Department of Energy.

<sup>(a)</sup>Permanent address: Institute of Theoretical Science, University of Oregon, Eugene, Ore. 97403.

<sup>(b)</sup>Permanent address: Centre de Physique Théorique, Ecole Polytechnique, 91120 Palaiseau, France.

<sup>1</sup>G. R. Kalbfleisch, Phys. Rev. D **10**, 916 (1974).

<sup>2</sup>Particle Data Group, Phys. Lett. **75B**, (1978).

<sup>3</sup>Riazuddin and S. Oneda, Phys. Rev. Lett. **27**, 549 (1971).

<sup>4</sup>P. Weisz, Riazuddin, and S. Oneida, Phys. Rev. D **5**, 2264 (1972).

<sup>5</sup>A. Ali, Phys. Rev. D **8**, 1441 (1973); N. G. Deshpande and D. Dicus, Phys. Rev. D **10**, 3048 (1974). See also the review by H. Pagels, Phys. Rep. **16C**, 219 (1975).

<sup>6</sup>The solution  $B = -A/m_\pi^2 = C/2$ ,  $D = E = 0$  yields the Schwinger model; see J. Schwinger, Phys. Rev. **167**, 1432 (1968). This model predicts a large slope  $\alpha \approx -0.44$ . With the value of  $A$  given in Eq. (3) and with  $C$  arbitrary, a large slope is required to increase the theoretical estimate, as discussed in Refs. 3 and 4.

<sup>7</sup>The effect of scalar mesons  $\delta$  and  $\epsilon$  on the  $\eta'$  decay was first considered in an effective-Lagrangian framework by J. Schechter and Y. Ueda, Phys. Rev. D **3**, 2874 (1971), and **9**, 987 (1973). The main difference between that work and the present approach involves the use of derivative couplings and the numerical estimates of the coupling constants.

<sup>8</sup>An attempt to resolve the  $\eta' \rightarrow \eta\pi\pi$  puzzle by considering a purely phenomenological amplitude dominated by the  $\delta$  pole was made by C. Singh and J. Pasupathy, Phys. Rev. Lett. **35**, 1193 (1975). Their amplitude does not manifestly satisfy the current-algebra constraints, and consequently predicts enhancement in the  $\eta-\pi$  distribution at large invariant mass, a feature totally absent from the data (see Ref. 1).

<sup>9</sup>H. Osborn and D. J. Wallace, Nucl. Phys. **B20**, 23 (1970).

## Observables for the Analysis of Event Shapes in $e^+e^-$ Annihilation and Other Processes

Geoffrey C. Fox and Stephen Wolfram

California Institute of Technology, Pasadena, California 91125

(Received 18 September 1978)

We present a set of rotationally invariant observables which characterizes the "shapes" of events, and is calculable in quantum-chromodynamics perturbation theory for final states consisting of quarks and gluons ( $G$ ). We include the effects of fragmentation to hadrons in comparing the shapes of events from the processes  $e^+e^- \rightarrow q\bar{q}$ ,  $e^+e^- \rightarrow q\bar{q}G$ , and  $e^+e^- \rightarrow \text{heavy resonance} \rightarrow GGG$ , and from heavy-quark and lepton production. We indicate how our analysis may be extended to deep-elastic lepton-hadron interactions and hadron-hadron collisions involving large transverse momenta.

Experiments<sup>1</sup> have shown that at high center-of-mass energies ( $\sqrt{s}$ ) the final states in  $e^+e^- \rightarrow \text{hadrons}$  usually consist predominantly of two jets of hadrons presumably resulting from the process  $e^+e^- \rightarrow q\bar{q}$ . Quantum chromodynamics (QCD) explains this basic two-jet structure,<sup>2</sup> but predicts that one of the outgoing quarks should sometimes emit a gluon ( $G$ ), tending to lead to three-jet final states.

Previous attempts<sup>3</sup> to discriminate between two- and three-jet events concentrated on finding a "jet axis" by minimization, and then measuring the collimation of particles with respect to it.

Instead, one may use observables which directly characterize the "shape" of each event. Since there is no natural axis defined in the final state of  $e^+e^-$  annihilation, it is convenient to consider rotationally invariant observables. A set of such observables is given by [ $Y_l^m(\Omega)$  are the usual spherical harmonics and  $P_l(\cos\varphi)$  the Legendre polynomials]

$$\begin{aligned}H_l &\equiv \left( \frac{4\pi}{2l+1} \right) \sum_{m=-l}^{+l} \left| \sum_i Y_l^m(\Omega_i) \frac{|\vec{p}_i|}{\sqrt{s}} \right|^2 \\ &= \sum_{i,j} \frac{|\vec{p}_i||\vec{p}_j|}{s} P_l(\cos\varphi_{ij}),\end{aligned}\quad (1)$$

where the indices  $i$  and  $j$  run over the hadrons which are produced in the event, and  $\varphi_{ij}$  is the angle between particles  $i$  and  $j$ . When the first for the  $H_i$  is used, one must choose a particular set of axes to evaluate the angles ( $\Omega_i$ ) of their momenta, but the values of the  $H_i$  deduced will be independent of the choice. Energy-momentum conservation requires  $H_1 = 0$  and  $H_0 = 1$ . In principle, all the other  $H_i$  carry independent information.<sup>4</sup> In practice, however, one need only consider the lower-order  $H_i$ ; in this paper we concentrate on  $H_2$  and  $H_3$ .

The information contained in the  $H_i$  may also be expressed by the "autocorrelation function"

$$F(\cos\beta) = 2 \int \rho(\hat{\Omega}) \rho(\hat{\Omega}\hat{R}(\alpha, \beta, \gamma)) \\ = \sum_l (2l+1) H_l P_l(\cos\beta), \quad (2)$$

where  $\rho(\hat{\Omega})$  is a continuous distribution of momentum and  $\hat{\Omega}, \hat{R}$  are operators in the rotation group. For particle events, we define the two-detector energy correlation<sup>5</sup>

$$\tilde{F}_2(\sigma_1, \sigma_2) = \frac{16\pi^2}{|\sigma_1||\sigma_2|} \frac{E_1 E_2}{s}, \quad (3)$$

where  $E_i$  are energies incident on detectors covering the regions  $\sigma_i$  of total solid angle  $|\sigma_i|$ . We form the rotationally invariant observable  $F_2$  by averaging  $\tilde{F}_2$  over all possible positions for the detectors, while maintaining their relative orientation. In  $e^+e^-$  annihilation events, this may be achieved (apart from correlations with the beam axis and polarization) by averaging over events. In the limit  $|\sigma_i| \rightarrow 0$ ,  $F_2$  becomes a function solely of the angle  $\beta$  between the two point detectors, and is identical to  $F(\cos\beta)$ .  $F_2$  may clearly be generalized to a correlation between  $n$  detectors ( $F_n$ ). However, unlike the case of the  $H_i$ , there are infrared difficulties when the  $F_n$  are calculated in QCD perturbation theory.<sup>6</sup>

The ability of the  $H_i$  to distinguish between different processes is illustrated in Fig. 1. Final states of the process  $e^+e^- \rightarrow q\bar{q}$  have  $H_i = 1$  for even  $l$  and  $H_i = 0$  for odd  $l$ . In contrast, the process  $e^+e^- \rightarrow q\bar{q}G$  gives events with a wide distribution of  $H_i$  values, corresponding to a range of shapes. For example, the dependence of  $H_2$  on the fractional energies  $x_i (= 2E_i/\sqrt{s})$  of the final quarks and gluons in this case is given by

$$H_2 = 1 - 6(1-x_1)(1-x_2)(1-x_3)/x_1x_2x_3. \quad (4)$$

Each kinematic configuration, labeled by the  $x_i$ , leads to an event of a different shape, and each is characterized by a particular value of  $H_2$ .




FIG. 1. The distributions in  $H_2$  and  $H_3$  for the processes  $e^+e^- \rightarrow q\bar{q}$  (dotted lines),  $e^+e^- \rightarrow q\bar{q}G$  (full lines), and  $e^+e^- \rightarrow$  heavy resonance  $\rightarrow GGG$  (dashed lines). The process  $e^+e^- \rightarrow q\bar{q}G$  alone yields an infinite total cross section, but when added to  $e^+e^- \rightarrow q\bar{q}$  calculated through  $O(g^2)$  the combination of processes [denoted by  $e^+e^- \rightarrow q\bar{q}(G)$ ] gives a finite cross section. We have taken  $\alpha_s = 0.25$  for the  $e^+e^- \rightarrow q\bar{q}G$  distribution.

The  $H_i$  do not discriminate between final states differing by the inclusion of a very low-energy particle or by the replacement of one particle by two collinear particles with the same total momentum. It is believed that these properties are sufficient to ensure that calculations involving the  $H_i$  are infrared finite in QCD perturbation theory.<sup>3,6</sup>

A convenient measure of the event shapes due to different processes is provided by the mean  $H_i$ . For the sum of the process  $e^+e^- \rightarrow q\bar{q}G$  and  $e^+e^- \rightarrow q\bar{q}$  calculated to lowest order in the QCD coupling constant  $\alpha_s = g^2/4\pi$ , we have

$$\langle H_2 \rangle = 1 + (2\alpha_s/3\pi)(33 - 4\pi^2) \simeq 1 - 1.4\alpha_s, \quad (5)$$

so that a center-of-mass energy  $\sqrt{s} = 40$  GeV,  $\langle H_2 \rangle \simeq 0.76$ .

QCD suggests that heavy  $Q\bar{Q}$  vector mesons (such as  $\psi, \Upsilon$ ) should decay to three gluons. Figure 1 shows that the  $H_2$  and  $H_3$  distributions due to this process are very different from those for  $e^+e^- \rightarrow q\bar{q}G$ . The flatter  $H_2$  distribution for the  $GGG$  decay is reflected in a lower  $\langle H_2 \rangle$ :

$$\langle H_2 \rangle \simeq \frac{103\pi^2 - 1008}{16(\pi^2 - 9)} \simeq 0.62. \quad (6)$$

Our results above were obtained by making the idealization that final states consist of free quarks and gluons. In reality, one must consider the "fragmentation" of these quarks and gluons into hadrons, although at sufficiently high energy the values of the  $H_i$  should be the same whether they are calculated from Eq. (1) using the momenta of the actual hadrons in each event, or of their par-

ent quarks and gluons. In order to estimate the shapes of realistic events at finite energy, one must go beyond the realms of present QCD theory and adopt an essentially phenomenological model for the generation of complete hadronic final states by the fragmentation of quarks and gluons. We use the model developed by Field and Feynman,<sup>7</sup> which agrees with available data.<sup>8</sup>

QCD predicts that, away from resonances,  $e^+e^-$  annihilation should be dominated by the processes  $e^+e^- \rightarrow q\bar{q}$  and  $e^+e^- \rightarrow q\bar{q}G$ . The processes  $e^+e^- \rightarrow q\bar{q}G$  can give rise to final states containing either two or three jets of hadrons. Two-jet events occur when some of the quarks and gluons have low energy or are nearly collinear<sup>9</sup> and they cannot be distinguished from  $e^+e^- \rightarrow q\bar{q}$  events by measurements on the hadron final state. Only when  $e^+e^- \rightarrow q\bar{q}G$  and  $e^+e^- \rightarrow q\bar{q}$  [calculated through  $O(g^2)$ ] are added is the jet-production cross section infrared finite. We denote this combination of processes by  $e^+e^- \rightarrow q\bar{q}(G)$ .

In Fig. 2, we present the  $H_2$  distributions for realistic hadronic events resulting from  $e^+e^- \rightarrow q\bar{q}$ ,  $e^+e^- \rightarrow q\bar{q}(G)$ , and  $e^+e^- \rightarrow \xi \rightarrow GGG$  ( $\xi$  is a heavy  $Q\bar{Q}$  resonance). The modifications to the results in Fig. 1 due to the fragmentation of the quarks and

gluons into hadrons are striking. (They also occur for the higher  $H_i$  and for other observables designed to identify three-jet events.<sup>3</sup>) Nevertheless, above  $\sqrt{s} \approx 10$  GeV, the  $H_2$  distributions for the different types of events are clearly distinguished. By  $\sqrt{s} \approx 40$  GeV, the predictions are similar to those obtained in the idealization of free quarks and gluons (Fig. 1).  $H_2$  and  $H_4$  distributions are particularly effective at distinguishing  $e^+e^- \rightarrow q\bar{q}(G)$  and  $e^+e^- \rightarrow \xi \rightarrow GGG$  events, while  $H_3$  distributions are very sensitive to the presence of any pure  $e^+e^- \rightarrow q\bar{q}$  component. The  $H_1$  distributions for realistic events may be made more similar to the idealized ones of Fig. 1 by using only the higher-momentum hadrons in each event for the computation of the  $H_i$ .<sup>10</sup> Even the cut  $|p_i| > 0.5$  GeV is sufficient to effect a great improvement. The  $H_i$  distributions<sup>10</sup> are little affected if only the charged particles in each event are detected. Our predictions are not particularly sensitive to the parameters of the jet development model (which may presumably in any case be determined from single-hadron momentum distributions), but it is still difficult to estimate the uncertainties in our results at energies where the fragmentation of the quarks and gluons has an important effect. Refinement of the jet model as further experimental data become available should allow more accurate predictions to be made.

The  $H_i$  are not specialized to the investigation of two- and three-jet events. They may also be used to identify events of other types. The pair production and weak decay of heavy mesons (containing a heavy quark  $Q$  and a light antiquark  $\bar{q}$ ) and heavy leptons ( $L$ ) should give events containing many hadron jets. For heavy leptons we assume the decay scheme  $L \rightarrow \nu_L u \bar{d}$ , while for heavy quarks (mesons) we consider the three possibilities  $Q \rightarrow q' u \bar{d}$ ,  $Q \rightarrow q' G$ , and  $Q\bar{q}_{\text{spect}} \rightarrow q' \bar{q}''$ . Figure 3 shows our predictions for the  $H_2$  distributions of heavy-quark and lepton production events. We take no account of hadron production by heavy quarks prior to their weak decays, so that our results for heavy-quark pair production should be valid only near threshold.

In addition to the  $H_i$ , one may consider the multipole moments<sup>5</sup>

$$B_i \equiv \sum_i (|\vec{p}_i|/\sqrt{s}) P_i(\cos \alpha_i), \quad (7)$$

where  $\alpha_i$  are the angles made by the particles in the event with the beam axis. A  $q\bar{q}$  final state with angular distribution  $1 + \lambda \cos^2 \alpha_i$  (the naive parton model predicts  $\lambda = 1$ ) gives a broad dis-




FIG. 2. The  $H_2$  distributions predicted for hadronic events resulting from the processes  $e^+e^- \rightarrow q\bar{q}$  (dotted lines),  $e^+e^- \rightarrow q\bar{q}(G)$  (full lines), and  $e^+e^- \rightarrow \text{heavy resonance} \rightarrow GGG$  (dashed lines), at various center-of-mass energies  $\sqrt{s}$ .

tribution in  $B_2$  with mean  $2\lambda/5(\lambda+3)$ , while the process  $e^+e^- \rightarrow q\bar{q}(G)$  gives  $\langle B_2 \rangle \approx 1/10 - 3\alpha_s/10\pi$ , corresponding to  $\lambda \approx 1 - 4\alpha_s/\pi$ , and  $e^+e^- \rightarrow \zeta - GGG$  gives  $\langle B_2 \rangle \approx (72 - 7\pi^2)/80(\pi^2 - 9)$  or  $\lambda \approx (72 - 7\pi^2)/(13\pi^2 - 120) \approx 0.35$ . The  $H_l$ , being rotational invariants, are of course insensitive to correlations with the beam direction. They are, however, far superior in identifying the shape of events and distinguishing competing processes.

The  $H_l$  may also be used to analyze three-jet effects in deep-inelastic lepton-nucleon scattering. Making the idealization of free final quarks and gluons, and treating the nucleon fragments as a single particle, we find that in the virtual photon- (or  $W$ -) nucleon rest frame, two-jet events arising from  $\gamma^*q \rightarrow q$  give  $H_l \approx 1$  for even  $l$  and  $H_l \approx 0$  for odd  $l$ , just as in  $e^+e^-$  annihilation. The three processes<sup>11</sup>  $\gamma^*q \rightarrow q$ ,  $\gamma^*q \rightarrow qG$ , and  $\gamma^*G \rightarrow q\bar{q}$  typically give a  $\langle H_2 \rangle$  which varies smoothly from  $1 - 0.5\alpha_s$  at Bjorken  $x$  around 0.1 to  $1 - 0.9\alpha_s$  at  $x = 0.8$ . The distributions in the  $H_l$  are similar to those in  $e^+e^-$  annihilation. The effects of fragmentation to hadrons are governed by  $s_{\gamma^*N} = Q^2(1/x - 1)$ .

For processes in which a natural plane ( $\Pi$ ) is defined it is convenient to use the two-dimensional analogs of the  $H_l$ :

$$C_l \equiv \left| \sum_i \frac{|\vec{p}_i|_{\text{proj}}}{\sqrt{s}} \exp il\varphi_i \right|^2, \quad (8)$$

where  $\varphi_i$  are the angles of the particles relative to an arbitrary axis in  $\Pi$ , and  $|\vec{p}_i|_{\text{proj}}$  are the




FIG. 3. The  $H_2$  distributions predicted for hadronic events resulting from the production and weak decay of heavy-quark ( $Q$ ) and -lepton ( $L$ ) pairs (dotted lines) at  $\sqrt{s} = 20$  GeV, and in the free-quark and gluon approximation ( $\sqrt{s} \rightarrow \infty$ ). Three mechanisms for heavy-quark decay are considered:  $Q \rightarrow q' u\bar{d}$  (full lines,  $Q \rightarrow q' G$  (dashed lines), and  $Q\bar{q}_{\text{spect}} \rightarrow q' \bar{q}''$  (dot-dashed lines). In the free-quark and gluon approximation the latter two processes give the same  $H_l$  distributions.

magnitudes of their momenta projected onto  $\Pi$ . In deep-inelastic scattering, it is best to take the plane  $\Pi$  to be orthogonal to the  $\gamma^*$  (or  $W^*$ ) direction. Then two-jet events give  $C_l \approx 0$ , while three-jet ones can give nonzero values of  $C_{2l}$ .<sup>12</sup> Typically, in the free-quark approximation,  $\langle C_{2l} \rangle$  is independent of  $l$ , and typically  $\langle C_{2l} \rangle \approx 0.06\alpha_s$  at  $x = 0.1$ , rising to  $0.15\alpha_s$  at  $x = 0.8$ . In hadron-hadron collisions involving high transverse momenta,  $\Pi$  should be chosen as the plane perpendicular to the incoming hadrons. Once again, the distributions in  $C_l/C_0$  distinguish two- and three-jet events. The obvious two-dimensional analog of  $F_2$  [as defined in Eq. (2)] will also be useful.

A detailed discussion of the work summarized here is given in Ref. 6.

This work was supported in part by the U. S. Department of Energy under Contract No. EY 76-C-03-0068. We are grateful to R. D. Field and R. P. Feynman for the use of their jet-development computer program, and to the MATHLAB group of the Massachusetts Institute of Technology Laboratory for Computer Science for the use of MACSYMA.

<sup>1</sup>See, for example, B. H. Wilk and G. Wolf, DESY Report No. 78-23, 1978 (to be published).

<sup>2</sup>G. Sterman and S. Weinberg, Phys. Rev. Lett. **39**, 1436 (1977); G. Sterman, Phys. Rev. D **17**, 2789 (1978).

<sup>3</sup>H. Georgi and M. Machacek, Phys. Rev. Lett. **39**, 1237 (1977); E. Farhi, Phys. Rev. Lett. **39**, 1587 (1977); A. De Rújula, J. Ellis, E. G. Floratos, and M. K. Gaillard, Nucl. Phys. **B138**, 387 (1978); S.-Y. Pi, R. L. Jaffe, and F. E. Low, Phys. Rev. Lett. **41**, 142 (1978). These observables appear to be inconvenient to measure experimentally [S. Brandt and H. D. Dahmen, Seigen Report No. SI-78-8, 1978 (to be published)].

<sup>4</sup>The  $H_l$  may be generalized by taking  $n$  multipole moments of  $\rho(\hat{\Omega})$  and combining them to give scalars under the rotation group using  $3-j$  symbols. The set of all such observables determines  $\rho(\hat{\Omega})$  up to an overall rotation. As we shall describe elsewhere, these observables allow precise tests for planes of particles in events. An obvious application is to  $e^+e^- \rightarrow \zeta - GGG$ .

<sup>5</sup>The mean values of the  $B_l$  and  $\tilde{F}_2$  for  $e^+e^- \rightarrow q\bar{q}(G)$  have also been considered from a rather different point of view in C. L. Basham, L. S. Brown, S. D. Ellis, and S. T. Love, Phys. Rev. D **17**, 2298 (1978), and Phys. Rev. Lett. **41**, 1585 (1978) (this issue).

<sup>6</sup>G. C. Fox and S. Wolfram, California Institute of Technology Report No. CALT-68-678, 1978 (to be published).

<sup>7</sup>R. D. Field and R. P. Feynman, Nucl. Phys. **B136**, 1 (1978).

<sup>8</sup>The model was adjusted so as to agree with observed single-hadron momentum distributions. Limited experimental tests of its predictions for the detailed structure of jets [W. G. Scott, in "Neutrinos—78," edited by Earle C. Fowler (Perdue Univ. Press, to be published)] have proved successful.

<sup>9</sup>The division between configurations of quarks and gluons which give two- and three-jet events is determined by the details of their fragmentation to hadrons. At present the division must be made almost arbitrarily, but our results are not sensitive to the choice (see Ref. 6).

<sup>10</sup>If incomplete final states are considered then only a

fraction of the true energy of the event will be measured, so that it is convenient to use the effective  $H_1/H_0$  rather than  $H_i$  for this case.

<sup>11</sup>All processes of  $O(g^2)$ , including those involving extra initial-state particles (e.g.,  $\gamma^*Gq \rightarrow q$ ), must be added in order to obtain an infrared-finite result. To  $O(g^2)$ , however, only the three-jet parts of  $\gamma^*q \rightarrow qG$  and  $\gamma^*G \rightarrow q\bar{q}$  contribute to  $\langle H_{2l+1} \rangle$  and  $\langle H_{2l} \rangle - 1$ .  $\gamma^*G \rightarrow q\bar{q}$  gives an insignificant contribution.

<sup>12</sup>One may also define two-dimensional analogs of the  $B_l$ . These provide an improved formulation of the tests of QCD proposed by H. Georgi and H. D. Politzer, Phys. Rev. Lett. **40**, 3 (1978).

## Energy Correlations in Electron-Positron Annihilation: Testing Quantum Chromodynamics

C. Louis Basham, Lowell S. Brown, Stephen D. Ellis, and Sherwin T. Love

*Department of Physics, University of Washington, Seattle, Washington 98195*

(Received 21 August 1978)

An experimental measure is presented for a precise test of quantum chromodynamics. This measure involves the asymmetry in the energy-weighted opening angles of the jets of hadrons produced in the process  $e^+e^- \rightarrow$  hadrons at energy  $W$ . It is special for several reasons: It is reliably calculable in asymptotically free perturbation theory; it has rapidly vanishing (order  $1/W^2$ ) corrections due to nonperturbative confinement effects; and it is straightforward to determine experimentally.

Quantum chromodynamics<sup>1</sup> (QCD) is an appealing candidate for the field theory of hadronic interactions, but precise tests are needed to establish its validity. Renormalization-group analyses<sup>2</sup> show that at large momentum transfer the effective coupling becomes small. This suggests that suitably chosen quantities are precisely calculable at high energies, using "asymptotically free perturbation theory." The central assumption here is that those quantities which are free from infrared mass divergences are reliably calculated at high energies in perturbation theory, using the running coupling constant which vanishes at infinite energy.<sup>3</sup> This requires, of course, that energies are well away from (new particle) thresholds (e.g., charm) and that nonperturbative confinement effects are negligible.

The major experimental tests of QCD have been of a semiquantitative nature, involving deeply inelastic lepton scattering from nuclear targets.<sup>4</sup> In general, such analyses suffer from the necessity of introducing arbitrary functions to describe the constituent distributions within the hadronic target. Electron-positron annihilation avoids such ambiguities since there are no hadrons in the initial state, and various features of this an-

nihilation process have been suggested<sup>5</sup> as possible precise tests of QCD. Such features must satisfy several criteria in order to be useful: (1) They must exhibit characteristic properties of QCD, e.g., the presence of both fermions and gauge vector bosons or the vanishing of the running coupling as the energy increases [see Eq. (3)]. (2) They must be reliably calculable. This requires freedom from infrared singularities and insensitivity to nonperturbative effects so as to allow a perturbative analysis. (3) They must be accessible to experiment. This constraint argues against studying quantities which are small effects on much larger backgrounds and quantities which are sensitive to the absence of complete (i.e., all particle) data for each annihilation event.

We have previously suggested<sup>6</sup> the study of the hadronic energy pattern, the "antenna pattern," produced in  $e^+e^-$  annihilation as a possible means to test QCD precisely. The experimental quantity to be measured is the hadronic energy radiated into an element of solid angle  $d\Omega$  (in direction  $\hat{n}$ ) divided by the incident energy per unit area in the  $e^+e^-$  beams. In terms of the partial cross section for the process  $e^+e^- \rightarrow N$  hadrons at