
 ERRATA

STARK EFFECT REVISITED. I. W. Herbst and B. Simon [Phys. Rev. Lett. 41, 67 (1978)].

Equation (2) should read:

$$a_{2n} = \frac{-1}{\pi} \int_0^R \frac{\Gamma(x)}{x^{2n+1}} dx + O(R^{-2n}).$$

HOW MANY NEUTRINOS? Ernest Ma and Jon Okada [Phys. Rev. Lett. 41, 287 (1978)].

On page 287, the last term $\frac{1}{4}[x^2y^2/(1-y^2)]$ in Eq. (3) should be absent, as recently pointed out by K. J. F. Gaemers, R. Gastmans, and F. M. Renard [Phys. Rev. D (to be published)].

SUPERFLUID ^3He IN NARROW CYLINDERS.

C. M. Gould and D. M. Lee [Phys. Rev. Lett. 41, 967 (1978)].

In the middle of the second paragraph on page 967 a line of type was lost. The sentence should have read as follows: "By confining the liquid to cylinders narrower than typical bending lengths ($\approx 10 \mu\text{m}$) we have constrained \hat{l} to lie in a plane,⁴ which necessarily results in at least one singularity somewhere in the plane."

IONIZATION AND ENERGY POOLING IN LASER-EXCITED Na VAPOR. G. H. Bearman and J. J. Leventhal [Phys. Rev. Lett. 41, 1227 (1978)].

Figure 2 was inadvertently printed as both Fig. 2 and Fig. 3. The correct Fig. 3 is given below.




FIG. 3. Na^+ and Na_2^+ ion signals as functions of pressure of various additive gases. Data are normalized to constant $\text{Na}(3p)$ population. The Na^+ data were taken only with N_2 . The arbitrary ordinates for Na_2^+ and Na^+ are unrelated.