

---

 ERRATA
 

---

POLARIZATION OF  $\Lambda$ 's AND  $\bar{\Lambda}$ 's PRODUCED BY 400-GeV PROTONS. K. Heller, P. T. Cox, J. Dworkin, O. E. Overseth, P. Skubic, L. Schachinger, T. Devlin, B. Edelman, R. T. Edwards, G. Bunce, R. Handler, R. March, P. Martin, L. Pondrom, and M. Sheaff. [Phys. Rev. Lett. **41**, 607 (1978)].

On page 610, line 8, in the next to the last paragraph (right-hand side),  $P_{\Sigma^+} = \frac{1}{3}P_{\Lambda}$  should read  $P_{\Sigma^+} = -\frac{1}{3}P_{\Lambda}$ . On page 611, footnote 7,  $P_n = \frac{17}{19}P_{\Lambda}$  should read  $P_n = \frac{20}{21}P_{\Lambda}$ .

MELTING IN "TWO-DIMENSIONAL" SOLID  $^3\text{He}$ . A. Widom, J. R. Owers-Bradley, and M. G. Richards [Phys. Rev. Lett. **43**, 1340 (1979)].

A figure [Fig. 2] crucial to the presentation of data in this Letter was inadvertently omitted and the original version of Fig. 1 was used in its place. The correct Fig. 2 is given below.




FIG. 2. Loci of the two anomalies in NMR data shown in Fig. 1 as  $x$ , the fractional monolayer coverage, is changed. Solid circles give the higher-temperature anomaly  $T_c^{**}$ ; open circles give the lower-temperature anomaly  $T_c^*$ . The dashed line shows the locus of specific-heat peaks (Ref. 12) and the solid line gives the theoretical curve for the Kosterlitz-Thouless model of melting (Ref. 1).