

published).

<sup>13</sup>J. M. Sanchez and D. de Fontaine, Phys. Rev. B 21, 216 (1980).

<sup>14</sup>R. C. Kittler and L. M. Falicov, Phys. Rev. B 18, 2506 (1978), and 19, 291 (1979).

<sup>15</sup>M. Hansen, *Constitution of Binary Alloys* (McGraw-Hill, New York, 1958).

<sup>16</sup>For an introduction to this method, see *Monte Carlo Methods in Statistical Physics*, edited by K. Binder (Springer, Berlin-Heidelberg-New York, 1979).

<sup>17</sup>Previous studies of this model were either restricted to  $c_B=0.5$  ( $H=0$ ),  $N=2048$  sites (Ref. 3) or to concentration  $c_B=0.25$ ,  $N=500$  sites [L. D. Fosdick, Phys. Rev. 116, 565 (1959)]. This lattice size clearly is too small to study the phase transitions, however.

<sup>18</sup>This estimate is in good agreement with Ref. 3, indicating that the inaccuracy due to finite lattice size is small.

<sup>19</sup>A significant change of interaction parameters with lattice spacing has recently been revealed even for

magnetic "model substances" like EuS, cf., U. Köbler and K. Binder, to be published.

<sup>20</sup>P. C. Clapp and S. C. Moss, Phys. Rev. 171, 754 (1968); S. C. Moss and P. C. Clapp, Phys. Rev. 171, 764 (1968); see also H. Chen and J. B. Cohen, Acta Metall. 27, 603 (1979).

<sup>21</sup>K. Binder, unpublished.

<sup>22</sup>W. Selke and M. E. Fisher, to be published.

<sup>23</sup>B. Derrida, Y. Pomeau, G. Toulouse, and J. Vannimenus, J. Phys. (Paris) 40, 67 (1979), and to be published; see also J. Villain, J. Phys. C 10, 1717 (1977); G. Toulouse, Commun. Phys. 2, 115 (1977).

<sup>24</sup>J. M. Cowley, Phys. Rev. 77, 669 (1950).

<sup>25</sup>D. T. Keating and B. E. Warren, J. Appl. Phys. 22, 286 (1951); see also S. C. Moss, J. Appl. Phys. 35, 3547 (1964).

<sup>26</sup>C. Sykes and F. W. Jones, Proc. Roy Soc. London, Ser. A 157, 213 (1938); R. L. Orr, Acta Metall. 8, 489 (1960); R. L. Orr, J. Luciat-Labry, and R. Hultgren, Acta Metall. 8, 431 (1960).

## Roughening Transition in the $^4\text{He}$ Solid-Superfluid Interface

J. E. Avron, L. S. Balfour, C. G. Kuper, J. Landau, S. G. Lipson, and L. S. Schulman

*Department of Physics, Technion-Israel Institute of Technology, Haifa, Israel*

(Received 20 March 1980)

Principal planes of the  $^4\text{He}$  solid-superfluid interface are expected to undergo roughening transitions at temperatures of about 1 K. An experiment is described in which two such transitions were observed in the hcp-superfluid interface: first for the basal plane at 1.08 K and second for an orthogonal face at 0.85 K.

PACS numbers: 68.45.-v, 67.80.-s, 67.40.-w

The roughening transition (RT) is a theoretical concept, known to apply to certain Ising models,<sup>1</sup> and to computer simulations of crystal growth.<sup>2,3</sup> In this Letter we discuss the applicability of the RT to a real physical system: the  $^4\text{He}$  solid-superfluid interface. We report the observation of morphological transitions in  $^4\text{He}$  crystals at 1.08 and 0.85 K which, we argue, are RT's. Parshin and his co-workers<sup>4,5</sup> have argued that the interface in this system is always rough, but their view is contrary to evidence presented here (and by Landau *et al*<sup>6</sup>). Balibar<sup>7</sup> has independently proposed the existence of a RT in the system.

The roughening temperature,  $T_R$ , is characterized by the vanishing of the step energy.<sup>1,2</sup> Above  $T_R$  the interface fluctuates macroscopically, and in the thermodynamic limit, translation invariance is restored.<sup>8</sup> Unfortunately, this aspect of the RT is not expected to be observable in practice. Fluctuations in ordinary macroscopic systems are microscopic; for the two-dimensional interface, the mean square amplitude fluctuation is of order  $\ln N$ , where  $N$  is the number of lattice points in the interface.

The interface is modeled by a two-dimensional (2D) lattice Hamiltonian with discrete, unbounded Ising spin  $Z$ : the vertical coordinate of the interface. A capillary-wave Hamiltonian<sup>9</sup> for the interface is

$$H = \sum_{\vec{i}} \{ 1 + [\nabla Z(\vec{i})]^2 \}^{1/2} [ \sigma_0 \Omega + (\rho_0 \hbar^2 / M^2) (\nabla \psi)^2 ]. \quad (1)$$

Here,  $\psi$  is the superfluid field,  $\rho_0$  its density,  $\nabla$  the discrete gradient,  $\Omega$  the area of the 2D unit cell, and  $\vec{i}$  a lattice bond. The term proportional to  $\sigma_0$  (a coupling constant) represents the energy density of the boundary of the solid, and the term proportional to  $\hbar^2$  is the energy density of the superfluid layer near the interface.<sup>10,11</sup> In the lowest approximation the  $\psi$  field integrates out. Similar conclusions hold for the Villain model<sup>12</sup> and for other models<sup>13</sup> of the interface. The surface tension is the free energy of the Hamiltonian  $H$ .

The specific choice of the interface Hamiltonian is of little importance here since the Kosterlitz-Thouless roughening transition<sup>14</sup> is a general feature of 2D models whose Hamiltonians are invari-

ant under discrete uniform shifts  $Z(\vec{i}) - Z(\vec{i} + \vec{a})$ .<sup>14,15</sup>

The quantum aspects of solid He will be taken into account only phenomenologically;  $\sigma_0$  is taken to be the measured surface tension of  $^4\text{He}$  at  $T = 0$ .<sup>6</sup> The roughening temperature is given by  $T_R = K\sigma_0\Omega$ , where  $K$  is a constant of order unity, whose value depends on the details of the interface model (the choice of Hamiltonian), the lattice structure, etc.<sup>16</sup> Taking  $\sigma_0$  and  $\Omega$  appropriate to the  $^4\text{He}$  interface<sup>6</sup> gives  $T_R \approx 1$  K.

Despite the difficulty of calculating  $T_R$ , we can estimate *relative* values of  $T_R$  on different planes and hence establish a *hierarchical ordering* of plane orientation according to roughening temperature. To accomplish this we identify the  $^4\text{He}$  interface with the interface of a 3D Ising model with Slater-Kirkwood<sup>17</sup> pair interaction and  $\pm$  boundary conditions. This is reasonable only below 1 K, where  $^4\text{He}$ , like the Ising model, has essentially no latent heat.<sup>18,19</sup> To estimate  $T_R$  we use a bound of van Beijern as if it were an equality. Van Beijern proved that for a given crystal orientation  $T_R \geq T_c$ , where  $T_c$  is the critical temperature of the 2D Ising model obtained by considering the restriction of the 3D Ising model to the plane of the interface. This reduces the problem to that of computing  $T_c$  for certain 2D Ising models, which represent sections of the 3D structure and therefore have different lattices and interatomic couplings.

We have estimated  $T_c$  for these 2D models with the Slater-Kirkwood interaction both by mean-field theory<sup>21</sup> and by reduction to soluble models.<sup>22</sup> The results are summarized in Table I, showing that the basal plane has the largest  $T_R$ .

The roughening transition implies<sup>23</sup> that faces which are flat below  $T_R$  will become curved (atom-

ically rough) above it.<sup>24</sup> This can happen with an arbitrarily small change in  $\sigma(\theta, \varphi)$  that smooths a cusp in the Wulff diagram.<sup>24</sup>

We have recently been investigating<sup>6</sup> experimentally the surface tension at the solid-superfluid helium interface. In particular, we found that  $\sigma$  is independent of  $T$  within a single crystallographic phase, and that large equilibrium crystals have apparently more facets at low temperatures than at high temperatures.<sup>25</sup> Evidence for a RT, and even for a hierarchy of such transitions, comes from observing the shape of a small<sup>26</sup> crystal as it was very slowly warmed up. The size of the crystal was comparable with the capillary length, 1.4 mm. A complete sequence of holographic interferograms<sup>27</sup> was photographed,




FIG. 1. Equilibrium holographic interferograms of the  $^4\text{He}$  crystal at constant volume: (a) at 0.6 K; (b) at 0.8 K below the lower transition; (c) at 0.9 K above the lower transition; (d) at 1.06 K before the upper transition; (e) at 1.09 K after the upper transition; (f) at 1.06 K after the reversal of (d)  $\rightarrow$  (e); (g) schematic of the planes in (a) (Ref. 28).

TABLE I. Estimates of the roughening temperature for various crystal planes. The temperature scale was chosen so that the highest  $T_R$  is unity. The mean-field values were obtained by summing the Slater-Kirkwood potential in the given plane. The nearest-neighbors estimate follows from nearest-neighbors soluble models with interaction fitted to the Slater-Kirkwood potential.

Miller indices	$T_R$ , arbitrary units	
	Mean field	Nearest neighbors
(0001)	1	1
(11 $\bar{2}$ 0)	0.5	0.5-1
(10 $\bar{1}$ 0)	0.35	0.2
(10 $\bar{1}$ 1)	0.34	0.16

of which several significant prints are assembled in Fig. 1. At the lowest temperature, 0.6 K, the interferogram (a) shows several plane interfaces  $A, B, C$  (g) to be present. The plane  $A$ , of type  $\langle 11\bar{2}0 \rangle$ , shrank and disappeared by 0.8 K (b). Shortly afterwards the crystallographically equivalent plane  $B$  dramatically disappeared; compare (b) and (c), at 0.9 K. As the temperature continued to rise the facet  $C$  started to shrink and the crystal was observed to undergo a sudden change in external dimensions at 1.8 K. This transition reversed as the temperature was reduced again, and occurred a second time at the same temperature as the heating was recommenced. Apart from the changes in extremal dimensions of the crystal [Figs. 1(d)–1(f)], the transition is less obvious at first glance, because the facet  $C$  is almost parallel to the line of sight. To make the effect clearer, we show in Fig. 2 the area of the face  $C$  as a function of  $T$ , and the transition at 1.08 K is obvious. These observations suggest that the first two transitions in Table I occur at  $1.08 \pm 0.02$  and  $0.85 \pm 0.05$  K.

We found that  $^3\text{He}$  fractions of the order of  $10^{-4}$  or larger enhanced the flat faces, and eliminated curves in the interface.<sup>5</sup> This presents two problems: (a) How does  $^3\text{He}$  stabilize facets? (b) How can such small concentrations of impurities affect the interface?

The answer to (a) is that  $^3\text{He}$  pins the interface. This happens because  $^3\text{He}$  is preferentially expelled from the solid phase by its zero-point energy. A capillary-wave description of the rough surface, following Ref. 4, shows that the time scale for interface fluctuations on an atomic scale is considerably shorter than that for diffusive motion of the  $^3\text{He}$  impurities in the superflu-




FIG. 2. The area of plane  $C$  as a function of  $T$ : open circles, increasing  $T$ ; closed circles, decreasing  $T$ .

id. To understand (b), note<sup>13</sup> that the mean-square amplitude fluctuation of the rough interface  $\langle Z(0)Z(r) \rangle$  is of the order  $(T/2\pi\sigma_0\Omega) \ln(r/a)$ . The interface flattens when the mean distance between the  $^3\text{He}$  impurities,  $r$ , leads to fluctuations small compared to unity. The consequent logarithmic dependence on the concentration explains (b).

In conclusion, the concept of a roughening transition accounts for all the phenomena we have observed in  $^4\text{He}$  hcp-superfluid interfaces.<sup>27</sup>

Two of us (J.E.A.) and (J.L.) thank S. Balibar for stimulating correspondence. One of us (J.L.) is indebted to D. O. Edwards for suggestions and one of us (J.E.A.) thanks E. Witten and R. K. D. Zia for discussion.

This work was supported in part by the U. S.–Israel Binational Science Foundation, Grants No. 1425 and No. 1573.

<sup>1</sup>H. Van Beijern, *Phys. Rev. Lett.* **38**, 992–996 (1977); see also H. Van Beijern, *Comm. Math. Phys.* **40**, 1–6 (1975): Extension to interactions with  $\infty$  range is given in J. Bricmont, J. L. Lebowitz, C. E. Pfister, and E. Olivieri, *Comm. Math. Phys.* **66**, 1–20 (1979); R. L. Dobrushin, *Theory Probab. Its Appl. (USSR)* **17**, 582 (1972).

<sup>2</sup>K. A. Jackson, *J. Cryst. Growth* **24/25**, 130–136 (1974); H. J. Leamy and G. H. Gilmer, *J. Cryst. Growth* **24/25**, 499–592 (1974).

<sup>3</sup>J. D. Weeks, G. H. Gilmer, and H. J. Leamy, *Phys. Rev. Lett.* **31**, 549–551 (1973), give evidence for RT based on low temperature expansion for the sc Ising model.

<sup>4</sup>A. F. Andreev and A. Ya. Parshin, *Zh. Eksp. Teor. Fiz.* **75**, 1511 (1978) [*Sov. Phys. JETP* **48**, 763–766 (1978)].

<sup>5</sup>K. O. Keshishev, A. Ya. Parshin, and A. V. Babkin, *Pis'ma Zh. Eksp. Teor. Fiz.* **30**, 63 (1979) [*JETP Lett.* **30**, 56 (1979)].

<sup>6</sup>J. Landau, S. G. Lipson, L. M. Määtänen, L. S. Balfour, and D. O. Edwards, *Phys. Rev. Lett.* **45**, 31 (1980).

<sup>7</sup>S. Balibar and B. Castaing, to be published.

<sup>8</sup>M. Aizenman, *Phys. Rev. Lett.* **43**, 407 (1979).

<sup>9</sup>D. J. Wallace and R. K. D. Zia, *Phys. Rev. Lett.* **43**, 808–811 (1979), and references therein.

<sup>10</sup>D. R. Nelson and J. M. Kosterlitz, *Phys. Rev. Lett.* **39**, 1201 (1977). We neglect superfluid excitations.

<sup>11</sup>For lattice superfluid models see D. D. Betts, in *Phase Transitions and Critical Phenomena*, edited by C. Domb and M. S. Green (Academic, New York, 1974), Vol. 3.

<sup>12</sup>J. Villain, *J. Phys. (Paris)* **36**, 581 (1975).

<sup>13</sup>Other choices are  $\Sigma(\sigma_0\Omega)|\nabla Z|$ , etc. See, e.g., H. J. F. Knops, *Phys. Rev. Lett.* **39**, 766–769 (1977).

<sup>14</sup>There is extensive literature on the Kosterlitz-Thouless transition, see D. J. Amit, Y. Y. Goldschmidt, and G. Grinstein, *J. Phys. A* **13**, 585-620 (1980); J. Jose, L. P. Kadanoff, S. Kirkpatrick, and D. R. Nelson, *Phys. Rev. B* **16**, 1217 (1977); *Phys. Today* **31**, No. 8, 17 (1978), and references therein. The RT is related to the Kosterlitz-Thouless transition by duality.

<sup>15</sup>S. Elitzur, R. Pearson, and J. Shigemitsu, *Phys. Rev. D* **19**, 3698 (1979).

<sup>16</sup>Rigorous estimates of the Kosterlitz-Thouless transition temperature are given in M. Aizenman and B. Simon, to be published. Accurate, but nonrigorous estimates of  $T_c$  are also quoted there.

<sup>17</sup>J. C. Slater and J. G. Kirkwood, *Phys. Rev.* **37**, 682 (1931).

<sup>18</sup>The latent heat vanishes at  $T=0$  and  $T=0.79$  K [E. R. Grilly, *J. Low Temp. Phys.* **11**, 33 (1973)].

<sup>19</sup> $T_c$  of the Ising model can be pushed to  $\infty$  by imposing a solid-on-solid condition preventing more than a single spin flip in any column. This does not affect our estimates of  $T_R$ .

<sup>20</sup>This means that, e.g., the 3D, sc, nearest-neighbor Ising model has  $T_R \neq 0$  only for the six principal planes. Compare with Ref. 1.

<sup>21</sup>Mean field is an upper bound on  $T_c$ . See R. Griffith, *Commun. Math. Phys.* **6**, 121 (1967). For extensions of

these results see, e.g., B. Simon, to be published, and references given there.

<sup>22</sup>For a given plane, we retain only nearest-neighbor interactions. This underestimates  $T_c$ .  $T_c$  for Ising 2D models are given by I. Syozi, in *Phase Transitions and Critical Phenomena*, edited by C. Domb and M. S. Green (Academic, New York, 1972), Vol. 1.

<sup>23</sup>The large heat conductivity of superfluid  $^4\text{He}$  ensures thermodynamic equilibrium.

<sup>24</sup>C. Herring, in *Structure and Properties of Solid Surfaces*, edited by R. Gomer and G. S. Smith (Univ. of Chicago Press, Chicago, 1955).

<sup>25</sup>If  $\sigma$  was  $T$  dependent no conclusion could be drawn. For a  $T$ -dependent  $\sigma$ , facets may disappear without RT, e.g., by adding a constant to  $\sigma$ .

<sup>26</sup>Gravity gives the vortex excitation a mass inversely proportional to the capillary length. Finite mass destroys RT in the strict sense. For crystals small compared to the capillary length this is irrelevant.

<sup>27</sup>J. Pipman, S. G. Lipson, J. Landau, and N. Bochner, *J. Low Temp. Phys.* **31**, 119 (1978).

<sup>28</sup>The crystallographic details were determined by observing the crystal during growth, when it has a simple pyramidal shape. The faces overlapping at  $D$  in Fig. 1(g) are not exact crystallographic planes, but are in contact with the glass windows.

## Theory of the Surface Tension of Simple Liquids: Application to Liquid Metals

K. K. Mon and D. Stroud

*Department of Physics, The Ohio State University, Columbus, Ohio 43210*

(Received 20 June 1980)

This Letter presents an approximate closed-form solution to the density-functional theory for the surface tension and surface width of simple liquids. The results for both are in excellent agreement with experiments on Ar over wide range of temperature and density. The theory is extended via an ion density-functional formalism to simple metals. Good agreement is obtained for the surface tensions of ten liquid metals and for the surface width of liquid Hg.

PACS numbers: 68.10.Cr, 61.25.Mv, 61.25.Bi

The thermodynamics and structure of liquid-vapor interfaces are subjects of intensive experimental and theoretical interest.<sup>1</sup> Although various theoretical methods have obtained very good agreement with experiment for the surface tension of liquid Ar,<sup>1,2</sup> it is commonly believed that the surface free energies of liquid metals are dominated by electronic effects and that, therefore, theoretical techniques which are successful in treating simple nonmetallic surfaces cannot be used for metals. The work of Evans and Kumaravadeivel,<sup>3</sup> which extends the theory of Lang and Kohn<sup>4</sup> to liquid metals, does lead to agreement with the surface tension in several cases but the predicted widths for the transition profile are small compared with those of most other

fluids.

In this Letter we present a simple and quite successful theory for the surface tension  $\tau$  and width  $l$  of simple liquids, both metallic and non-metallic. It involves a number of approximations which allow one to solve a density-functional theory in closed form and obtain simple expressions for  $\tau$  and  $l$ . The results for both are in excellent agreement with experiment for liquid Ar over wide range of temperature and reproduce the scaling results of Fisk and Widom.<sup>5</sup> The results for the surface tensions of liquid metals are in good agreement with experiment, for all metals studied. The surface widths emerging from the calculations are of the order of one to two atomic layers in all cases and for Hg, the one




FIG. 1. Equilibrium holographic interferograms of the  $^4\text{He}$  crystal at constant volume: (a) at 0.6 K; (b) at 0.8 K below the lower transition; (c) at 0.9 K above the lower transition; (d) at 1.06 K before the upper transition; (e) at 1.09 K after the upper transition; (f) at 1.06 K after the reversal of (d)  $\rightarrow$  (e); (g) schematic of the planes in (a) (Ref. 28).