

PHYSICAL REVIEW LETTERS

VOLUME 46

6 APRIL 1981

NUMBER 14

Critical Behavior of Branched Polymers and the Lee-Yang Edge Singularity

Giorgio Parisi

*Istituto di Fisica della Facoltà di Ingegneria, Università degli Studi di Roma, I-00185 Roma, Italy, and
Istituto Nazionale di Fisica Nucleare, Laboratori Nazionali di Frascati, I-00044 Frascati, Italy*

and

Nicolas Sourlas

*Laboratoire de Physique Théorique de l'Ecole Normale Supérieure, F-75231 Paris 05, France
(Received 29 December 1980)*

The exponents for large branched dilute polymers (which are also connected with the exponents of the lattice animals) are related in D dimensions to the exponents of the Lee-Yang edge singularity of the Ising model in $D - 2$ dimensions. From the exact solution of the Ising model in zero and one dimension, one gets the polymer exponents in two and three dimensions, $\theta(D=2) = 1$ and $\theta(D=3) = \frac{3}{2}$.

PACS numbers: 05.50.+q, 05.70.Jk, 61.40.Km, 75.10.Hk

In this Letter we study the statistics of the dilute limit of branched polymers in a good solvent. This problem belongs to the same universality class as the statistics of lattice animals.¹ The two systems have the same critical exponents which have been computed at first order in $\epsilon = 8 - D$ (D is the space dimension) by Lubensky and Isaacson¹ using a field-theory formalism.

It is well known that in order to study the statistics of polymers, one can represent them as a self-avoiding chain on a regular lattice. Occupied bonds on the lattice represent monomers and connected clusters of occupied bonds represent polymers. In the limit of very dilute branched polymers in a good solvent we can neglect the influence which different polymers exercise on each other and study the statistics of a single cluster.

The statistics of clusters is also studied in the context of the percolation problem. A single connected cluster is usually called a lattice animal in the percolation terminology. The particular properties of the polymers we will study are the number \mathcal{N} of possible configurations of a branched

polymer made from N monomers and the mean end-to-end distance R . The behavior for large N is expected to be the following¹:

$$\mathcal{N} \sim N^{-\theta} \lambda^N, \quad R \sim N^\nu. \quad (1)$$

The critical exponents θ and ν are universal but the constant λ is not.

We show that the critical exponents θ and ν in D dimensions are related to the critical exponent σ which controls the behavior of the magnetization near the Lee-Yang edge singularity² in the Ising model in the presence of an imaginary external field in $d = D - 2$ dimensions

$$\theta(D) = \sigma(d) + 2. \quad (2)$$

The Josephson scaling law (or modern field-theory arguments) connects the singularity of the free energy F with the singularity of the correlation length ξ ($F \sim \xi^{-d}$). We get

$$\nu(D) = [\sigma(d) + 1]/d. \quad (3)$$

In the case of the Ising model it is well known from the Lee-Yang theorem that in the presence

of an external magnetic field $H = H_1 + iH_2$ the correlation functions may be singular only for $H_1 = \text{Re}(H) = 0$. Above the critical temperature T_c there are singularities only for $H_2 = \text{Im}(H) \neq 0$. Let H_0 be the distance from the real axis of the closest singularity, $H = iH_0$. Fisher² has studied this singularity (which he calls the Lee-Yang edge singularity) and has shown that in its neighborhood the magnetization behaves like

$$m \equiv M - M_0 \sim (H - iH_0)^\sigma,$$

where M_0 is the magnetization at iH_0 .

While H_0 is not universal (it depends on the particular lattice, on the temperature, etc.), the exponent σ is universal like the ordinary critical exponents and has been computed by renormalization-group methods^{2,3} in the $\epsilon = d - 6$ expansion.

Because of the shift of the dimensionality appearing in Eqs. (2) and (3) from the exact solution of the Ising model (in the presence of an external field) in $d = 0$ and $d = 1$ dimensions, we get the exact values of θ and ν in the interesting case of three dimensions: $\theta(3) = \frac{3}{2}$, $\nu(3) = \frac{1}{2}$, and $\theta(2) = 1$. These values can be compared with the results of "high-temperature expansions"⁴ (enumeration of the bond animals up to $N = 9$) $\theta(2) = 1.00 \pm 0.01$ and $\theta(3) = 1.55 \pm 0.05$ and the result of Monte Carlo simulations $\nu(3) = 0.45 \pm 0.06$,⁵ or $\nu(3) = 0.53 \pm 0.02$.⁶ From the $\epsilon' = 6 - d$ expansion for σ ,³ we also obtain the first three terms in the $\epsilon = 8 - D$ expansion for θ :

$$\theta = \frac{5}{2} - \frac{\epsilon}{12} - \frac{79}{3888}\epsilon^2 + \left(\frac{\zeta(3)}{81} - \frac{10445}{1259712} \right) \epsilon^3. \quad (4)$$

In order to establish this connection between θ and σ we show that the effective Hamiltonian written by Lubensky and Isaacson for the description of the critical properties of the dilute branched polymers is equivalent to the effective Hamiltonian for the Ising model in a quenched random external field. Then we use the equivalence—in the critical region—of a D -dimensional magnetic system in a quenched random external field to a $d = D - 2$ magnetic system without random external field.

Our method therefore relies entirely on renormalization-group ideas and field theory. We show that the two problems, which look very different, belong to the same universality class. The excellent agreement we find for $\theta(2)$ where the numerical analysis is more accurate may be viewed as another verification of renormalization-group ideas.

For the polymer problem the effective interaction has been shown to be¹

$$\mathcal{H} = \frac{1}{2} \sum_i [(\partial_r \varphi_i)^2 + r\varphi_i^2] - w \sum_i \varphi_i^3 + u \left(\sum_i \varphi_i \right)^2 + H \sum_i \varphi_i, \quad (5)$$

where $i = 1, \dots, n$ and the limit $n \rightarrow 0$ has to be taken at the end. The field φ_i appearing in Eq. (5) is to be identified with the field ψ_{1i} of Ref. 1. The other components ψ_{ji} , $j \neq 1$, do not become critical at the phase transition and their presence can be neglected for the computation of the critical exponents, as was already noticed by the authors of Ref. 1. It was also shown in Ref. 1 that for the Hamiltonian (5) the first infrared singularities in the one-loop diagram level appear for $D = 8$, contrary to naive expectations ($D = 6$). This result holds only for $n = 0$.

As usual, we first shift the field φ_i ($\varphi_i \rightarrow \varphi_i + Q$) in order to eliminate the linear term in (5) and we get the effective Hamiltonian

$$\mathcal{H} = \frac{1}{2} \sum_i [(\partial_r \varphi_i)^2 + m^2 \varphi_i^2 - \lambda \varphi_i^3] + 4uQ^2 \sum_{i,j} \varphi_i \varphi_j + 4uQ \sum_{i,j} \varphi_i^2 \varphi_j. \quad (6)$$

For the computation of the critical exponents one can take account of the leading infrared divergences only, neglecting power corrections. [This is why we omitted the quartic term from Eq. (6).]

The propagator P_{ij} given by Eq. (6) is not diagonal in the fields but has the form

$$P_{ij} = P_{ij}^{(1)} + P_{ij}^{(2)}, \quad P_{ij}^{(1)} = \delta_{ij} / (p^2 + m^2), \\ P_{ij}^{(2)} = \Pi_{ij} / [(p^2 + m^2)(p^2 + m^2 + nuQ^2)], \quad (7)$$

where $\Pi_{ij} = 1$, $\Pi^2 = n\Pi$.

There are thus two propagators $P^{(1)}$ and $P^{(2)}$ and two cubic vertices. Figure 1 represents three Feynman diagrams contributing to the self-energy.




FIG. 1. Feynman diagrams contributing to the self-energy.

gy. Continuous lines represent the $P^{(1)}$ propagator and dashed lines represent the $P^{(2)}$ propagator. In the limit $n \rightarrow 0$, $P^{(2)}$ becomes more singular as $m^2 \rightarrow 0$ and has to be taken a maximum number of times in order to get the most singular diagram. [The diagram 1(b) is more singular than 1(a).] The diagram 1(c) is proportional to n (because $\Pi^2 = n\Pi$) and does not contribute to the limit $n \rightarrow 0$ although more singular (one first takes $n \rightarrow 0$ and then $m^2 \rightarrow 0$). This limits the number of $P^{(2)}$ propagators. An analysis similar to that of Houghton, Reeve, and Wallace⁷ can be carried out. One easily sees that only the $\sum_i \varphi_i^3$ cubic vertex contributes to the most singular diagrams; the vertex $\sum_{i,j} \varphi_i^2 \varphi_j$ combined with $P^{(2)}$ giving additional n factors is suppressed by the $n \rightarrow 0$ limit. If we omit this term, the Hamiltonian (6) also describes the φ^3 theory with a quenched random imaginary external field h (with a Gaussian distribution of width uQ^2) in the replica formalism⁸ (h must be imaginary because of the interaction of polymers being repulsive, $u > 0$).

For real magnetic field it was proven in a perturbative framework⁹ that, at the leading divergence level, the correlation functions of the quenched system in D dimensions are the same as those of a magnetic system without the random external field in dimension $d = D - 2$. As we have previously noted,¹⁰ the leading infrared diagrams correspond to the solution of the stochastic equation

$$-\Delta\varphi + m^2\varphi + w\varphi^2 = ih, \quad \langle h(x)h(y) \rangle = \delta(x - y), \quad (8)$$

where $h(x)$ is a random magnetic field. By redefining φ as $i\varphi$, we get

$$-\Delta\varphi + m^2\varphi + iw\varphi^2 = h. \quad (9)$$

If we write this equation in a field-theory formalism, a hidden supersymmetry appears which allows us to prove the shift of dimensionality in a compact nonperturbative way.¹⁰ Therefore the problem has been reduced to the computation of

the critical exponents of a φ^3 interaction with an imaginary coupling constant. Fisher² has proved that the same interaction appears in the study of the Lee-Yang edge singularity in the presence of an imaginary magnetic field. In this case, one finds that the magnetization has a branch point of the form²

$$m \sim (h - h_0)^\sigma, \quad \sigma = (d - 2 + \eta)/(d + 2 + \eta). \quad (10)$$

Consequently, the free energy has a singularity of the form $(h - h_0)^{\sigma+1}$. The singularity of the free energy of the d -dimensional Ising model is therefore the same as for the D -dimensional polymer problem ($D = d + 2$). From Eq. (1) it follows that the polymer generating function $Z(K) = \sum_n \mathcal{X}(N)K^N$ has a singularity of the form $(K - 1/\lambda)^{\theta-1}$ and this proves Eq. (2).

The exponent σ has been computed up to the third order in the ϵ expansion³ and from this we obtain Eq. (4). The Ising model with an external field is soluble for $d = 0$ (one site) and $d = 1$. One finds that $\sigma(0) = -1$ and $\sigma(1) = -\frac{1}{2}$, from which the exact values for θ and ν in two and three dimensions follow as listed in the introduction. Using the $\epsilon = 8 - D$ expansion and the exact value of θ for $D = 3$, we find the following parametrization for θ :

$$\theta = 2.5 - 0.0833\epsilon \times \frac{1 + 0.5661\epsilon}{1 + 0.3223\epsilon - 0.0406\epsilon^2}, \quad (11)$$

from which we find, using Eq. (3), that $\nu(2) = [d\theta/dD]_{D=2} = 0.61$. If we add the numerical results of series expansion for the Ising model in two and three dimensions,² we get the estimates for θ in four and five dimensions shown in Table I. From ν , the Hausdorff dimension of the animals $d_H = 1/\nu$ follows. Going from eight to three dimensions, the Hausdorff dimension is reduced from 4 to 2.

We would like to thank T. C. Lubensky for useful discussions. One of us (N.S.) would like to thank the Theory Groups at Rome and Frascati, where this work was done, for their warm hospi-

TABLE I. Values of the exponent θ at different dimensions. θ_1 is the value obtained in Ref. 3 from bond animal enumeration. θ_2 is obtained from the high-temperature series evaluation of σ given in Ref. 2 through Eq. (2); and θ_3 is given by Eq. (11).

| D | 2 | 3 | 4 | 5 | 6 | 7 |
|------------|-----------------|-----------------|-----------------|---------------|---------------|---------------|
| θ_1 | 1.00 ± 0.01 | 1.55 ± 0.05 | 1.90 ± 0.07 | 2.2 ± 0.1 | 2.3 ± 0.2 | 2.4 ± 0.2 |
| θ_2 | 1 | 1.5 | 1.85 | 2.10 | ... | ... |
| θ_3 | 1.006 | 1.5 | 1.84 | 2.08 | 2.26 | 2.40 |

tality.

The Laboratoire de Physique Théorique is a Laboratoire Propre du Centre National de Recherche Scientifique, associé à l'Ecole Normale Supérieure et à l'Université de Paris-Sud, 24 Rue Lhomond, F-75231 Paris 05, France.

¹T. C. Lubensky and J. Isaacson, Phys. Rev. Lett. 41, 829 (1978), and Phys. Rev. A 20, 2130 (1979). References to previous work are found in these papers.

²M. E. Fisher, Phys. Rev. Lett. 40, 1610 (1978); D. A. Kurze and M. E. Fisher, Phys. Rev. B 20, 2785 (1979).

³O. F. de Alcantara Bonfim, J. E. Kirkham, and A. J. McKane, J. Phys. A 13, L247 (1980).

⁴D. S. Gaunt, J. Phys. A 13, L97 (1980).

⁵S. Redner, J. Phys. A 12, L239 (1979).

⁶K. Holl, Diploma Thesis, Köln University, Köln-Lindenthal, Germany, 1980 (unpublished).

⁷A. Houghton, J. R. Reeve, and D. J. Wallace, Phys. Rev. B 17, 2956 (1978).

⁸See, for example, T. C. Lubensky, in *Ill-Condensed Matter*, edited by R. Balian, R. Maynard, and G. Toulouse (North-Holland, Amsterdam, 1979).

⁹Y. Imry and S.-k. Ma, Phys. Rev. Lett. 35, 1399 (1975); A. P. Young, J. Phys. C 10, L257 (1977).

¹⁰G. Parisi and N. Sourlas, Phys. Rev. Lett. 43, 744 (1979).