field by surface charges. If the experimental results are interpreted in terms of surface effects, the broadening if the DTA signal towards lower temperatures indicates a lower disordering temperature of the dipoles in the surface region as compared to the dipoles in the interior. The number of dipoles in the surface region may be taken from the area under the broadened shoulder of the endothermal signal. Provided the latent heat of disordering is identical to that of the bulk, the fraction of "surface dipoles" is estimated to be about 30% to 50%. This value coincides with the number of NaNO₂ molecules located at the free surface of a 5-nm crystal. In other words, surface effects are restricted to about one monolayer. There is little evidence for the existence of "dead layers"² at the surface and for particlesize effects on the disordering of the "volume dipoles" as concluded from the unshifted peak temperature.

The interpretation of the experimental data by analogy to superparamagnetism appears unlikely as it requires the "surface" and "volume" dipoles to disorder simultaneously. Simultaneous disorder of both types of dipoles implies an enhanced dipole coupling in the surface region⁴ which appears implausible in view of the reduced coordination number. In addition, it would be hard to understand in terms of this model why the DTA peak is broadened but not shifted.

The financial support by the Deutsche Forschungsgemeinschaft Sonderforschungsbereich 130 is gratefully acknowledged. The authors would like to thank Dr. J. Petermann for many helpful discussions and the suggestion to use PVA for the coprecipitation experiments.

¹C. Jaccard, W. Känzig, and M. Peter, Helv. Phys. Acta <u>26</u>, 521 (1953).

²W. Buchheit, V. Kreibig, D. Müller, and A. Voigt, Z. Phys. B <u>32</u>, 83 (1978).

³D. Y. Parpia, to be published.

⁴K. Binder, in Proceedings of the Fifth International Meeting on Ferroelectricity, University Park, Pennsylvania, 1981 (to be published).

Efficacious Form for Model Pseudopotentials

Leonard Kleinman and D. M. Bylander

Department of Physics, The University of Texas, Austin, Texas 78712 (Received 26 February 1982)

A simple way has been discovered to put model pseudopotentials, $V(\vec{r}) = \sum_{lm} |Y_{lm}\rangle V_l(r)$ × $\langle Y_{lm}|$, into a form which reduces the number of integrals of $V(\vec{r})$ required for an energyband calculation from mn(n+1)/2 to mn for each l in the sum (where n is the number of plane waves used in the expansion and m the number of points in the Brillouin zone at which the calculation is performed). The new form may be chosen to improve the accuracy of the pseudopotential when used in other chemical environments.

PACS numbers: 71.10.+x, 31.15.+q, 71.25.Cx

Pseudopotentials may be divided into three classes, of which one is empirical¹ and the other two are based on first principles. One of the first-principles pseudopotentials is obtained by adding some core functions ψ_c to the wave function ψ to obtain a smooth pseudo wave function,

$$|\varphi\rangle = |\psi\rangle + \sum_{c} \alpha_{c} |\psi_{c}\rangle, \tag{1}$$

and in its most general (non-Hermitian) form is [see Austin, Heine, and Sham² (AHS)]

$$V_{Ps}^{AHS} = V + \sum_{c} |\psi_{c}\rangle \langle F_{c}|, \qquad (2)$$

where $F_c(\mathbf{\tilde{r}})$ is an arbitrary function. By substituting Eq. (1) into the Schrödinger equation, the

relation

$$\alpha_c = \langle F_c | \varphi \rangle / (E - E_c) \tag{3}$$

is obtained. Zwicker³ has shown that in general the AHS pseudopotential is nonvariational, i.e., first-order errors in a trial pseudofunction φ_t cause first-order errors of either sign in the energy E_t . The original Phillips-Kleinman (PK) pseudopotential^{4,5}

$$V_{\rm Ps}^{\rm PK} = V + \sum_{c} (E - E_{c}) |\psi_{c}\rangle \langle\psi_{c}|$$
(4)

has a nonunique φ because $\alpha_c^{PK} = \langle \psi_c | \varphi \rangle$ makes $\varphi' = \varphi + \sum_c \beta_c \psi_c$ a solution of Eq. (1). Zwicker³ was able to show that V_{Ps}^{PK} is variational and

1425

when used with a variational φ_t causes φ to become that unique φ which most closely approximates φ_t . Furthermore, it is obvious from

$$|\psi_t|^{\rm PK} = |\varphi_t\rangle - \sum_c \langle \psi_c |\varphi_t\rangle |\psi_c\rangle \tag{5}$$

that the trial PK wave function is always orthogonal to the core eigenfunctions. The AHS pseudopotential may also be made variational by making it Hermitian by setting $F_c = A_c \psi_c$, with A_c being a constant. In that case it is easy to see that ψ_t is not orthogonal to the ψ_c unless φ_t has sufficient variational freedom in the core region to make it so. Since the whole purpose of using a pseudopotential is to eliminate the need for core variational freedom in φ_t , we conclude that of this class of pseudopotentials, the PK form is the most efficacious. Pseudopotentials in this class have three shortcomings: (1) For valence functions which are orthogonal to all core functions such as oxygen 2p or iron 3d, V_{Ps} reduces to V; about this nothing can be done. (2) V_{PS}^{PK} depends explicitly on the eigenvalue E. This ultimately arises from the nonorthogonality of orthogonalized plane waves and causes no more trouble than the off-diagonal energy dependence of the secular equation obtained from any expansion in nonorthogonal basis functions. (3) Outside the core region the normalized φ is proportional to but not equal to ψ . Thus to perform an accurate self-consistent calculation, one must construct ψ using Eq. (1) or (5) and then renormalize it.

The "model" pseudopotential of Abarenkov and Heine⁶ is based upon the idea that the core is a black box from which the valence wave functions emanate with some logarithmic derivative and that any core pseudopotential which yields that logarithmic derivative is a valid first-principles pseudopotential. Thus the model pseudopotential may be written⁶

$$V_{M} = \begin{cases} V_{val}(\mathbf{\hat{r}}) - \sum_{lm} |Y_{lm}\rangle A_{l}(E) \langle Y_{lm}| \text{ for } r < R_{m}, \\ -Z/r + V_{val}(\mathbf{\hat{r}}) \text{ for } r > R_{m}, \end{cases}$$
(6)

where V_{val} is the Coulomb and exchange potential due to the valence electrons, Z is the valence

$$V_{M} = V_{va1}(\mathbf{\hat{r}}) + \sum_{lm} |Y_{lm}\rangle [V_{l}^{so}(r)\mathbf{\hat{L}} \cdot \mathbf{\hat{S}} + \overline{V}_{l}^{ion}(r)] \langle Y_{lm} |,$$

where

$$V_{l}^{so} = \frac{2}{2l+1} \left[V_{l+1/2}^{HSC} - V_{l-1/2}^{HSC} \right],$$

$$\overline{V}_{l}^{ion} = \frac{1}{2l+1} \left[(l+1) V_{l+1/2}^{HSC} + l V_{l-1/2}^{HSC} \right].$$

charge, and $A_{i}(E)$ is a constant (in space) function of E chosen to make the pseudofunction logarithmic derivative equal that of the true eigenfunction at several atomic eigenvalues. To a high degree of accuracy $A_{i}(E)$ may be linearized:

$$A_{l}(E) = A_{l} + B_{l}E.$$
⁽⁷⁾

This model pseudopotential contains two defects which in principle are irreparable although in practice are not too serious. Because the average potential in an infinite crystal is arbitrary, E in Eq. (6) is not uniquely determined when V_M is used in energy-band calculations. Note that this problem does not arise with V_{Ps}^{PK} because $E - E_c$ is independent of a constant shift in the crystal potential. A second problem is the nonequality of the normalized pseudo wave functions and true wave functions for $r > R_M$. Because in this case the pseudofunction is not obtained from the wave function by the addition of core orbitals, there is no way to construct the wave function from a knowledge of the pseudofunction.

Recently a giant step forward in model pseudopotentials was taken by Hamann, Schlüter, and Chiang⁷ (HSC) who inverted the Schrödinger equation to obtain a model pseudopotential,

$$V_{M}^{HSC} = V_{val}(\mathbf{\hat{r}}) + \sum_{lm} |Y_{lm}\rangle V_{l}^{HSC}(\mathbf{r})\langle Y_{lm}|, \quad (8)$$

where $V_{M}^{HSC}(r) \rightarrow -Z/r$ for $r > r_{c}$ (where r_{c} is a core cutoff radius). The real and pseudo wave functions of V_{M}^{HSC} are identical (not merely proportional) for $r > r_c$. Thus the need to calculate the true wave function is minimized.⁸ More importantly, it can be proven⁷ that as a consequence of the identity of the functions, the logarithmic derivatives and the first energy derivatives of the logarithmic derivatives of the true and pseudo wave functions are identical for $r > r_c$. Hence the energy dependence of V_{M}^{HSC} is second order and in most cases may be ignored. Kleinman⁹ has shown how $V_{l\pm 1/2}^{HSC}(r)$ may be obtained, which when used in the Schrödinger equation yields eigenvalues of the Dirac equation with $j = l \pm \frac{1}{2}$. Thus very accurate self-consistent relativistic energy-band calculations can be made with the following simple model pseudopotential used in the Schrödinger equation⁹:

(11)

(10)

VOLUME 48, NUMBER 20

When attempting to use Eq. (9) for an energy-band calculation in which our basis functions were expanded in plane waves, we were thwarted by the huge number of integrals necessitated by the semilocal (i.e., nonlocal in angular coordinates but local in radial coordinate) part of the potential. Note that the integral of $|Y_{lm}\rangle V(r)\langle Y_{lm}|$ between a pair of plane waves is proportional to

$$\int j_{l}(kr)V(r) j_{l}(k'r)r^{2} dr P_{l}(\cos\theta_{kk'}),$$

where the j_l are spherical Bessel functions and $P_l(\cos\theta_{kk'})$ is the Legendre polynomial of the angle between the wave vectors \vec{k} and $\vec{k'}$. If one has *n* different wave vectors at *m* points in the Brillouin zone, then for each *l* one has mn(n+1)/2 separate integrals to evaluate. In a thin-film calculation *n* can be as large as 1000 and *m* about 20, yielding 10⁷ integrals. Note that this problem does not arise with the fully nonlocal pseudopotentials of the AHS or PK type. The integral of $|\psi_c\rangle\langle\psi_c|$, appearing in Eq. (4), between a pair of plane waves is proportional to

$$\int j_{l}(kr)\psi_{c}(r)r^{2}dr\int j_{l}(k'r)\psi_{c}(r)r^{2}dr P_{l}(\cos\theta_{kk'});$$

this factorization into a product of integrals involving only a single k requires the evaluation of only mn separate integrals for each ψ_c .

In order to circumvent this difficulty we first add and substract a completely arbitrary function $V_L(r)$ from the relativistic form of Eq. (8) [or equivalently from Eq. (6)] to obtain

$$V_{M}^{j=l\pm 1/2} = V_{val}(\mathbf{\hat{r}}) + V_{L}(\mathbf{r}) + \sum_{lm} |Y_{lm}\rangle \,\delta V_{l\pm 1/2} \langle Y_{lm}|, \qquad (12)$$

where $\delta V_{l\pm 1/2} = V_{l\pm 1/2}^{\text{HSC}}(r) - V_L(r)$. With $\Phi_{l\pm 1/2, m}(\mathbf{\hat{r}}) = \varphi_{l\pm 1/2}(r)Y_{lm}(\theta, \varphi)$, we write down a nonlocal pseudo-potential to replace the semilocal term, $V_{\text{SL}}^{j=l\pm 1/2}$, in Eq. (12):

$$V_{\rm NL}^{j=l\pm1/2}(r) = \sum_{lm} \frac{\left| \delta V_{l\pm1/2}(\Phi^0)_{l\pm1/2,m} \right\rangle \langle (\Phi^0)_{l\pm1/2,m} \delta V_{l\pm1/2} |}{\langle (\Phi^0)_{l\pm1/2,m} | \delta V_{l\pm1/2} | \langle (\Phi^0)_{l\pm1/2,m} \rangle} .$$
(13)

The superscript zero indicates that $(\Phi^0)_{l\pm 1/2,m}$ is the psuedo eigenfunction for which $V_{SL}^{j=l\pm 1/2}$ was calculated. Note that

$$V_{\rm NL}^{j=l\pm 1/2} | (\Phi^0)_{l'\pm 1/2,m'} \rangle = |\delta V_{l'\pm 1/2}(\Phi^0)_{l'\pm 1/2,m'} \rangle$$

and that

$$V_{SL}{}^{j=l\pm 1/2} | (\Phi^0)_{l'\pm 1/2, m'} \rangle = | Y_{l'm'} \rangle \delta V_{l'\pm 1/2} | (\varphi^0)_{l'\pm 1/2} \rangle = | \delta V_{l'\pm 1/2} (\Phi^0)_{l'\pm 1/2, m'} \rangle$$

so that 10

$$V_{\rm NL}^{j=l\pm 1/2} | (\Phi^0)_{l'\pm 1/2, m'} \rangle = V_{\rm SL}^{j=l\pm 1/2} | (\Phi^0)_{l'\pm 1/2, m'} \rangle.$$

TABLE I. Dirac-equation eigenvalues for W and two of its ions and the errors obtained by using the semilocal and nonlocal pseudopotentials. There are no errors in the W⁺ pseudopotentials because they were constructed from that configuration. There are no bound 5f states in the $5d^46s^2$ configuration.

	$W(5d^46s^2)$			$W^+(5d^46s^1)$	$W^{++}(5a^46s^0)$		
	Dirac (eV)	Δ (SL) (meV)	Δ (NL) (meV)	Dirac (eV)	Dirac (eV)	Δ (SL) (meV)	Δ (NL) (meV)
$6s_{1/2}$	- 5.890	- 13	- 9	-12.509	-19.637	42	33
$6p_{1/2}$	-1.988	- 5	- 4	-7.710	-14.084	33	26
$6p_{3/2}$	-1.491	-4	- 4	-6.897	-12.962	31	24
$5d_{3/2}$	-5.144	-14	-1	- 12.415	-20.589	33	8
$5d_{5/2}$	-4.421	- 15	-2	-11.644	-19.762	35	11
$5f_{5/2}$	•••	•••	•••	-0.981	-4.404	11	12
$5f_{7/2}$	•••	• • •	• • •	- 0.981	-4.400	10	11

(14)



FIG. 1. The ionic pseudopotentials $V_{l\pm 1/2}^{\text{HSC}}(r)$ and $V_L(r)$ all of which approach -2Z/r for large r.

By this trivial replacement of $V_{\rm SL}$ by the factorized $V_{\rm NL}$ we reduce the number of plane-wave integrals¹¹ required for each l from mn(n+1)/2 to mn. Note also that when the pseudopotential is transported to another atomic configuration or to a crystalline environment $(\Phi^0)_{l'\pm 1/2,m'}$ is replaced by $\Phi_{l'\pm 1/2,m'}$ and Eq. (14) does not hold. Because one has a great deal of freedom in choosing $V_{\rm NL}$ (due to the arbitrariness of V_L), the fact that $V_{\rm NL}{}^{j=l\pm 1/2} |\Phi_{l'\pm 1/2,m'}\rangle$ is not equal to $V_{\rm SL}{}^{j=l\pm 1/2}$ $\times |\Phi_{l'\pm 1/2,m'}\rangle$ may be used to reduce the error incurred in transporting the pseudopotential.

We have constructed relativistic 6s, 6p, 5d, and 5f pseudopotentials for singly ionized tungsten in the configuration¹² $5d^46s^1$. The Dirac eigenvalues listed in Table I were obtained with use of the Wigner interpolation formula for the correlation potential and the relativistic^{13,14} Kohn-Sham exchange potential. The pseudopotentials which are displayed in Fig. 1 were constructed with cutoff functions⁷ of the form $\exp[-(r/r_c)^{3+5}]$ with r_c =1.20, 1.75, 1.85, and 1.85 bohrs for the 5d, 6s, 6p, and 5f potentials. V_L was taken to have the form (in rydbergs)

$$V_{L} = -\frac{2Z}{r} \left\{ 1 - \exp\left[-\left(\frac{r}{\alpha}\right)^{3\cdot 5} \right] + \gamma \gamma \exp\left[-\left(\frac{r}{\beta}\right)^{3\cdot 5} \right] \right\},$$
(15)

with Z = 6, $\alpha = 1.75$, $\beta = 1.8$, and $\gamma = 0.25$. Both $V_{\rm SL}$ and $V_{\rm NL}$ yield the same eigenvalues as the Dirac equation for this configuration. When transported to the $5d^46s^2$ atomic and $5d^46s^0$ doubly ionized configurations the eigenvalues of these pseudopotentials differ from the Dirac eigenvalues by the small amounts listed in Table I. The parameters in V_L have been chosen to make the $V_{\rm NL}$ eigenvalues more nearly correct than the $V_{\rm SL}$. This particular choice of V_L makes $V_{\rm NL}$ worse with respect to $V_{\rm SL}$ when transported to configurations with a different number of d electrons. However, with a few more parameters we were able to construct a V_L which yielded a $V_{\rm NL}$ that was better than $V_{\rm SL}$ for both classes of configurations.

We would like to thank Dr. M. Schlüter for making his relativistic tungsten pseudopotential available to us. This work was supported by the National Science Foundation under Grant No. DMR-80 19518.

 1 M. L. Cohen and T. K. Bergstresser, Phys. Rev. <u>141</u>, 789 (1966).

²B. J. Austin, V. Heine, and L. J. Sham, Phys. Rev. <u>127</u>, 276 (1962).

³R. D. Zwicker, Phys. Rev. B <u>12</u>, 5502 (1975).

⁴J. C. Phillips and L. Kleinman, Phys. Rev. <u>116</u>, 287 (1959).

⁵E. Antoncik, J. Phys. Chem. Solids <u>10</u>, 314 (1959).

⁶I. V. Abarenkov and V. Heine, Philos. Mag. <u>12</u>, 529 (1965).

⁷D. R. Hamann, M. Schlüter, and C. Chiang, Phys. Rev. Lett. <u>43</u>, 1494 (1979).

⁸Although the true and pseudo valence charge densities are not identical in the core region, the integrated charges over the core region are equal. Furthermore, $V_{\rm M}^{\rm HSC}$, like all model pseudopotentials, is designed to yield correct eigenvalues when $V_{\rm val}$ is obtained from the pseudo valence charge density.

⁹L. Kleinman, Phys. Rev. B <u>21</u>, 2630 (1980).

¹⁰Equation (14) insures that if $(\Phi^{0})_{l'\pm l/2}$ is an eigenfunction of the semilocal pseudo Hamiltonian, it is also an eigenfunction of the nonlocal Hamiltonian. However, if $\delta V_{l\pm l/2}$ is very large (and positive) for any *l*, there may exist spurious eigenfunctions of the nonlocal Hamiltonian. These spurious solutions do not occur for the $\delta V_{l\pm l/2}$ with which we calculate in this paper.

¹¹If the integrals are to be calculated in real space, we have reduced some n^3 three-center integrals down to n^2 two-center integrals.

¹²We arbitrarily take the number of $j = l + \frac{1}{2}$ and $j = l - \frac{1}{2}$ electrons to be proportional to l + 1 and l.

¹³A. K. Rajagopal, J. Phys. C 11, L943 (1978).

¹⁴A. H. MacDonald and S. H. Vosko, J. Phys. C <u>12</u>, 2977 (1979).