

Memory Index of First-Passage Time: A Simple Measure of Non-Markovian Character

Peter Hänggi^(a) and Peter Talkner

Institut für Theoretische Physik der Universität Basel, CH-4056 Basel, Switzerland

(Received 17 October 1983)

The theory is developed for the second moment of the first-passage-time variable of statistical flows with memory (non-Markovian dynamics) and applied to a model of free non-Markovian Brownian motion. More importantly, a novel quantity, called memory index σ , is introduced, giving a quantitative characterization of the influence on non-Markovian behavior. Thus, a theoretical or experimental evaluation of σ allows one to assess the quality of widely used approximative (Markovian) Fokker-Planck-type descriptions of the dynamics.

PACS numbers: 05.40.+j, 05.20.-y, 42.50.+q

The concept of the mean of the first-passage-time variable^{1,2} enjoys great popularity among many physicists, chemists, and engineers. Its great practical use for estimating relevant time scales in nonlinear dynamical problems such as they occur in chemical kinetics, decay of metastable states, or decay of unstable states in non-equilibrium systems has been demonstrated many times.¹⁻¹⁰

The main purpose of this paper is the promotion of yet another quantity: the memory index σ [defined below in (20)] of the first-passage-time random variable. A measurement of σ does not prevent major difficulties for experimentalists, yet provides a novel characterization of the quality of an *a priori* Markovian approximation of the coarse-grained dynamics of nonlinear statistical systems. A researcher interested in the theoretical modeling of the dynamics of statistical systems will almost always choose an approximative Markovian theory such as a nonlinear Fokker-Planck modeling.¹⁻¹⁰ It should be borne in mind, however, that such an approach has no divine right. A Markovian modeling merely serves as an *a priori* approximation over a more accurate but generally rather complex non-Markovian modeling.^{8,11,12} It is of course generally understood that "strong" deviations from a Markovian description can be traced back to the coarse graining over a slow variable. In complex systems (e.g., biological systems) it can become rather difficult to collect the complete set of *all* relevant slow variables forming the approximative Markovian dynamics. A quantitative criterion testing the validity of a chosen Markovian approximation appears thus very desirable.

Limiting for the sake of simplicity the following discussion to a one-dimensional stochastic variable $x(t)$, we unavoidably must first develop a small amount of stochastic calculus. Consider

an interval $I = [x_1, x_2]$ out of the state space of the dynamical variable $x(t)$. If initially the random variable $x(t)$ assumes the value $x(t_0=0) = x_0$ in I , the first passage time $\tau(x_0)$ (a random variable) is the time which elapses before $x(t)$ leaves the interval I for the first time. If $F_t(x_0)$ denotes the probability for a random walker, $x(t)$, which started out at time $t_0=0$ at x_0 to be still in interval I without ever having left this interval, one obtains for the probability density $\rho_t(x_0)$ that the first passage time $\tau(x_0)$ lies in the interval $(t, t+dt)$

$$\rho_t(x_0) = -(\partial/\partial t)F_t(x_0), \quad F_{t_0=0}(x_0) = 1. \quad (1)$$

The moments $T_n(x_0)$ of the first-passage-time random variable are defined by

$$T_n(x_0) = \int_0^\infty t^n \rho_t(x_0) dt, \quad n = 1, 2, \dots; \quad (2)$$

$$T_0(x_0) = 1.$$

Now let us first consider a Markovian dynamics; i.e., in terms of a master operator Γ the time evolution of the single-event probability density $p_t(x)$ of the stochastic variable $x(t)$ is in operator notation given by

$$\dot{p}_t = \Gamma p_t. \quad (3)$$

Then, the moments T_n in (2) obey a rather simple equation^{1,2}

$$\bar{\Gamma}^\dagger T_n = -n T_{n-1}, \quad n = 1, 2, \dots; \quad T_0 = \underline{1}. \quad (4)$$

Hereby, $\bar{\Gamma}^\dagger$ denotes the adjoint operator with the kernel $\bar{\Gamma}^\dagger(x, y) = \Gamma(y, x)$ and the bar over $\bar{\Gamma}^\dagger$ denotes the properly adjusted operator so as to prevent backflow of probability into the interval I . In particular, transition rates $\Gamma(y-x) = \Gamma(x, y)$, with x in I and y not in I , are set equal to zero.¹¹ With such boundary conditions included in the definition of $\bar{\Gamma}^\dagger$, T_n is formally given by iterates of the Green's function $[\bar{\Gamma}^\dagger]^{-1}$ (inverse of adjusted

rate matrix^{1,11}):

$$T_n = (-1)^n n! [\bar{\Gamma}^\dagger]^{-n} \underline{1}. \quad (5)$$

A non-Markovian dynamics is characterized by a retarded master operator^{12,13}

$$\dot{p}_t = \int_0^t \underline{K}_{t-s} p_s ds, \quad (6a)$$

where

$$\underline{K}_{t-s} p_s |_{x} = \int K_{t-s}(x, y) p_s(y) dy. \quad (6b)$$

It has been shown previously,¹¹ that the probability $F_t(x_0)$ satisfies the relation

$$\dot{F}_t = \int_0^t \bar{K}_{t-s}^\dagger F_s ds, \quad (7)$$

with the operator \bar{K}_t^\dagger being adjusted as above.

It then follows from (1) and (2) that the mean first passage time T_1 of a non-Markovian process obeys^{14,14}

$$\Omega_0^\dagger T_1 = -\underline{1}, \quad (8a)$$

where Ω_0^\dagger is given by (zeroth moment)

$$\Omega_0^\dagger = \int_0^\infty \bar{K}_s^\dagger ds. \quad (8b)$$

Often, the time integral of the kernel $K_t(x, y)$ is identified with a transition kernel $\Gamma(x, y)$ of a Markovian process; i.e., $\langle x | \int_0^\infty K_t dt | y \rangle \equiv \Gamma(x, y) > 0$, $x \neq y$.¹² Hence, the mean first passage time T_1 of the non-Markovian dynamics, K_t , coincides precisely with T_1 of the commonly used Markovian approximation given by¹²

$$\underline{\Gamma}_{\text{Markov}} \equiv \int_0^\infty \underline{K}_s ds. \quad (9)$$

Clearly this remarkable fact cannot hold true for all moments T_n because processes generated from a common initial probability p_0 (p_0 not the stationary probability \bar{p}) via (6), or via (3) with $\underline{\Gamma}$ given by (9), are markedly different. In particular, let us focus now on the second moment

$$T_2(x_0) = \int_0^\infty t^2 [-\partial F_t(x_0)/\partial t] dt, \quad (10)$$

which after a partial integration¹⁵ equals

$$T_2(x_0) = 2 \int_0^\infty t F_t(x_0) dt. \quad (11)$$

Next we use the trivial identity [see (7)]

$$-\int_0^\infty t \{ \partial F_t / \partial t - \int_0^t \bar{K}_{t-s}^\dagger F_s ds \} dt = 0, \quad (12)$$

which readily yields

$$T_1 + \mathcal{L} \{ \bar{K}_t^\dagger t * F_t; z=0 \} + \mathcal{L} \{ \bar{K}_t^\dagger * F_t t; z=0 \} = 0. \quad (13)$$

The notation $\mathcal{L} \{ f_t; z=0 \}$ denotes the Laplace transform of the function f_t evaluated at $z=0$ and $f * g$ stands for a convolution between f and

g . Introducing the operator Ω_1^\dagger (first moment)

$$\Omega_1^\dagger = \int_0^\infty s \bar{K}_s^\dagger ds, \quad (14)$$

one can recast (13) as

$$\Omega_0^\dagger T_2 = -2 [T_1 + \Omega_1^\dagger T_1]. \quad (15)$$

This relation is a central result of the paper. It determines the second moment of the first-passage-time variable and notably differs from the corresponding Markovian relation in (4) for $n=2$ if Ω_0^\dagger is identified with $\bar{\Gamma}^\dagger$ [see (9)].

Before turning to the interpretation of the result in (15) we apply (15) to the following model of non-Markovian dynamics:

$$\dot{p}_\tau(x) = \gamma^2 \int_0^\tau \left\{ \frac{J_1(\tau - \mu)}{\tau - \mu} \right\} \frac{\partial}{\partial x} \left(\frac{\partial}{\partial x} + x \right) p_\mu(x) d\mu \quad (16)$$

with J_1 being a first-order Bessel function. Equation (16) can be regarded as a model for free Brownian motion presented here in absence of any rigorous derivation. Actually, (16) results as the approximative (Born approximation) generalized Fokker-Planck equation for Rubin's model¹⁶⁻¹⁸ of the Brownian motion of a heavy particle (mass M) in a linear harmonic chain (angular frequency coupling ω_0) of light particles (mass m) with γ^2 being the mass ratio m/M , x the dimensionless velocity, $x = (m/kT)^{1/2} v$, of the heavy particle, and $\tau = \omega_0 t$ the dimensionless time variable.¹⁷ Although (16) is not the exact generalized Fokker-Planck equation for Rubin's model,¹⁷ (16) yields the *exact non-Markovian moments*, $\langle x(\tau) \times |x(0) = x_0 \rangle$ and the *exact stationary velocity correlation function* $\varphi(\tau) = \langle x(\tau)x(0) \rangle$.^{17,19} Observing the integrals $\int_0^\infty [J_1(\tau)/\tau] d\tau = 1$, $\int_0^\infty J_1(\tau) d\tau = 1$, one obtains

$$\Omega_0^\dagger = \Omega_1^\dagger = \gamma^2 (x \partial / \partial x + \partial^2 / \partial x^2) = \underline{\Gamma}_{\text{Fokker-Planck}}^\dagger, \quad (17)$$

or for (15), utilizing (8a),

$$\Omega_0^\dagger T_2 = -2 [T_1 - \underline{1}], \quad (18)$$

which can easily be integrated. If $x = x_1$ is a reflecting boundary, $\partial T_2(x) / \partial x |_{x=x_1} = 0$, and $x_2 > x_1$ absorbing, $T_2(x = x_2) = 0$, one obtains in terms of the stationary probability $\bar{p}(x) = (2\pi)^{-1/2} \exp(-x^2/2)$

$$T_1(x_0) = \int_{x_0}^{x_2} [dy / \bar{p}(y)] \int_{x_1}^y \bar{p}(z) dz / \gamma^2, \quad (19a)$$

$$T_2(x_0) = 2 \int_{x_0}^{x_2} [dy / \bar{p}(y)] \int_{x_1}^y \bar{p}(z) \times \{ T_1(z) - 1 \} dz / \gamma^2, \quad (19b)$$

or

$$[T_2^{NM}(x_0) - T_2^M(x_0)]/T_1^2(x_0) = -2/T_1(x_0). \quad (19c)$$

The superscript NM denotes non-Markovian dynamics given by (6) and (16), and M Markovian dynamics, (9) and (17). Note that $(T_2^{NM} - T_2^M)/T_1^2$ is proportional to $\gamma^2 = m/M$ [compare with (22) below].

The quantity

$$\begin{aligned} \sigma(x_0) &\equiv \frac{T_2^{NM}(x_0) - T_2^M(x_0)}{T_1^2(x_0)} \\ &= \frac{-2}{T_1^2(x_0)} [(\Omega_0^\dagger)^{-1} \Omega_1^\dagger T_1] |_{x_0}, \end{aligned} \quad (20a)$$

or its average over a probability p_0 of starting values x_0

$$\bar{\sigma} = \int_I p_0(x) \sigma(x) dx / \int_I p_0(x) dx, \quad (20b)$$

will be called *memory index of the first-passage-time random variable*. By use of the Markovian approximation (9), or equivalently $\underline{K}_t = 2 \Gamma \delta(t)$, the operator Ω_1^\dagger vanishes, $\Omega_1^\dagger = 0$; i.e., the measure σ vanishes for all x_0 . Hence, the quantity σ can be used as a practical quantitative measure of non-Markovian character and can be evaluated either theoretically or experimentally⁵ measuring the moments $T_2^{NM}(x_0)$, $T_1(x_0)$, and calculating $T_2^M(x_0)$ from the Markov approximation. As mentioned earlier, the quantity σ should be "small" if there exists a clear-cut separation between macroscopic and microscopic time scales; i.e., if both correlations are rapidly decaying.¹² However, the actual magnitude of such "small deviations from Markovian behavior" not only depends on the relaxation-time scale of both correlations but is also a function of the strength of the heat-bath noise. This noise strength can be measured by the value of the macroscopic diffusion coefficient

$$D(x) = \frac{1}{2} \int (y-x)^2 \Gamma_{\text{Markov}}(y,x) dy. \quad (21)$$

In most cases of practical interest, the diffusion of the intensive macrovariable [i.e., $x(t) = \langle x(t) \rangle + O(\Phi^{-1/2})$] scales as the inverse of the system size Φ ,^{8,9,20} e.g., $\Phi = 1/\gamma^2 = M/m$ in our Brownian motion example (16). If we can assume that the random walker starting at x_0 on its motion towards the exit point at x_2 (absorbing boundary) passes a region in state space where the walker must eventually overcome a barrier, or at least passes a region where it moves freely, the first moment $T_1(x_0)$ is growing at least inversely to D . Thus, for vanishing diffusion $D(x)$ ($\Phi \rightarrow \infty$), σ in

(20) approaches zero:

$$\sigma(x) \approx O(1/T_1(x)) \rightarrow 0 \text{ as } D(x) \rightarrow 0. \quad (22)$$

In conclusion, appreciable deviations from the approximative Markovian dynamics are expected to occur predominantly in systems of small size. Most interesting candidates for observing an appreciable non-Markovian character as characterized by a finite memory index are quantum-optical systems^{5-8,10} such as photon statistics in a single-mode laser at threshold or in absorptive optical bistability, magnetic hysteresis, noisy electronic oscillators, transport in Josephson junctions of small size, or ligand migration in biomolecules. The theoretical modeling of the dynamics of those nonlinear systems is usually based on a heuristic nonlinear Fokker-Planck equation^{1-10,21} and most researchers tacitly assume that such an approximative Markovian description will model satisfactorily not only the statics and quantities like T_1 but also the dynamics as, e.g., higher-order correlation functions or higher moments T_n .

The concept put forward in this paper can also be utilized to characterize the degree to which a high-dimensional nonlinear discrete dynamics exhibiting strange attractors can be approximated by a one-dimensional map model.

This work has been supported by the Swiss National Science Foundation.

(a)Permanent address: Department of Physics, Polytechnic Institute of New York, Brooklyn, N.Y. 11201.

¹G. H. Weiss, *Adv. Chem. Phys.* **13**, 1 (1966).

²N. W. Goel and N. Richter-Dyn, *Stochastic Models in Biology* (Academic, New York, 1974).

³E. W. Montroll and K. E. Shuler, *Adv. Chem. Phys.* **1**, 361 (1958).

⁴K. Schulten, Z. Schulten, and A. Szabo, *J. Chem. Phys.* **74**, 4426 (1981).

⁵F. T. Arecchi and A. Politi, *Phys. Rev. Lett.* **45**, 1219 (1980), and *Phys. Lett.* **77A**, 312 (1980).

⁶F. Haake, J. W. Haus, and R. Glauber, *Phys. Rev. A* **23**, 3255 (1981).

⁷P. Hänggi, A. Bulsara, and R. Janda, *Phys. Rev. A* **22**, 671 (1980); M. Delle-Donne, P. H. Richter, and J. Ross, *Z. Phys. B* **42**, 271 (1981), and **45**, 84 (1981).

⁸P. Hänggi and H. Thomas, *Phys. Rep.* **88C**, 209 (1982).

⁹N. G. Van Kampen, *Stochastic Processes in Physics and Chemistry* (North-Holland, Amsterdam, 1981), - Chaps. 6 and 11.

¹⁰H. Risken, in *The Fokker-Planck Equation: Methods of Solution and Applications*, Springer Series in Synergetics, edited by H. Haken (Springer-Verlag, New York,

1983).

¹¹P. Hänggi and P. Talkner, *Z. Phys.* **B45**, 79 (1981).

¹²R. Zwanzig, *Phys. Rev.* **124**, 983 (1961); F. Haake, in *Quantum Statistics in Optics and Solid-State Physics*, Springer Tracts in Modern Physics Vol. 66 (Springer-Verlag, New York, 1973), p. 98, see Eqs. (2b.1)–(2c.5); J. T. Hynes and J. M. Deutch, in *Physical Chemistry: An Advanced Treatise in Eleven Volumes*, edited by D. Henderson (Academic, New York, 1975), Vol. 11B; H. Grabert, in *Projection Operator Techniques in Non-equilibrium Statistical Mechanics*, Springer Tracts in Modern Physics Vol. 95 (Springer-Verlag, New York, 1982), p. 1.

¹³Generally Eq. (6) does contain an inhomogeneous term. If, however, the physical preparation procedure is taken into account explicitly in the definition of the projection operator, this inhomogeneity disappears; see H. Grabert, P. Hänggi, and P. Talkner, *Z. Phys.* **B26**, 398 (1977), and **29**, 273 (1978), and *J. Stat. Phys.* **22**, 537 (1980).

¹⁴For unit-step non-Markovian birth and death processes (including those with two-step transitions) one obtains from Eq. (8a) closed-form expressions for $T_1(x_0=i)$, i in I ; see Eqs. (3.7) and (3.20) in Ref. 11.

¹⁵With $F_{t=\infty}(x_0) = 0$, one utilizes

$$\lim_{t \rightarrow \infty} t^2 F_t(x_0) = \lim_{t \rightarrow \infty} t^2 \int_t^\infty \rho_s(x_0) ds \\ \leq \lim_{t \rightarrow \infty} \int_t^\infty s^2 \rho_s(x_0) ds = 0$$

for bounded $T_2(x_0)$.

¹⁶R. Rubin, *J. Math. Phys. (N.Y.)* **1**, 309 (1960).

¹⁷E. L. Chang, R. M. Mazo, and J. T. Hynes, *Mol. Phys.* **28**, 997 (1974).

¹⁸H. Nakazawa, *Prog. Theor. Phys., Suppl.* **36**, 172 (1966); R. I. Cukier, *Physica (Utrecht)* **61**, 321 (1972); J. M. Deutch and L. Silbey, *Phys. Rev. A* **3**, 2049 (1971).

¹⁹The exact non-Markovian dynamics for Rubin's model is governed by a Gaussian non-Markovian conditional probability which obeys an exact time-convolutionless (but time dependent) master equation; cf. S. A. Adelman, *J. Chem. Phys.* **64**, 124 (1976); R. F. Fox, *J. Math. Phys. (N.Y.)* **18**, 2331 (1977); P. Hänggi, H. Thomas, H. Grabert, and P. Talkner, *J. Stat. Phys.* **18**, 155 (1978). Moreover, the exact generalized Langevin equation is known (Ref. 18) as well as an exact connection [H. Grabert, P. Hänggi, and P. Talkner, *J. Stat. Phys.* **22**, 537 (1980), relations 55 and 56] between the nonlocal retarded kernels in the generalized Fokker-Planck equation and the retarded kernels appearing in the generalized Langevin equation. For the community of statistical physicists it is a rather disturbing fact that the exact time-retarded generalized Fokker-Planck equation for this simple linear model is still not known at present time. In contrast, the exact generalized Fokker-Planck equation is known for nonlinear flows driven by dichotomic noise; cf. P. Hänggi and P. Riseborough, *Phys. Rev. A* **27**, 3379 (1983).

²⁰See also the recent important results by O. J. Eder and T. Lackner, *Phys. Rev. A* **28**, 952 (1983).

²¹See Ref. 8, Sec. 6, and the comprehensive review by Risken (Ref. 10) on the Fokker-Planck equation which gives many examples.