

Experimental Study of Inclusive Muon Spectra from Electron-Positron Collisions in the Energy Region $33 \leq \sqrt{s} \leq 38.54$ GeV

B. Adeva, D. P. Barber, U. Becker, J. Berdugo, G. Berghoff, A. Boehm, J. G. Branson, J. D. Burger, M. Capell, M. Cerrada, C. C. Chang, H. S. Chen, M. Chen, M. L. Chen, M. Y. Chen, Y. S. Chu, R. Clare, E. Deffur, M. Demarteau, P. Duinker, Z. Y. Feng, H. S. Fesefeldt, D. Fong, M. Fukushima, D. Harting, T. Hebbeker, G. Hertel, M. C. Ho, M. M. Ilyas, D. Z. Jiang, W. Krenz, P. Kuijer, Q. Z. Li, D. Luckey, E. J. Luit, C. Mana, M. A. Marquina, G. G. G. Massaro, R. Mount, H. Newman, M. Pohl, F. P. Poschmann, R. R. Rau, J. P. Revol, S. Rodriguez, M. Rohde, J. A. Rubio, H. Rykaczewski, J. Salicio, I. Schulz, K. Sinram, M. Steuer, G. M. Swider, H. W. Tang, D. Teuchert, Samuel C. C. Ting, K. L. Tung, M. Q. Wang, M. White, H. G. Wu, S. X. Wu, B. Wyslouch, B. Zhou, R. Y. Zhu, and Y. C. Zhu

III. Physikalisches Institut, Technische Hochschule, D-5100 Aachen, Federal Republic of Germany, and Deutsches Elektronen Synchrotron DESY, D-2000 Hamburg 52, Federal Republic of Germany, and Laboratory for Nuclear Science, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, and Junta de Energia Nuclear, Madrid, Spain, and Nationaal Instituut voor Kernfysica en Hoge-Energiefysica, 1009-DB Amsterdam, The Netherlands, and Institute of High Energy Physics, Chinese Academy of Science, Beijing, People's Republic of China, and California Institute of Technology, Pasadena, California 91125

(Received 22 April 1983)

The results of a high-statistics study of inclusive muon spectra at PETRA are reported. Improved mass limits have been obtained for heavy quarks, heavy leptons, and charged Higgs particles. It is shown that the fragmentation properties of b quarks and c quarks are different, with the mean fragmentation variables $\langle z_b \rangle = 0.75 \pm 0.03 \pm 0.06$, $\langle z_c \rangle = 0.46 \pm 0.02 \pm 0.05$ and the average semileptonic branching ratio for the B and C hadrons $R(B) = (10.5 \pm 1.5 \pm 1.3)\%$, $R(C) = (11.5 \pm 1.0 \pm 1.7)\%$.

PACS numbers: 13.10.+q, 13.65.+i, 14.80.Dg

In the last twenty years experimental studies of leptons from synchrotrons and colliders have yielded important information in our understanding of particle physics. The discovery of the J ($c\bar{c}$),¹ the Υ ($b\bar{b}$),² and the heavy lepton τ ³ are good examples. It follows naturally that we can use the study of leptons at higher energies to search for new quark states,⁴ new heavy leptons (L^\pm),⁵ and charged Higgs particles (H^\pm).⁶ Further, these studies can provide information on the semileptonic branching ratios and fragmentation functions of known heavy quarks.

New heavy leptons and charged Higgs particles would be pair produced in e^+e^- . The heavy lepton should have a branching ratio into $\mu\bar{\nu}_\mu\nu_L$ of about 14%, and into ν_L +hadrons of about 60%. The charged Higgs particle is expected to decay dominantly into $\tau\bar{\nu}_\tau$ and $s\bar{c}$ (or $b\bar{c}$), with the relative branching ratios of these modes not clearly predicted. Both types of new particles will then give rise to events with an isolated muon roughly opposite to a hadron jet. In the case of the charged Higgs particle, the probability of producing such an event will depend upon the relative branching ratio into $\tau\nu_\tau$.

The production mechanism for hadron events

containing muons,

$$e^+e^- \rightarrow \text{jet}_1(\mu) + \text{jet}_2, \quad (1)$$

can be visualized via the following sequence⁷:

(i) Heavy-quark pairs $Q\bar{Q}$ ($Q=b, c$ quarks) are produced via QED $e^+e^- \rightarrow Q\bar{Q}$. (ii) The fragmentation of $Q \rightarrow M+q$ produces hadron M , which carries the signature of Q , and $q=u, d, s$ quarks which fragment into more hadrons. (iii) The hadron M decays semileptonically, $M \rightarrow \mu + \pi$.

The hadron M can be identified because its decay μ has a distinctive large transverse momentum (with respect to the jet axis), $P_T \sim m_Q/4$. Since the production rates for the heavy quarks are known, a measurement of the rate of muons with transverse momenta in this region determines the semileptonic branching ratio of M . Further, the momentum of the μ along the jet direction provides information on the momentum carried by M and thus on the fragmentation functions of the heavy quarks.

For the present studies we have used data from the MARK-J detector, a 4π calorimeter which measures the directions and energies of e, μ, γ , and jets.⁸ We collected a total of 76 pb^{-1} integrated luminosity in the region $33 \leq \sqrt{s} \leq 38.54$

GeV. From 25 000 hadron events we select inclusive muon events with the criteria described in detail earlier.⁸

The inclusive muon sample contains 806 events with an isolated muon opposite to a hadron jet, dominantly due to τ pair production. By searching for events where the muon and the hadron jet are acoplanar ($>30^\circ$), we unambiguously limit the mass of a new sequential heavy lepton to be greater than 18 GeV. The increase in this limit over that reported previously⁵ is mainly due to a recent increase in the PETRA beam energy. Our limit⁶ on the mass of a charged Higgs particle, assuming that its branching ratio into $\tau\nu_\tau$ is greater than 25%, is slightly improved to $M_{H^\pm} > 14$ GeV.

To obtain mass limits on the production of new quarks and to study the fragmentation processes, we select⁸ the sample of hadron events containing muons in one of the jets, and reduce the effect of hard-gluon emission by collecting only those events with a broad jet oblateness $O_b < 0.3$. After cuts to reduce the background from hadron punchthrough, we are left with 850 events. The remaining punchthrough (13%) and decay (19%) backgrounds are then statistically subtracted from the data. The punchthrough background has been simulated by a Monte Carlo calculation. We reject most of this background by requiring that the muon make a smooth series of chamber hits in two dimensions as it passes through the iron spectrometer. The punchthrough Monte Carlo calculation agrees with the number of rejected events. The efficiency for real muons to pass these cuts has been tested with our data sample on $e^+e^- \rightarrow e^+e^-\mu^+\mu^-$. Background from π and K decay is accurately computed in the Monte Carlo program. We include in our systematic errors (quoted below) 50% uncertainty in the punchthrough background and 20% in the decay background. The background P_T spectrum peaks 300 MeV lower than that from charm and the $P_{||}$ spectrum is peaked at much lower values than that of charm or bottom.

The Monte Carlo calculation includes radiative corrections, detector simulation, and the assumption that B decays via $B \rightarrow C + x$ only. A spectator model⁹ of the B -meson decay was used and the mass of the final charm system was varied between 1.9 and 2.4 GeV, given an average C (D meson) momentum in the B rest frame approximately 1.6 GeV/ c . The rms resolution $\Delta P_T/P_T$ is about 0.5, due to a momentum resolution $\Delta P/P \sim 0.25$ and the error in the reconstruction of the

jet axis.

In the framework of perturbative QCD, we use a simple Feynman-Field⁷ model of fragmentation. Let E_M and E_Q be the energy of M and Q and P_M and P_Q be their momentum along the Q direction. The quantity $z \equiv (E_M + P_M)/(E_Q + P_Q)$ is a measure of the energy which the heavy quark retains after fragmentation. On the basis of the uncertainty principle¹⁰ the fragmentation function for $Q \rightarrow M + q$ can be parametrized¹¹ in terms of h_Q as

$$f_Q(z) = \frac{1}{z} \left(1 - \frac{1}{z} - \frac{h_Q^2}{1-z} \right)^{-2}.$$

Let B and C be the hadronic states with b and c quark signatures, respectively, and let $R(B)$ and $R(C)$ be their average semileptonic-decay branching ratios. Figure 1(a) shows the differential cross section in P_T^2 compared with the pion distribution.¹² The muon P_T^2 distribution is distinctly different from the pion distribution, with a shoulder in the range $2 \lesssim P_T^2 \lesssim 5$ GeV² which is indicative of the decay of a particle of mass around 5 GeV. Figure 1(b) shows the data compared with the Monte Carlo predictions for the total μ rate and its components $B \rightarrow \mu + x$ and $C \rightarrow \mu + x$. We have grouped the small $B \rightarrow C \rightarrow \mu + x$ contribution into the $C \rightarrow \mu + x$ sample. The shoul-




FIG. 1. (a) The P_T^2 distributions of muons normalized to the total hadronic cross section σ_T . The open circles are the cross section with $P > 2$ GeV; the solid points are the cross section extrapolated to all P . They differ only in the first two points with $P_T^2 < 2$ (GeV)². The inclusive π spectrum is also indicated as a dashed curve, scaled by 10^{-2} . (b) Comparison of the data with Monte Carlo predictions, including individual $B \rightarrow \mu$ and $C \rightarrow \mu$ contributions.




FIG. 2. The normalized thrust distribution of the B - and C -enriched samples and all hadronic events. The dashed curve is the Monte Carlo prediction for the B -enriched sample with an additional charge- $\frac{1}{3}$ quark (b') of mass 16 GeV.

der is well explained by B decays, and the Monte Carlo prediction agrees with the data. A P_T cut is used to divide the data into B -enriched and C -enriched samples. The B -enriched sample ($P_T > 1.2$ GeV) has a composition, according to Monte Carlo calculations, of 64% B decays, while the expected composition of the C -enriched sample ($P_T < 1.2$ GeV) is 85% C decays.

Figure 2 shows the thrust distribution for the B - and C -enriched samples and all hadron events. The B -enriched sample exhibits a broader distribution than does the sample of hadrons. No significant difference exists between the C -enriched sample and the hadron sample. Similar to Fig. 1 the thrust distributions of the B -enriched and C -enriched samples can be explained by QCD with known quarks. The P_T and thrust distributions enable us to obtain a limit on the mass of new charge- $\frac{1}{3}$ quarks (b') of $m > 16$ GeV (Fig. 2)⁴ and on charge- $\frac{2}{3}$ quarks (t) of $m > 19$ GeV.⁸ Of course, b' and t quarks with masses only slightly greater than the b quark cannot be ruled out by this method alone. Their existence, however, is inconsistent with the agreement between our measured B semileptonic branching ratio and that measured on the $\Upsilon(4s)$.¹³

Figures 3(a) and 3(b) show the $x = 2P/\sqrt{s}$ distributions for B -enriched and C -enriched samples, respectively, together with the Monte Carlo predictions. The x distribution depends mainly on the fragmentation properties of the quarks. The fact that the average x of the B sample is significantly higher than that of the C sample implies that $\langle z_b \rangle$ is much larger than $\langle z_c \rangle$.

As seen from Figs. 1 and 2 the P_T and thrust distributions separate B and C samples independently. Figure 3 shows that the x (or P) distribu-




FIG. 3. The x distributions for (a) the B -enriched sample, and (b) the C -enriched sample. The Monte Carlo curves are also shown.

tion is a measure of $\langle z_b \rangle$ and $\langle z_c \rangle$. To obtain the values of $R(B)$, $R(C)$, and h_Q we divide the data into an $8 \times 8 \times 8$ three-dimensional grid in the variables of P and P_T of the muon and the thrust. From the maximum-likelihood method we find¹³⁻¹⁵

$$R(C) = (11.5 \pm 1.0 \pm 1.7)\%,$$

$$R(B) = (10.5 \pm 1.5 \pm 1.3)\%,$$

$$|h_c| = 0.8 \pm 0.1 \pm 0.2,$$

$$|h_b| = 0.15 \pm 0.03 \pm 0.05.$$

This yields $\langle z_c \rangle = 0.46 \pm 0.02 \pm 0.05$ and $\langle z_b \rangle = 0.75 \pm 0.03 \pm 0.06$. The ratio $|h_c|/|h_b|$ is $5.3 \pm 1.3 \pm 2.2$, and may be compared with the ratio of the quark masses¹¹ m_b/m_c , which is approximately 3. The first errors¹⁶ are statistical. The second errors are systematic, including uncertainty in the background subtraction, the simulation of detector resolution and acceptance, the variation of mean transverse momenta of the primary particles B and C from 200 to 500 MeV, and the uncertainties in the decay $B \rightarrow \mu + \nu + (C + x)$. Our value of $\langle z_b \rangle/\langle z_c \rangle \approx 1.6$ shows that b and c quarks have different fragmentation properties, with the b quark retaining 75% of its energy in fragmentation.

The quantity $\langle z_Q \rangle$ may be determined without assumption of functional form by dividing the available z region into many intervals and fitting the data bin by bin. With ten equal bins, this yields $\langle z_b \rangle = 0.74 \pm 0.10$, $\langle z_c \rangle = 0.46 \pm 0.05$.

We thank Professor V. Soergel and Professor P. Soeding for their support. This work was supported in part by the Deutsches Bundesministerium für Forschung und Technologie.

¹J. J. Aubert *et al.*, Phys. Rev. Lett. **33**, 1404 (1974); J. E. Augustin *et al.*, Phys. Rev. Lett. **33**, 1406 (1974); J. M. Feller *et al.*, Phys. Rev. Lett. **40**, 274 (1978); R. H. Schindler *et al.*, Phys. Rev. D **24**, 78 (1981).

²S. W. Herb *et al.*, Phys. Rev. Lett. **39**, 252 (1977); W. R. Innes, Phys. Rev. Lett. **39**, 1240 (1977); C. Bebek *et al.*, Phys. Rev. Lett. **46**, 84 (1981); K. Chadwick, *et al.*, Phys. Rev. Lett. **46**, 88 (1981); L. J. Spencer *et al.*, Phys. Rev. Lett. **47**, 771 (1981).

³M. L. Perl *et al.*, Phys. Lett. **63B**, 466 (1976).

⁴M. Chen, in Proceedings of the Banff Summer Institute in Particles and Fields, Banff, Canada, August 1981 (unpublished); J. Buerger, in *Proceedings of the International Symposium on Lepton and Photon Interactions*, edited by W. Pfeil (Physikalisches Institut, Universität Bonn, Bonn, 1981), p. 115; P. Duinker, Rev. Mod. Phys. **54**, 325 (1982).

⁵B. Adeva *et al.*, Phys. Rev. Lett. **48**, 967 (1982); D. P. Barber *et al.*, Phys. Rev. Lett. **45**, 1904 (1980); R. Brandelik *et al.*, Phys. Lett. **99B**, 163 (1981); Ch. Berger *et al.*, Phys. Lett. **99B**, 489 (1981).

⁶A. Ali *et al.*, Nucl. Phys. **B191**, 93 (1981); B. Adeva *et al.*, Phys. Lett. **115B**, 345 (1982); W. Bartel *et al.*, Phys. Lett. **114B**, 211 (1982); H. J. Behrend *et al.*, Phys. Lett. **114B**, 287 (1982); R. Brandelik *et al.*, Phys.

Lett. **122B**, 95 (1983).

⁷R. D. Field and R. P. Feynman, Nucl. Phys. **B136**, 1-76 (1978).

⁸D. P. Barber *et al.*, Phys. Rep. **63**, 337 (1980).

⁹A. Ali *et al.*, Phys. Lett. **93B**, 155 (1980), and Nucl. Phys. **B168**, 409 (1980).

¹⁰M. Suzuki, Phys. Lett. **71B**, 139 (1977); A. Ali, in *Physics and Contemporary Needs* (Plenum, New York, 1982), Vol. 5.

¹¹C. Peterson *et al.*, Phys. Rev. D **27**, 105 (1983).

¹²R. Brandelik *et al.*, Phys. Lett. **113B**, 98 (1982); G. Wolf, DESY Report No. 82-077, 1982 (unpublished).

¹³Our $R(B)$ can be compared with the value $(12.4 \pm 1.7 \pm 3.1)\%$ measured on the resonance Υ''' . K. Chadwick *et al.*, Phys. Rev. D **27**, 473 (1983).

¹⁴The world average branching ratio is $19^{+4}_{-3}\%$ for $D^{\pm} \rightarrow e + x$ and $< 6\%$ for D^0 . M. Roos *et al.* (Particle Data Group), Phys. Lett. **111B**, 1 (1982).

¹⁵For earlier measurements on h_c , see W. Bartel *et al.*, DESY Report No. 80/86, 1980 (unpublished); K. Kleinknecht and B. Renk, Z. Phys. C **17**, 325 (1983); D. Lueke *et al.*, DESY Report No. 82/073, 1982 (unpublished). For preliminary results on h_b , see G. H. Trilling, J. Phys. (Paris), Colloq. **43**, C3-58 (1982); J. M. Yelton *et al.*, Phys. Rev. Lett. **49**, 430 (1982); M. Althoff *et al.*, DESY Report No. 83/010, 1983 (unpublished).

¹⁶These errors include the effects of correlations between the parameters and represent the 68% (one-standard-deviation) confidence interval in each parameter, with all other parameters free to vary. It should be noted that the probability distribution is approximately Gaussian in the h 's, not in the more commonly quoted parameter h^2 .