

---

**ERRATA**


---

**X-Ray Resonance Exchange Scattering.** J. P. HANNON, G. T. TRAMMELL, M. BLUME, and DOON GIBBS [Phys. Rev. Lett. **61**, 1245 (1988)]

The factor  $P$  appearing in Eq. (4) should be  $P = \{n_e(\uparrow) - [\Delta/\Gamma(x-i)]n_h\}$ . Because of the additional frequency factor,  $(x-i)^{-1}$ , the exchange splitting  $\Delta$  and the induced moment  $n_e(\uparrow)$  can be separately determined. This correction was pointed out to us by M. Altarelli.

In Eq. (2),  $Y_{LM}(\hat{r}_i)$  should be  $Y_{LM}(\hat{r}_i)^*$ .

In the first equation on p. 1247,  $C^2(1,1,2;m_l - M, m_s, m_l)$  should be  $C^2(1,1,2;m_l - M, M, m_l)$ , and the factor  $\frac{24}{5}$  in  $|\chi|^2$  should be  $\frac{8}{15}$ .

In the seventh line from the bottom of p. 1247, "symmetry" should be "asymmetry."

**Possible Observation of Light Neutral Bosons in Nuclear Emulsions.** F. W. N. DE BOER and R. VAN DANTZIG [Phys. Rev. Lett. **61**, 1274 (1988)].

The vertical scale in Fig. 2 (and in Fig. 8 of Ref. 10) has to be divided by a factor of 2.

In the dashed curve in Fig. 2 obscuration has been taken into account but not the finite grain density.

**Spectrum of  $J^P=2^+$  Mesons.** S. K. BOSE and E. C. G. SUDARSHAN [Phys. Rev. Lett. **62**, 1445 (1989)].

The terms  $P^*3$  and  $Q^*3$  that appear under the integral sign of Eq. (3) should correctly read as  $(P^*)^3$  and  $(Q^*)^3$ , respectively.

**Two-Photon Absorption of Nonclassical Light.** J. GEABANACLOCHE [Phys. Rev. Lett. **62**, 1603 (1989)].

In Fig. 1, the dashed line is for *amplitude*-squeezed light and the dash-dotted line is for *phase*-squeezed light, contrary to what the figure caption reads.