

### Renormalization of the Statistics Parameter in Three-Dimensional Electrodynamics

Gordon W. Semenoff,<sup>(1)</sup> Pasquale Sodano,<sup>(2),(a)</sup> and Yong-Shi Wu<sup>(3)</sup>

<sup>(1)</sup>*Department of Physics, University of British Columbia, Vancouver, Canada V6T 2A6*

<sup>(2)</sup>*Department of Physics, Boston University, 590 Commonwealth Avenue, Boston, Massachusetts 02215*

<sup>(3)</sup>*Department of Physics, University of Utah, Salt Lake City, Utah 84112*

(Received 17 October 1988)

We argue that the  $\beta$  function for the statistics parameter in (2+1)-dimensional electrodynamics is zero to all orders in perturbation theory beyond one loop. We show that there can be finite radiative corrections from massless charged scalar fields.

PACS numbers: 12.20.Ds, 05.30.-d, 11.10.Gh, 11.15.Bt

Gauge theories in (2+1) dimensions have several features not seen in their (3+1)-dimensional counterparts. These center on the possible appearance of the Chern-Simons three-form in the action<sup>1</sup> and are connected with parity anomalies<sup>2</sup> and exotic charge,<sup>2</sup> angular momentum,<sup>3</sup> and statistics.<sup>4</sup> Original motivation for their study was their resemblance to the high-tem-

perature limit of gauge theories in (3+1) dimensions. More recently, there has been interest in (2+1)-dimensional electrodynamics in models of certain condensed-matter phenomena, particularly high- $T_c$  superconductors<sup>5-7</sup> and the quantized Hall effect.<sup>8</sup>

The Euclidean action of scalar electrodynamics with both Chern-Simons and Maxwell kinetic terms for the gauge fields is

$$S = \int d^3x \left[ (D_\mu \phi_0)^* (D_\mu \phi_0) + m_0^2 \phi_0^* \phi_0 + i \frac{\alpha_0}{4\pi} \epsilon_{\mu\nu\lambda} A_\mu^0 \partial_\nu A_\lambda^0 + \frac{1}{4e_0^2} F_{\mu\nu}^0 F_{\mu\nu}^0 \right], \tag{1}$$

where  $D_\mu = \partial_\mu + ig_0 A_\mu^0$ . This is a prototype for the effective field theory of the  $CP^1$  model discussed in Ref. 9 (see also Ref. 10). The Chern-Simons term gives the  $\phi$  quanta fractional statistics.<sup>4</sup> (With the Maxwell term this is only true at large distances.) The gauge field and  $e_0^2$  have dimension 1 and  $g_0$  and  $\alpha_0$  are dimensionless. This field theory is superrenormalizable with computable, finite physical parameters. There is considerable interest in a model with only Chern-Simons kinetic terms.<sup>4-10</sup> It is defined by the limit  $e_0^2 \rightarrow \infty$  where it is renormalizable with dimensionless bare coupling  $g_0 |\alpha_0|^{-1/2}$  and  $F_{\mu\nu}^0 F_{\mu\nu}^0$  is an irrelevant operator. Since all physical parameters are not finite in this limit, renormalization is required. In Ref. 9 it was argued that the

dimensionless statistics parameter  $g_0^2/\alpha_0$  can have an interesting renormalization flow. We show in this paper that this is not the case. When the renormalized mass  $m^2 \neq 0$ , the corresponding  $\beta$  function is zero to all orders in loop expansion. For the critical case  $m^2 = 0$  (although there is no symmetry which prevents radiative mass generation, it is natural at a critical point for the scalar to be massless) we demonstrate that  $\beta$  is zero to two-loop order and we conjecture that it is zero to all orders. We also show that, when  $m^2 = 0$ ,  $g_0^2/\alpha_0$  has finite renormalization in higher loops. With  $\phi_0 = Z^{1/2} \phi$ ,  $A_\mu^0 = Z_3^{1/2} A_\mu$ ,  $g_0 = Z^{-1} Z_3^{-1/2} Z_2 g$ ,  $\alpha_0 = Z_3^{-1} Z_a \alpha$ ,  $m_0^2 = m^2 + \delta m^2$ , and the Ward-Takahashi identity  $Z = Z_2$ , the action is

$$S = \int \left\{ Z [\partial_\mu \phi^* \partial_\mu \phi + (m^2 + \delta m^2) \phi^* \phi] + \frac{Z_3}{4e_0^2} F_{\mu\nu} F_{\mu\nu} + Z_a i \frac{\alpha}{4\pi} \epsilon_{\mu\nu\lambda} A_\mu \partial_\nu A_\lambda + Z g A_\mu \phi^* i \overleftrightarrow{\partial}_\mu \phi + Z g^2 A_\mu A_\mu \phi^* \phi \right\}. \tag{2}$$

$Z_a$  is fixed by the requirement that the renormalized Chern-Simons term has coefficient  $\alpha/4\pi$ . The physical statistics parameter is  $g^2/\alpha = Z_a g_0^2/\alpha_0$ . Propagators, vertices, and counterterms are summarized in Fig. 1. We use  $e_0^2$  as a gauge-invariant cutoff to be taken to  $\infty$  at the end of computations. To see that when  $m^2 > 0$ ,  $Z_a$  receives no contributions beyond one loop,<sup>11</sup> consider the irreducible Euclidean  $N$ -photon correlation function for  $N \geq 4$  in the one-loop approximation  $\Gamma_{\mu_1 \dots \mu_N}(p_1; \dots; -\sum_1^{N-1} p_k)$  shown in Fig. 2.  $\Gamma$  is symmetric. Radiative corrections to  $Z_a$  with two or more loops are one-photon-irreducible graphs obtained either by sewing together all but two legs of  $\Gamma$  [see Fig. 3(a)], then finding the term linear in the external momentum and antisymmetric in vector indices of the resulting two-point function,

$$\lim_{p \rightarrow 0} \epsilon_{\mu\nu\rho} \frac{\partial}{\partial p_\rho} \int dp_3 dp_4 \dots \Gamma_{\mu\nu\lambda_3 \dots \lambda_N} \left( p; -p; p_3; \dots; -\sum_3^{N-1} p_k \right) \mathcal{K}_{\lambda_3 \dots \lambda_N}(p_3; p_4; \dots), \tag{3}$$




FIG. 1. Propagators and vertices for scalar electrodynamics. The topological mass is  $\mu = ae\delta/2\pi$ ,  $e\delta \rightarrow \infty$ , and  $\gamma$  is the gauge-fixing parameter.




FIG. 3. How  $\Pi_{\mu\nu}$  is constructed from  $\Gamma$ .  $\tilde{\mathcal{H}}$  depends on  $p$  whereas  $\mathcal{H}$  is independent of  $p$ .

or by sewing two  $\Gamma$ 's together in all possible (one-photon irreducible) ways [Fig. 3(b)],

$$\lim_{p \rightarrow 0} \epsilon_{\mu\nu\rho} \frac{\partial}{\partial p_\rho} \int dl_1 \cdots dq_1 \cdots \Gamma_{\mu\lambda\dots}(p;l_1;\dots) \Gamma_{\nu\rho\dots}(-p;q_1;\dots) \tilde{\mathcal{H}}_{\lambda\dots\rho\dots}(p;l_1;\dots;q_2;\dots). \tag{4}$$

When  $m^2 > 0$ ,  $\Gamma$  is analytic at  $p_i^{\mu i} = 0$ .  $\tilde{\mathcal{H}}$  depends on  $p$  where as  $\mathcal{H}$  does not. The Ward-Takahashi identity is

$$0 = p_i^{\nu i} \Gamma_{\mu_1 \dots \nu_1 \dots \mu_N} \left[ p_1; \dots; p_i; \dots; - \sum_k^{N-1} p_k \right].$$

Taking a derivative by  $p_i$ , setting  $p_i = 0$ , and using analyticity yields

$$0 = \Gamma_{\mu_1 \dots \nu_1 \dots \mu_N} \left[ p_1; \dots; 0; \dots; - \sum_1^{N-1} p_k \right],$$

This immediately implies that (4) vanishes. It further implies

$$\lim_{p \rightarrow 0} \frac{\partial}{\partial p_\rho} \Gamma_{\mu\nu\dots}(p; -p; \dots) = \lim_{p \rightarrow 0} \left[ \frac{\partial}{\partial p_\rho} \Gamma_{\mu\nu\dots}(p; 0; \dots) - \frac{\partial}{\partial p_\rho} \Gamma_{\mu\nu\dots}(0; -p; \dots) \right] = 0,$$

and (3) also vanishes. (This generalizes to other massive matter such as spinor and vector fields. It is also valid at finite temperature. There the Euclidean momentum vectors have one component the discrete Matsubara frequency  $p_i^0 = 2\pi k_B T n_i$  with  $T$  the temperature,  $n_i$  an integer, and the other two components continuous variables. For  $N > 4$ , Euclidean  $N$ -photon functions have unique extensions to functions of complex momenta  $p_i^0$ . With massive matter, these functions are analytic at zero frequency and momentum and the Ward-Takahashi identity also holds. The above reasoning shows that no two-photon vertex with one power of derivatives is generated by radiative corrections beyond one loop at finite temperature.) Thus, given the known *finite* one-loop corrections from both scalars and spinors,<sup>2</sup> their contributions to the  $\beta$  function for the




FIG. 2. The  $N$ -photon correlation function at one-loop order is the symmetrized sum with all combinations of both three- and four-point vertices.




FIG. 4. One-loop photon self-energy.

statistics parameter is zero.

This leaves the possibility of a nonzero  $\beta$  function from massless charged fields where the  $N$ -point functions  $\Gamma$  cannot be assumed analytic at zero external momenta. The zero mass limit which defines a massless field theory may not commute with the zero momentum limit which determines the renormalization of a coupling constant. However, since the  $\beta$  function is independent of momentum, it will be zero in the massless limit as it is in massive theory. This will be verified by explicit calculation to two-loop order in the following. We conjecture that it holds to all orders. This argument does not exclude finite renormalizations and we shall find explicit nonzero finite corrections at two-loop order from massless scalars.

The one-loop correction to the photon two-point function from charged scalars (Fig. 4) is

$$\Pi_{\mu\nu}^{(1)}(p) = (p^2 \delta_{\mu\nu} - p_\mu p_\nu) \frac{g^2}{8\pi} \left\{ -\frac{2\sqrt{m^2}}{p^2} + \frac{p^2 + 4m^2}{p^2 \sqrt{p^2}} \arcsin \left[ \frac{p^2}{p^2 + 4m^2} \right]^{1/2} \right\}. \quad (5)$$

Although the loop integral is potentially divergent and requires a cutoff at intermediate stages of calculation, the result is finite. Its antisymmetric part vanishes and when  $m^2 > 0$  its symmetric part has a local limit  $(p^2 \delta_{\mu\nu} - p_\mu p_\nu) \times g^2/24\pi\sqrt{m^2}$  which contributes to  $Z_3$ . Figure 5 depicts two-loop self-energy diagrams. The corresponding integrals are finite for  $e_0^2 < \infty$  and by power counting are logarithmically divergent when  $e_0^2 \rightarrow \infty$ . The contributions of Figs. 5(b)-5(e) to  $(1/2p^2)\epsilon_{\mu\nu\lambda}\Pi_{\mu\nu}^{(2)}p_\lambda$  cancel immediately upon combining the integrals. Figure 5(a) can be reduced to the Feynman parameter integral

$$\frac{g^4}{8\pi^2\alpha} \int_0^1 dx_1 \cdots dx_4 \theta(1-x_1-\cdots-x_4) \frac{1}{[(1-x_1-x_2)(x_1+\cdots+x_4)-(x_3+x_4)^2]^{5/2}} \times \ln \frac{\mu^2(1-x_1-\cdots-x_4)(1-x_1-x_2)+m^2(x_1+\cdots+x_4)(1-x_1-x_2)+p^2\Omega}{m^2(x_1+\cdots+x_4)(1-x_1-x_2)+p^2\Omega} \quad (6)$$

where  $\mu = ae_0^2/2\pi$  and

$$\Omega = [(1-x_1-x_2)(x_2+x_3)-(x_3+x_4)^2] - \frac{[(1-x_1-x_2)(x_2+x_3)-(x_3+x_4)x_3]^2}{(1-x_2-x_3)(x_1+\cdots+x_4)-(x_3+x_4)^2}. \quad (7)$$

Figure 5(f) reduces to

$$-\frac{g^4}{8\pi^2\alpha} \int_0^1 dx_1 dx_2 \theta(1-x_1-x_2) \frac{x_1 x_2}{[(1-x_1-x_2)(x_1+x_2)-x_1 x_2]^{5/2}} \times \ln \frac{\mu^2(1-x_1-x_2)(1-x_1)+m^2(1-x_1)(x_1+x_2)+p^2\tilde{\Omega}}{m^2(1-x_1)(x_1+x_2)+p^2\tilde{\Omega}}, \quad (8)$$

$$\tilde{\Omega} = x_2(1-x_1-x_2) - \frac{x_2^2(1-x_1-x_2)^2}{(1-x_1-x_2)(x_1+x_2)+x_1 x_2}. \quad (9)$$

If we keep  $m^2 > 0$  and put  $p^2 = 0$ , these two contributions cancel for all values of  $\mu^2 > 0$ , consistent with the no-renormalization argument above. On the other hand, if we first put  $m^2 = 0$  and then  $\mu^2/p^2$  large (since  $e_0^2 \rightarrow \infty$ ), the divergent leading terms proportional to  $\ln(\mu^2/p^2)$  cancel. Therefore, as expected, there is no infinite contribution to  $Z_\alpha$ . After some standard manipulations,<sup>12</sup> the finite parts are

$$\frac{g^4}{8\pi^2\alpha} \int_0^1 d\rho d\lambda dx dy \theta(1-y-\rho) \frac{\lambda^{1/2}\rho}{[1-\lambda+\lambda\rho(1-\rho)]^{5/2}} \times \ln \frac{\rho(1-\rho)(1-\lambda)(1-\lambda\rho)}{[(1-\lambda\rho)(\rho x+y)-\lambda y^2][1-\lambda+\lambda\rho(1-\rho)] - [\rho(1-\lambda\rho)x+(1-\lambda)y]^2} \approx -2.74 \frac{g^4}{8\pi^2\alpha}, \quad (10)$$

where the approximate result is obtained numerically.

To couple spinors we add  $\int \psi^\dagger [\gamma_\mu (i\partial_\mu - i\tilde{g}_0 A_\mu^0) + M_0] \psi_0$  to (1). Generalization of the arguments above shows that




FIG. 5. Two-loop photon self-energy.

there are no contributions to  $Z_\alpha$  beyond one loop at either zero or finite temperature when  $M \neq 0$ . At one loop, the correction is known<sup>2</sup> at  $T=0$  and can be obtained at  $T \neq 0$  as

$$\lim_{p \rightarrow 0} \frac{1}{2p^2} \epsilon_{\mu\nu\lambda} \Pi_{\mu\nu}^{(1)} p_\lambda = \frac{\tilde{g}^2}{8\pi} \tanh \frac{M}{k_B T}.$$

When  $M=0$  the one-loop correction vanishes but there could be finite corrections from higher loops.

Equation (5) indicates a radiatively generated Maxwell term at one-loop (and higher) order. We determine  $Z_3$  so that the renormalized Maxwell term has coefficient  $1/4e^2$ . Even when the bare action has only a Chern-Simons term, a Maxwell term is generated in the effective action. In that case it is possible to resume perturbation theory using the renormalized Maxwell term in the Feynman rules which are then superrenormalizable. The present analysis also applies to the singular case where  $1/e^2=0$  which remains marginally renormalizable. In summary, for scalars when  $m=0$ ,  $Z_\alpha=1+2.74g^4/2\pi\alpha^2+\dots$ ,  $Z_3=e_0^2/e^2+\dots$ ; when  $m^2 \neq 0$ ,  $Z_\alpha=1$ ,  $Z_3=e_0^2/e^2-e_0^2g^2/24\pi\sqrt{m^2}+\dots$ . For spinors when  $M=0$ ,  $\delta Z_\alpha$ =(finite corrections from  $\geq$  two loops),  $Z_3$  is affected at two and higher loops; when  $M^2 \neq 0$ ,  $\delta Z_\alpha = -(\tilde{g}^2/2\alpha) \tanh(M/k_B T)$ ,  $\delta Z_3 = -e_0^2\tilde{g}^2/12\pi\sqrt{M^2}+\dots$  from one- and higher-loop vacuum polarization. In the massive theories we have  $Z_\alpha$  exactly. In both cases there are no infinite corrections to  $\alpha_0$  and the corresponding  $\beta$  function vanishes. However, there is finite renormalization from massless particles in higher loops and from massive spinors at one loop. There can be an infinite renormalization from charged massive vectors at one loop.<sup>13</sup> The latter [as well as the  $CP^1$  model to which (1) is approximate] is a nonrenormalizable theory and the perturbative expansion is unreliable. Also, our results for renormalizable theories are perturbative and would be improved by demonstrating Borel summability of the perturbation series. Finally, we speculate on hysteresis where we vary parameters, i.e.,  $T$ ,  $m^2$ ,  $e^2$  of scalar electrodynamics. The statistics parameter is unaffected if we avoid critical points. If we go to a critical point the statistics parameter gets finite computable corrections. We then leave the critical point with renormalized fractional statistics.

G.W.S. is supported by the Natural Sciences and Engineering Research Council of Canada. P.S. is supported by U.S. Department of Energy Grant No. DE/87/AC02/86ER40284 and the Istituto Nazionale di

Fisica Nucleare. Y.-S.W. is supported by U.S. National Science Foundation Grant No. PHY-8706501.

<sup>(a)</sup>Permanent address: Dipartimento di Fisica, Universita di Perugia, and INFN Sezione di Perugia, Via Elce di Sotto 10, 06100 Perugia, Italy.

<sup>1</sup>W. Siegel, Nucl. Phys. **B156**, 135 (1979); J. Schonfeld, Nucl. Phys. **B185**, 157 (1981); R. Jackiw and S. Templeton, Phys. Rev. D **4**, 2291 (1981); S. Deser, R. Jackiw, and S. Templeton, Phys. Rev. Lett. **48**, 975 (1982), and Ann. Phys. (N.Y.) **140**, 372 (1982).

<sup>2</sup>A. Niemi and G. Semenoff, Phys. Rev. Lett. **51**, 2077 (1983), and Phys. Rep. **135**, 3 (1987); N. Redlich, Phys. Rev. D **29**, 2366 (1984).

<sup>3</sup>M. Paranjape, Phys. Rev. Lett. **55**, 3290 (1985); G. Semenoff and R. Wijerardhanna, Phys. Lett. B **184**, 397 (1987).

<sup>4</sup>G. Semenoff, Phys. Rev. Lett. **61**, 517 (1988); J. Leinaas and J. Myrheim, Nuovo Cimento **B37**, 1 (1977); Y.-S. Wu, Phys. Rev. Lett. **52**, 2102, 111 (1984); F. Wilczek, Phys. Rev. Lett. **49**, 957 (1982); F. Wilczek and A. Zee, Phys. Rev. Lett. **51**, 2250 (1983), and **52**, 2111 (1984); Y.-S. Wu and A. Zee, Phys. Lett. **147B**, 325 (1984); D. Arovas, R. Schrieffer, F. Wilczek, and A. Zee, Nucl. Phys. **B251 FS13**, 117 (1985); G. Semenoff and P. Sodano, Phys. Rev. Lett. **57**, 1195 (1986).

<sup>5</sup>P. Anderson, Science **235**, 1196 (1986); S. Kivelson, D. Rokhsar, and J. Sethna, Phys. Rev. B **35**, 8865 (1987); R. Schrieffer, X. Wen, and S. Zhang, Phys. Rev. Lett. **60**, 944 (1988).

<sup>6</sup>X. Wen and A. Zee, Phys. Rev. Lett. **61**, 1025 (1988); D. Haldane, Phys. Rev. Lett. **61**, 1029 (1988).

<sup>7</sup>I. Dzyaloshinski, A. Polyakov, and P. Wiegmann, Phys. Lett. A **127**, 112 (1988); P. Wiegmann, Phys. Rev. Lett. **60**, 821 (1988); A. Polyakov, Mod. Phys. Lett. A **3**, 325 (1988).

<sup>8</sup>S. Zhang, H. Hansson, and S. Kivelson, Santa Barbara report, 1988 (to be published); S. Girvin, in *The Quantum Hall Effect*, edited by R. Prange and S. Girvin (Springer-Verlag, Berlin, 1986), and Chap. 10, references therein.

<sup>9</sup>A. Polyakov, in *Fields Strings and Critical Phenomena*, Les Houches, France, 1988 (to be published); D. Deser and N. Redlich, Phys. Rev. Lett. **61**, 1541 (1988).

<sup>10</sup>A. Polyakov, *Gauge Fields and Strings* (Harwood, London, 1987).

<sup>11</sup>S. Coleman and B. Hill, Phys. Lett. **159B**, 184 (1985); Y. Kao and M. Suzuki, Phys. Rev. D **31**, 2137 (1985); M. Bernstein and T. Lee, Phys. Rev. D **32**, 1020 (1985).

<sup>12</sup>Y. B. Dai and Y.-S. Wu, Scientia Sinica **19**, 65 (1976).

<sup>13</sup>C. Hagen, P. Panigrahi, and S. Ramaswamy, Phys. Rev. Lett. **61**, 389 (1988).