

## *A Dependence of $J/\psi$ and $\psi'$ Production at 800 GeV/c*

D. M. Alde, H. W. Baer, T. A. Carey, G. T. Garvey, A. Klein, C. Lee, M. J. Leitch, J. Lillberg,  
P. L. McGaughey, C. S. Mishra, J. M. Moss, and J. C. Peng  
*Los Alamos National Laboratory, Los Alamos, New Mexico 87545*

C. N. Brown, W. E. Cooper, and Y. B. Hsiung  
*Fermilab, Batavia, Illinois 60510*

M. R. Adams  
*University of Illinois at Chicago, Chicago, Illinois 60680*

R. Guo and D. M. Kaplan  
*Northern Illinois University, DeKalb, Illinois 60115*

R. L. McCarthy  
*State University of New York at Stony Brook, Stony Brook, New York 11794-3800*

G. Danner and M. Wang  
*Case Western Reserve University, Cleveland, Ohio 44106*

M. Barlett and G. Hoffmann  
*University of Texas, Austin, Texas 78712*  
(Received 31 July 1990)

The yield of  $J/\psi$  and  $\psi'$  vector-meson states has been measured for 800-GeV protons incident on deuterium, carbon, calcium, iron, and tungsten targets. A depletion of the yield per nucleon from heavy nuclei is observed for both  $J/\psi$  and  $\psi'$  production. This depletion exhibits a strong dependence on  $x_F$  and  $p_T$ . Within experimental errors the depletion is the same for the  $J/\psi$  and  $\psi'$ .

PACS numbers: 13.85.Ni, 25.40.Ve

The recent observation<sup>1</sup> of  $J/\psi$  suppression in relativistic heavy-ion central collisions, predicted as a signature of quark-gluon-plasma formation,<sup>2</sup> has generated renewed interest in the subject of hadron-induced  $J/\psi$  production.<sup>3-7</sup> In order to elucidate the origin of  $J/\psi$  suppression, it is desirable to examine the characteristics of hadronic  $J/\psi$  production in heavy versus light nuclei, which may mirror certain aspects of conditions encountered in central versus peripheral collisions in heavy-ion interactions. Since existing theoretical models predict that gluon-gluon fusion is the dominant mechanism for hadron-induced quarkonium production at 800 GeV,<sup>8,9</sup> measurement of the nuclear dependence of  $J/\psi$  production might also reveal possible modification of the gluon distribution in nuclei.<sup>10</sup>

We report here, from Fermilab experiment 772, the first  $A$ -dependence measurements of  $J/\psi$  and  $\psi'$  production in 800-GeV proton-nucleus collisions. Previous studies of nuclear effects on  $J/\psi$  production, performed at lower energies,<sup>11-14</sup> did not have adequate statistics or mass resolution to extract information on the  $\psi'$  resonance.

The E605/772 spectrometer<sup>15</sup> at Fermilab was used to measure dimuons produced in 800-GeV proton-nucleus collisions. A 5.2-m-thick absorber composed of copper,

graphite, and polyethylene, located at the exit aperture of the 15-m spectrometer magnet, absorbed low-energy backgrounds and allowed incident proton intensities of up to  $10^{11}$  protons per second to interact with targets of  $\sim 10$  g/cm<sup>2</sup> thickness. Solid nuclear targets (C, Ca, Fe, W) and a liquid-deuterium target were interchanged frequently during the experiment. A detailed Monte Carlo calculation was performed to correct for the small acceptance variations among different targets. This procedure, together with many beam-intensity and electronic lifetime monitors, yielded a systematic uncertainty on the ratios of yields from the various targets of  $\sim 2\%$ . Only statistical errors are shown in the figures. Further details on the measurements may be found in Ref. 16.

The data correspond to  $\sim 8 \times 10^{15}$  incident protons on target yielding  $\sim 10^5$   $J/\psi$ 's and  $\sim 1.2 \times 10^4$   $\psi'$ 's. The mass resolution of  $\sim 150$  MeV at a mass of 3 GeV gives excellent separation between the  $J/\psi$  and  $\psi'$  peaks (Fig. 1, inset). To extract the peak areas the spectrum was fitted with a combination of asymmetric Gaussian peaks plus a polynomial to represent the Drell-Yan (DY) continuum. Nuclear dependence of the DY continuum has been presented elsewhere.<sup>16</sup>

Figure 1 shows the heavy-nucleus to deuterium ratio per nucleon  $R$  integrated over Feynman  $x$  ( $x_F$ ) and




FIG. 1. The ratios of heavy-nucleus to deuterium integrated yields for the  $J/\psi$  and  $\psi'$  resonances and the Drell-Yan continuum. Inset: The raw (no acceptance correction) dimuon invariant-mass spectrum.

transverse momentum ( $p_t$ ) for the  $J/\psi$ ,  $\psi'$ , and the DY continuum ( $4 \leq M_{\mu\mu} \leq 9$  and  $M_{\mu\mu} \geq 11$  GeV) versus  $A$ . Mean values of the kinematic variables for the  $J/\psi$  and  $\psi'$  resonances, averaged over the spectrometer acceptance, are  $\langle x_F \rangle \sim 0.27$  and  $\langle p_t \rangle \sim 0.7$  GeV. In contrast to the DY data, which give a value of  $R$  very close to unity, a large depletion of the  $J/\psi$  and  $\psi'$  yields is found in the nuclear targets. A significant new result seen in Fig. 1 is that the depletion is the same within errors for the  $J/\psi$  and  $\psi'$ . To describe the  $A$  dependence of the  $J/\psi, \psi'$  depletion, we use the usual parametrization

$$\sigma_A = \sigma_N A^\alpha. \quad (1)$$

The curve in Fig. 1 corresponds to the best fit with  $\alpha = 0.92$ . The  $\sim 2\%$  normalization error translates into an approximately constant systematic error in  $\alpha$  of  $\sim 0.008$ . The validity of Eq. (1) for  $J/\psi$  production has not been well tested, since previous experiments typically measured only two targets. Katsanevas *et al.*<sup>14</sup> noted that the form  $A^\alpha$  fails to describe their data, when combined with earlier NA data.<sup>13</sup> Figure 1 shows that  $A^\alpha$  gives an adequate, though not excellent, description of the present data.

Figure 2 shows  $R$  for the  $J/\psi$  as a function of  $x_F$  and  $p_t$ . The  $J/\psi$  depletion in heavy targets is most pronounced at larger values of  $x_F$  and at smaller values of  $p_t$ . This statement also applies to  $R(x_F; p_t)$  for the  $\psi'$ . The observed  $x_F$  and  $p_t$  dependences are in qualitative agreement with previous proton-induced  $J/\psi$  production data,<sup>13,14</sup> as well as the pion- and antiproton-induced  $J/\psi$  production data.<sup>13,14</sup>

In Fig. 3 we show  $\alpha$  for the  $J/\psi$  vs  $x_F, x_2$  (defined




FIG. 2. The ratios of heavy-nucleus to deuterium yields for the  $J/\psi$  and  $\psi'$  resonances vs  $x_F$  and  $p_t$ .

below), and  $p_t$ , as determined by fits to  $R$  for all targets. Equation (1) describes the data quite well when binned versus  $x_F$  or  $p_t$ . Also shown are  $\alpha(x_F)$  and  $\alpha(x_2)$  for 200-GeV proton production of the  $J/\psi$  from NA3 (Ref. 13) based on two targets,  $^2\text{H}$  and Pt. Comparison of the two data sets shows that  $\alpha(x_F)$  depends little on beam energy.

In the simplest gluon-gluon fusion models,<sup>8,9</sup> the quarkonium cross section is given by the convolution of the process  $gg \rightarrow Q\bar{Q}$  with the gluon structure functions  $G(x_1)$  and  $G(x_2)$ , where  $x_1$  and  $x_2$  are the Bjorken  $x$  of the gluons in the beam and target hadrons, respectively. This simple model has been used to extract information on the gluon structure functions from  $J/\psi$  and  $\Upsilon$  production data.<sup>13,17</sup> Here it is assumed that  $x_1$  and  $x_2$  are related to the observed quantities  $m$  and  $x_F$  through the relations

$$m^2 = x_1 x_2 s, \quad x_F = x_1 - x_2, \quad (2)$$




FIG. 3. The parameter  $\alpha$  for the  $J/\psi$  resonance (circles) and the Drell-Yan continuum (diamonds) as determined from fits to all four heavy-target ratios. Also shown is  $\alpha(x_F, x_2)$  for the 200-GeV data from NA3 (Ref. 13) (squares).

where  $s$  is the center-of-mass energy squared. Strictly speaking,  $m$  in Eq. (2) can be the mass of any  $c\bar{c}$  state produced by gluon fusion (or  $q\bar{q}$  fusion) which subsequently decays into the  $J/\psi$ . We use the mass of the  $J/\psi$  to calculate  $x_2$  in Fig. 3. Comparison of the 800- and 200-GeV results clearly indicates that the data do not scale with  $x_2$ , a parameter intrinsic to the target-parton structure function. Changing the choice of mass in Eq. (2) to 3.5 GeV shifts the 200-GeV data by  $x_2 \sim 0.01$  further to the right relative to the 800-GeV data. Thus the  $J/\psi$  suppression is not a simple manifestation of the small- $x_2$  shadowing seen in deep-inelastic lepton scattering.<sup>18</sup>

Several models, aimed at a unified description of  $J/\psi$  production in hadron-nucleus and nucleus-nucleus collisions, have considered the effect of attenuation of  $c\bar{c}$  states by secondary reactions of the  $J/\psi$  with some combination of the remaining nucleons of the target plus hadronic debris formed in the collision (comovers).<sup>3,5-7</sup> The evolution of the  $J/\psi$  from the initial  $c\bar{c}$  state, where the interaction cross section may be very small due to color-transparency effects,<sup>19</sup> to the final state of hadronic dimensions is characterized by an exponential time dependence. At present, this time dependence has not been characterized experimentally. Although attenuation models have been directed primarily toward the central production region, their extension to the present  $x_F$  range is straightforward. It is clear that these models predict a value of  $\alpha$  nearer unity at large  $x_F$  for two reasons. First, the more energetic the  $J/\psi$ , the longer it stays in its (presumed) spatially small, color-transparent state. Second, for the most energetic  $J/\psi$ 's, the density of comovers decreases. The observation of a significant suppression in the yield of the  $J/\psi$  at large  $x_F$  implies that attenuation cannot be the complete explanation of the  $A$  dependence of hadronic  $J/\psi$  production. Additional evidence against the comover picture is found in beam-dump measurements of the  $A$  dependence of inclusive charm production.<sup>20,21</sup> There it is found that  $\alpha$  is substantially less than unity. Presumably open-charm channels should not suffer attenuating reactions in the same way as  $c\bar{c}$  states. The suppression of open-charm production may be connected to the well-known suppression of strangeness production<sup>22,23</sup> at large  $x_F$ .

One model<sup>4</sup> is apparently able to qualitatively reproduce the  $x_F$  dependence of  $\alpha$ . The authors of this model postulate intrinsic  $c\bar{c}$  components in the wave function of the incident hadron to achieve these features. It remains to be determined whether or not the magnitude of the intrinsic charm in the proton can account for the present data.

The fact that the  $A$  dependence of  $J/\psi$  and  $\psi'$  production is the same within errors provides an additional constraint on the hadronic attenuation picture. The radii of the  $J/\psi$  and  $\psi'$  differ by almost a factor of 2 in potential models.<sup>24</sup> Direct interpretation of this difference is complicated by the fact that the  $J/\psi$  is probably produced in

part by decays from  $\chi_c$  states which have radii comparable to the  $\psi'$ . Nevertheless, the present data indicate no dependence on final hadronic size.

Finally, we turn to the  $p_t$  dependence of  $\alpha$ . Figure 3 shows that the increase in  $\alpha$  at large  $p_t$  is somewhat greater for the  $J/\psi$  than for the DY continuum. This has been anticipated by models<sup>25,26</sup> which describe the  $p_t$  dependence of hadronic  $J/\psi$  production in terms of initial-state-final-state partonic multiple scattering. The ratio of the  $J/\psi$  to DY  $p_t$  dependence plays an important role in understanding the significance of  $J/\psi$  production in heavy-ion collisions. Although detailed model analyses of the NA38 results<sup>1</sup> are still being debated, the results seen here are in qualitative agreement with those from heavy-ion-induced  $J/\psi$  production, possibly indicating a common origin.

In summary, we have observed that production of the  $J/\psi$  and  $\psi'$  resonances with 800-GeV protons is strongly suppressed in heavy nuclei. The  $A$  dependence is well described by the simple expression  $A^\alpha$ , with a value of  $\alpha$  which depends on  $x_F$  and  $p_t$ . Comparison of the present data with the previous 200-GeV data shows that  $\alpha$  scales well with  $x_F$ , but not with  $x_2$ . Two popular models, small- $x$  shadowing and hadronic attenuation, have difficulties explaining either the  $x_2$  or the  $x_F$  dependence of existing data. It is clear that much remains to be understood about the nature of charmonium production and propagation in nuclei. In particular, it is important to have measurements of the  $A$  dependence of  $J/\psi$  and  $\psi'$  production over a much wider kinematic range, especially at negative  $x_F$  where almost no experimental data exists. In addition to charmonium production, measurement of the  $A$  dependence of open-charm channels, e.g.,  $D$  mesons, would greatly clarify such issues as intrinsic charm in light hadrons and comover attenuation.

We would like to acknowledge the efforts of the Fermilab Research and Accelerator Divisions and funding from the U.S. Department of Energy and the National Science Foundation. We also thank the Japanese Ministry of Education for providing parts of the spectrometer.

<sup>1</sup>C. Baglin *et al.*, Phys. Lett. B **220**, 471 (1989).

<sup>2</sup>T. Matsui and H. Satz, Phys. Lett. B **178**, 416 (1986).

<sup>3</sup>S. J. Brodsky and A. H. Mueller, Phys. Lett. B **206**, 685 (1988).

<sup>4</sup>S. J. Brodsky and P. Hoyer, Phys. Rev. Lett. **63**, 1566 (1989).

<sup>5</sup>J. P. Blaizot and J. Y. Ollitrault, Phys. Lett. B **217**, 386 (1989).

<sup>6</sup>S. Gavin and M. Gyulassy, Phys. Lett. B **207**, 257 (1988).

<sup>7</sup>S. Gavin and R. Vogt (to be published).

<sup>8</sup>R. Baier and R. Ruckl, Z. Phys. C **19**, 251 (1983).

<sup>9</sup>V. Barger, W. Y. Keung, and R. J. N. Phillips, Z. Phys. C **6**, 169 (1980).

<sup>10</sup>F. E. Close, J. Qiu, and R. G. Roberts, Phys. Rev. D **40**,

2820 (1989).

- <sup>11</sup>T. M. Antipov *et al.*, Phys. Lett. **76B**, 235 (1978).  
<sup>12</sup>M. J. Corden *et al.*, Phys. Lett. **110B**, 415 (1982).  
<sup>13</sup>J. Badier *et al.*, Z. Phys. C **20**, 101 (1983).  
<sup>14</sup>S. Katsanevas *et al.*, Phys. Rev. Lett. **60**, 2121 (1988).  
<sup>15</sup>D. E. Jaffe *et al.*, Phys. Rev. D **40**, 2777 (1989).  
<sup>16</sup>D. M. Alde *et al.*, Phys. Rev. Lett. **64**, 2479 (1990).  
<sup>17</sup>A. D. Martin, R. G. Roberts, and W. J. Stirling, Phys. Rev. D **37**, 1161 (1988).  
<sup>18</sup>J. Ashman *et al.*, Phys. Lett. B **202**, 603 (1988).  
<sup>19</sup>G. Bertsch *et al.*, Phys. Rev. Lett. **47**, 297 (1981).  
<sup>20</sup>H. Cobbaert *et al.*, Phys. Lett. B **191**, 456 (1987).  
<sup>21</sup>M. E. Duffy *et al.*, Phys. Rev. Lett. **55**, 1816 (1985).  
<sup>22</sup>K. B. Luke, in *Proceedings of Hadronic Matter in Collisions*, edited by J. Rafelski and P. Carruthers (World Scientific, Singapore, 1989), p. 571.  
<sup>23</sup>S. Fredericksson *et al.*, Phys. Rep. **114**, 189 (1987).  
<sup>24</sup>W. Kwong, J. L. Rosner, and C. Quigg, Annu. Rev. Nucl. Part. Sci. **37**, 325 (1987).  
<sup>25</sup>S. Gavin and M. Gyulassy, Phys. Lett. B **214**, 241 (1988).  
<sup>26</sup>J. Hufner, Y. Krihara, and H. J. Pirner, Phys. Lett. B **215**, 218 (1988).