

### Measurement of the Ratio $B(W \rightarrow \tau \nu)/B(W \rightarrow e \nu)$ in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV

F. Abe,<sup>(9)</sup> D. Amidei,<sup>(4)</sup> G. Apollinari,<sup>(16)</sup> M. Atac,<sup>(4)</sup> P. Auchincloss,<sup>(15)</sup> A. R. Baden,<sup>(6)</sup> N. Bacchetta,<sup>(11)</sup> M. W. Bailey,<sup>(14)</sup> A. Bamberger,<sup>(4),(a)</sup> P. de Barbaro,<sup>(15)</sup> B. A. Barnett,<sup>(8)</sup> A. Barbaro-Galtieri,<sup>(10)</sup> V. E. Barnes,<sup>(14)</sup> T. Baumann,<sup>(6)</sup> F. Bedeschi,<sup>(13)</sup> S. Behrends,<sup>(2)</sup> S. Belforte,<sup>(13)</sup> G. Bellettini,<sup>(13)</sup> J. Bellinger,<sup>(21)</sup> D. Benjamin,<sup>(20)</sup> J. Bensinger,<sup>(2)</sup> A. Beretvas,<sup>(4)</sup> J. P. Berge,<sup>(4)</sup> S. Bertolucci,<sup>(5)</sup> S. Bhadra,<sup>(7)</sup> M. Binkley,<sup>(4)</sup> R. Blair,<sup>(1)</sup> C. Blocker,<sup>(2)</sup> K. Bloom,<sup>(3)</sup> V. Bolognesi,<sup>(13)</sup> A. W. Booth,<sup>(4)</sup> C. Boswell,<sup>(8)</sup> G. Brandenburg,<sup>(6)</sup> D. Brown,<sup>(6)</sup> E. Buckley-Geer,<sup>(17)</sup> H. S. Budd,<sup>(15)</sup> G. Busetto,<sup>(11)</sup> A. Byon-Wagner,<sup>(4)</sup> K. L. Byrum,<sup>(21)</sup> C. Campagnari,<sup>(3)</sup> M. Campbell,<sup>(3)</sup> A. Caner,<sup>(4)</sup> R. Carey,<sup>(6)</sup> W. Carithers,<sup>(10)</sup> D. Carlsmith,<sup>(21)</sup> J. T. Carroll,<sup>(4)</sup> R. Cashmore,<sup>(4),(a)</sup> A. Castro,<sup>(11)</sup> F. Cervelli,<sup>(13)</sup> K. Chadwick,<sup>(4)</sup> G. Chiarelli,<sup>(5)</sup> W. Chinowsky,<sup>(10)</sup> S. Cihangir,<sup>(4)</sup> A. G. Clark,<sup>(4)</sup> D. Connor,<sup>(12)</sup> M. Contreras,<sup>(2)</sup> J. Cooper,<sup>(4)</sup> M. Cordelli,<sup>(5)</sup> D. Crane,<sup>(4)</sup> M. Curatolo,<sup>(5)</sup> C. Day,<sup>(4)</sup> F. DeJongh,<sup>(4)</sup> S. Dell'Agnello,<sup>(13)</sup> M. Dell'Orso,<sup>(13)</sup> L. Demortier,<sup>(2)</sup> B. Denby,<sup>(4)</sup> P. F. Derwent,<sup>(3)</sup> T. Devlin,<sup>(17)</sup> D. DiBitonto,<sup>(18)</sup> M. Dickson,<sup>(15)</sup> R. B. Drucker,<sup>(10)</sup> K. Einsweiler,<sup>(10)</sup> A. Etchegoyen,<sup>(4),(a)</sup> J. E. Elias,<sup>(4)</sup> R. Ely,<sup>(10)</sup> S. Eno,<sup>(3)</sup> S. Errede,<sup>(7)</sup> B. Esposito,<sup>(5)</sup> B. Flaughner,<sup>(4)</sup> G. W. Foster,<sup>(4)</sup> M. Franklin,<sup>(6)</sup> J. Freeman,<sup>(4)</sup> H. Frisch,<sup>(3)</sup> T. Fuess,<sup>(4)</sup> Y. Fukui,<sup>(9)</sup> A. F. Garfinkel,<sup>(14)</sup> A. Gauthier,<sup>(7)</sup> S. Geer,<sup>(4)</sup> D. W. Gerdes,<sup>(3)</sup> P. Giannetti,<sup>(13)</sup> N. Giokaris,<sup>(16)</sup> P. Giromini,<sup>(5)</sup> L. Gladney,<sup>(12)</sup> M. Gold,<sup>(10)</sup> K. Goulios,<sup>(16)</sup> H. Grassmann,<sup>(11)</sup> C. Grosso-Pilcher,<sup>(3)</sup> C. Haber,<sup>(10)</sup> S. R. Hahn,<sup>(4)</sup> R. Handler,<sup>(21)</sup> K. Hara,<sup>(19)</sup> R. M. Harris,<sup>(4)</sup> J. Hauser,<sup>(4)</sup> C. Hawk,<sup>(17)</sup> T. Hessian,<sup>(18)</sup> R. Hollebeek,<sup>(12)</sup> L. Holloway,<sup>(7)</sup> P. Hu,<sup>(17)</sup> B. Hubbard,<sup>(10)</sup> B. T. Huffman,<sup>(14)</sup> R. Hughes,<sup>(12)</sup> P. Hurst,<sup>(5)</sup> J. Huth,<sup>(4)</sup> J. Hylen,<sup>(4)</sup> M. Incagli,<sup>(13)</sup> T. Ino,<sup>(19)</sup> H. Iso,<sup>(19)</sup> H. Jensen,<sup>(4)</sup> C. P. Jessop,<sup>(6)</sup> R. P. Johnson,<sup>(4)</sup> U. Joshi,<sup>(4)</sup> R. W. Kadel,<sup>(10)</sup> T. Kamon,<sup>(18)</sup> S. Kanda,<sup>(19)</sup> D. A. Kardelis,<sup>(7)</sup> I. Karliner,<sup>(7)</sup> E. Kearns,<sup>(6)</sup> L. Keeble,<sup>(18)</sup> R. Kephart,<sup>(4)</sup> P. Kesten,<sup>(2)</sup> R. M. Keup,<sup>(7)</sup> H. Keutelian,<sup>(4)</sup> D. Kim,<sup>(4)</sup> S. Kim,<sup>(19)</sup> L. Kirsch,<sup>(2)</sup> K. Kondo,<sup>(19)</sup> J. Konigsberg,<sup>(6)</sup> E. Kovacs,<sup>(4)</sup> S. E. Kuhlmann,<sup>(1)</sup> E. Kuns,<sup>(17)</sup> A. T. Laasanen,<sup>(14)</sup> J. I. Lamoureux,<sup>(21)</sup> S. Leone,<sup>(13)</sup> J. Lewis,<sup>(4)</sup> W. Li,<sup>(1)</sup> T. M. Liss,<sup>(7)</sup> P. Limon,<sup>(4)</sup> N. Lockyer,<sup>(12)</sup> C. B. Luchini,<sup>(7)</sup> P. Lukens,<sup>(4)</sup> P. Maas,<sup>(21)</sup> M. Mangano,<sup>(13)</sup> J. P. Marriner,<sup>(4)</sup> M. Mariotti,<sup>(13)</sup> R. Markeloff,<sup>(21)</sup> L. A. Markosky,<sup>(21)</sup> R. Mattingly,<sup>(2)</sup> P. McIntyre,<sup>(18)</sup> A. Menzione,<sup>(13)</sup> T. Meyer,<sup>(18)</sup> S. Mikamo,<sup>(9)</sup> M. Miller,<sup>(3)</sup> T. Mimashi,<sup>(19)</sup> S. Miscetti,<sup>(5)</sup> M. Mishina,<sup>(9)</sup> S. Miyashita,<sup>(19)</sup> Y. Morita,<sup>(19)</sup> S. Moulding,<sup>(2)</sup> J. Mueller,<sup>(17)</sup> A. Mukherjee,<sup>(4)</sup> L. F. Nakae,<sup>(2)</sup> I. Nakano,<sup>(19)</sup> C. Nelson,<sup>(4)</sup> C. Newman-Holmes,<sup>(4)</sup> J. S. T. Ng,<sup>(6)</sup> M. Ninomiya,<sup>(19)</sup> L. Nodulman,<sup>(1)</sup> S. Ogawa,<sup>(19)</sup> R. Paoletti,<sup>(13)</sup> V. Papadimitriou,<sup>(4)</sup> A. Para,<sup>(4)</sup> E. Pare,<sup>(6)</sup> S. Park,<sup>(4)</sup> J. Patrick,<sup>(4)</sup> T. J. Phillips,<sup>(6)</sup> R. Plunkett,<sup>(4)</sup> L. Pondrom,<sup>(21)</sup> J. Proudfoot,<sup>(1)</sup> F. Ptohos,<sup>(6)</sup> G. Punzi,<sup>(13)</sup> D. Quarrie,<sup>(4)</sup> K. Ragan,<sup>(12)</sup> G. Redlinger,<sup>(3)</sup> J. Rhoades,<sup>(21)</sup> M. Roach,<sup>(20)</sup> F. Rimondi,<sup>(4),(a)</sup> L. Ristori,<sup>(13)</sup> T. Rodrigo,<sup>(4)</sup> T. Rohaly,<sup>(12)</sup> A. Roodman,<sup>(3)</sup> W. K. Sakumoto,<sup>(15)</sup> A. Sansoni,<sup>(5)</sup> R. D. Sard,<sup>(7)</sup> A. Savoy-Navarro,<sup>(4)</sup> V. Scarpine,<sup>(7)</sup> P. Schlabach,<sup>(6)</sup> E. E. Schmidt,<sup>(4)</sup> O. Schneider,<sup>(10)</sup> M. H. Schub,<sup>(14)</sup> R. Schwitters,<sup>(6)</sup> A. Scribano,<sup>(13)</sup> S. Segler,<sup>(4)</sup> Y. Seiya,<sup>(19)</sup> M. Shapiro,<sup>(10)</sup> N. M. Shaw,<sup>(14)</sup> M. Sheaff,<sup>(21)</sup> M. Shochet,<sup>(3)</sup> J. Siegrist,<sup>(10)</sup> P. Sinervo,<sup>(12)</sup> J. Skarha,<sup>(8)</sup> K. Sliwa,<sup>(20)</sup> D. A. Smith,<sup>(13)</sup> F. D. Snider,<sup>(8)</sup> L. Song,<sup>(12)</sup> M. Spahn,<sup>(10)</sup> P. Sphicas,<sup>(4)</sup> R. St. Denis,<sup>(6)</sup> A. Stefanini,<sup>(13)</sup> G. Sullivan,<sup>(3)</sup> R. L. Swartz, Jr.,<sup>(7)</sup> M. Takano,<sup>(19)</sup> F. Tartarelli,<sup>(13)</sup> K. Takikawa,<sup>(19)</sup> S. Tarem,<sup>(2)</sup> D. Theriot,<sup>(4)</sup> M. Timko,<sup>(18)</sup> P. Tipton,<sup>(4)</sup> S. Tkaczyk,<sup>(4)</sup> A. Tollestrup,<sup>(4)</sup> J. Tonnison,<sup>(14)</sup> W. Trischuk,<sup>(6)</sup> N. Turini,<sup>(13)</sup> Y. Tsay,<sup>(3)</sup> F. Ukegawa,<sup>(19)</sup> D. Underwood,<sup>(1)</sup> S. Vejck, III,<sup>(8)</sup> R. Vidal,<sup>(4)</sup> R. G. Wagner,<sup>(1)</sup> R. L. Wagner,<sup>(4)</sup> N. Wainer,<sup>(4)</sup> J. Walsh,<sup>(12)</sup> T. Watts,<sup>(17)</sup> R. Webb,<sup>(18)</sup> C. Wendt,<sup>(21)</sup> H. Wenzel,<sup>(13)</sup> W. C. Wester, III,<sup>(10)</sup> T. Westhusing,<sup>(13)</sup> S. N. White,<sup>(16)</sup> A. B. Wicklund,<sup>(1)</sup> H. H. Williams,<sup>(12)</sup> B. L. Winer,<sup>(15)</sup> J. Wyss,<sup>(11)</sup> A. Yagil,<sup>(4)</sup> K. Yasuoka,<sup>(19)</sup> G. P. Yeh,<sup>(4)</sup> J. Yoh,<sup>(4)</sup> M. Yokoyama,<sup>(19)</sup> J. C. Yun,<sup>(4)</sup> A. Zanetti,<sup>(13)</sup> F. Zetti,<sup>(13)</sup> and S. Zucchelli<sup>(4),(a)</sup>

(CDF Collaboration)

<sup>(1)</sup>Argonne National Laboratory, Argonne, Illinois 60439

<sup>(2)</sup>Brandeis University, Waltham, Massachusetts 02254

<sup>(3)</sup>University of Chicago, Chicago, Illinois 60637

<sup>(4)</sup>Fermi National Accelerator Laboratory, Batavia, Illinois 60510

<sup>(5)</sup>Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, Frascati, Italy

<sup>(6)</sup>Harvard University, Cambridge, Massachusetts 02138

<sup>(7)</sup>University of Illinois, Urbana, Illinois 61801

<sup>(8)</sup>The Johns Hopkins University, Baltimore, Maryland 21218

<sup>(9)</sup>*National Laboratory for High Energy Physics (KEK), Tsukuba, Ibaraki, Japan*

<sup>(10)</sup>*Lawrence Berkeley Laboratory, Berkeley, California 94720*

<sup>(11)</sup>*Università di Padova, Istituto Nazionale di Fisica Nucleare, Sezione di Padova, I-35131 Padova, Italy*

<sup>(12)</sup>*University of Pennsylvania, Philadelphia, Pennsylvania 19104*

<sup>(13)</sup>*Istituto Nazionale di Fisica Nucleare, University and Scuola Normale Superiore of Pisa, I-56100 Pisa, Italy*

<sup>(14)</sup>*Purdue University, West Lafayette, Indiana 47907*

<sup>(15)</sup>*University of Rochester, Rochester, New York 14627*

<sup>(16)</sup>*Rockefeller University, New York, New York 10021*

<sup>(17)</sup>*Rutgers University, Piscataway, New Jersey 08854*

<sup>(18)</sup>*Texas A&M University, College Station, Texas 77843*

<sup>(19)</sup>*University of Tsukuba, Tsukuba, Ibaraki 305, Japan*

<sup>(20)</sup>*Tufts University, Medford, Massachusetts 02155*

<sup>(21)</sup>*University of Wisconsin, Madison, Wisconsin 53706*

(Received 26 December 1991)

We have observed over  $10^2$  events of the type  $W \rightarrow \tau \nu$  followed by  $\tau \rightarrow$  hadrons, where the taus are identified by their decay into one or three charged particles. We measure the cross section times branching ratio for  $p\bar{p} \rightarrow W \rightarrow \tau \nu$  and compare it to the value for  $W \rightarrow e \nu$  to directly measure the ratio of weak coupling constants  $g_\tau/g_e$ . We find  $g_\tau/g_e = 0.97 \pm 0.07$ , consistent with lepton universality.

PACS numbers: 13.38.+c, 13.10.+q, 13.85.Qk, 14.60.-z

The universality of lepton couplings to the vector bosons is a direct consequence of SU(2) gauge invariance in the standard model [1]. The magnitude of the  $W$ - $\tau$  coupling at low  $Q^2$  has been extensively studied in the decays of the tau; a recent compilation of results gives  $g_\tau/g_e = 0.967 \pm 0.017$  [2]. A previous direct measurement of  $g_\tau/g_e$  using  $W$  decays is consistent with universality [3]. Lepton universality in the neutral current channel has also been tested in decays of the  $Z$  [4].

In this paper we describe a measurement of the cross section times branching ratio,  $\sigma B$ , for production of  $W$  bosons and their decay  $W \rightarrow \tau \nu$ , which affords a test of lepton universality in the charged current channel according to the relation

$$(g_\tau/g_e)^2 = \sigma B(W \rightarrow \tau \nu) / \sigma B(W \rightarrow e \nu).$$

Unlike tests of universality derived from the tau lifetime, this constitutes a direct measurement of  $g_\tau/g_e$  which is free from the uncertainty related to the tau branching ratios. Our measurement is dependent upon the direct identification of tau leptons among the significant background present in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8$  TeV. In particular, the taus are positively identified by the observation of characteristic peaks in charged multiplicity at one and three.

This measurement used  $4.05 \text{ pb}^{-1}$  of  $p\bar{p}$  collisions at the Collider Detector at Fermilab (CDF), taken in the 1989 run of the Fermilab Tevatron. The CDF is described fully elsewhere [5]. The event selection relies on the characteristic features of the decay  $W \rightarrow \tau \nu$  and the subsequent hadronic decay of the energetic tau. These include a high transverse momentum track, in a narrow low multiplicity jet, in association with missing transverse energy  $E_T$  [6].

Two triggers [7] were used in the selection of  $W \rightarrow \tau \nu$  events. The first, a missing- $E_T$  trigger optimized to select

$W \rightarrow e \nu$  events, required  $E_T \geq 25$  GeV and electromagnetic  $E_T \geq 8$  GeV in the most energetic cluster. A second selection, the tau trigger [8], was implemented to lower the  $E_T$  threshold and to explicitly select  $W \rightarrow \tau \nu$  events. This trigger required a "tau trigger cluster" along with  $E_T \geq 20$  GeV. A tau trigger cluster required a charged track with transverse momentum  $P_T \geq 4.8$  GeV/c, matched in azimuth to a cluster in the calorimeter with transverse energy  $E_T \geq 10$  GeV, the ratio of energy in the hadronic to electromagnetic calorimeter  $\text{Had/EM} \geq 0.125$ , and the number of calorimeter towers  $N_{\text{towers}} \leq 2$ , where a tower subtends  $\Delta\phi$  (azimuthal angle) =  $15^\circ$  and  $\Delta\eta$  (pseudorapidity) =  $0.2$ . The tau trigger was in operation for one third of the data taking.

Tau decay reconstruction is designed to distinguish the hadronic final state of the  $W \rightarrow \tau \nu$  decay from the QCD jet background. The reconstruction algorithm begins by searching for a seed track in the central tracking chamber with  $P_T > 5$  GeV/c. Additional tracks with  $P_T \geq 1$  GeV/c are included in the cluster if they are within  $30^\circ$  of the seed track. The charged particle multiplicity of the tau is given by the number of tracks within  $10^\circ$  of the seed track ( $N_{\text{tracks}}$ ), where the choice of  $10^\circ$  gives excellent efficiency for tau decay products with  $P_T \geq 1$  GeV/c. The number of tracks between  $10^\circ$  and  $30^\circ$  from the seed track ( $N_{\text{isolation}}$ ) is used as a measure of the isolation of the tau cluster. The energy deposited in the tau calorimeter region, a rectangular region  $0.6$  in pseudorapidity ( $\eta$ ) by  $30^\circ$  in azimuthal angle ( $\phi$ ) centered on the seed track, is associated with the tau cluster. Electromagnetic showers, presumably from  $\pi^0$ 's, are identified by reconstructing any clusters in a wire and pad strip chamber at electromagnetic shower maximum which are in the tau's calorimeter region. The energy from electromagnetic showers is found using the energy in the calorimeter tower containing the strip chamber cluster, corrected for any charged tracks pointing at the same tower. Two

measures of the transverse energy are used: The  $E_T(\tau)$  is defined as the calorimeter transverse energy, while  $P_T(\tau)$  is defined as the scalar sum of the transverse momentum of charged particles in the  $10^\circ$  cone and the transverse energy of reconstructed electromagnetic showers in the tau's calorimeter region.

The  $W \rightarrow \tau\nu$  sample is selected by requiring  $E_T \geq 25$  GeV for events from the missing  $E_T$  trigger, and  $E_T \geq 20$  GeV for events from the tau trigger; for both there must be one tau cluster with  $E_T(\tau) \geq 15$  GeV and  $|\eta| \leq 1.1$  in the event. To reduce contamination from QCD backgrounds, the  $W \rightarrow \tau\nu$  events must also satisfy the following requirements on global event characteristics: no clusters [9] with  $E_T \geq 10$  GeV besides the tau cluster, and no clusters with  $E_T \geq 5$  GeV and  $\Delta\phi \geq 150^\circ$  from the tau cluster. After these cuts there remains a residual background of very  $\tau$ -like QCD events, which can be suppressed with the  $P_T(\tau)$  variable. Events from the tau trigger with one, two, or three tracks in the signal cone are required to have  $P_T(\tau) \geq 17.5, 20,$  or  $22.5$  GeV/c, respectively. Finally events where the tau cluster also satisfies a subset of the CDF electron selection cuts [10] are removed to eliminate contamination of the  $W \rightarrow \tau\nu$  sample by  $W \rightarrow e\nu$  decays.

The  $W \rightarrow \tau\nu$  sample is divided into signal and background regions. The signal region includes those clusters with  $N_{\text{track}} \leq 3$  and  $N_{\text{isolation}} = 0$ , while the background region includes those clusters with  $N_{\text{track}} \geq 4$  and any  $N_{\text{isolation}}$ , plus those clusters with  $N_{\text{track}} = 2$  and  $N_{\text{isolation}} \geq 1$ . There are 207 events in the signal region from the missing  $E_T$  trigger, and 77 from the tau trigger. Of these, 22 events are common to both samples. The sizes of the signal and background are listed in Table I. The charged multiplicity for those events with  $N_{\text{isolation}} = 0$  is shown in Fig. 1; direct evidence for the presence of taus is seen in the excess of events at one and three tracks, as expected for tau decays.

The primary background to the  $W \rightarrow \tau\nu$  signal comes from QCD jet events in which one parton fragments to satisfy the tau cluster requirement and the other partons are mismeasured, giving the event a substantial amount of  $E_T$ . The clusters in Fig. 1 with four or more tracks are due to these QCD backgrounds. The amount of background in the  $W \rightarrow \tau\nu$  signal region can be estimated by extrapolating from the number of events in the background region using the measured multiplicity distribution of the QCD background. We measure the multiplicity distribution of QCD background by applying the tau clustering algorithm to a large sample of QCD dijet events with  $E_T$  in the same range, from 15 to 55 GeV, as the  $W \rightarrow \tau\nu$  sample. From this sample, the probability for a QCD jet to satisfy the tau cluster requirements, that is the trigger cuts,  $E_T \geq 15$  GeV,  $N_{\text{track}} \leq 3$ ,  $N_{\text{isolation}} = 0$ , and the  $P_T(\tau)$  cuts, is 3.8% for the missing- $E_T$  sample and 0.66% for the tau trigger. The probability for a QCD jet to fake a tau cluster does not vary greatly with  $E_T$ . The expected number of background events is found

TABLE I. Summary of the  $W \rightarrow \tau\nu$  analysis.

	$E_T$ Sample	Tau Sample
Number of Events		
$W \rightarrow \tau\nu$ Data sample	207	77
QCD background	$63 \pm 3 \pm 8$	$26 \pm 2 \pm 4$
$Z \rightarrow \tau\tau$ background	$7 \pm 2$	$4 \pm 1$
$W \rightarrow e\nu$ background	$5 \pm 1$	-
$N_{W \rightarrow \tau\nu}$	$132 \pm 14 \pm 8$	$47 \pm 9 \pm 4$
$A \cdot \epsilon(W \rightarrow \tau\nu)$		
$A(W \rightarrow \tau\nu)$		.396
$\tau \rightarrow$ hadrons B.R.		.639
$\epsilon(W \rightarrow \tau\nu)$	.0636	.0679
$A \cdot \epsilon(W \rightarrow \tau\nu)$	.0161	.0172
Systematic Errors on $A \cdot \epsilon$		
M.C. statistics	$\pm .0005$	$\pm .0005$
$N_{\text{lower}}, P_T(\tau)$ cuts	-	$\pm .0008$
B.R. (correlated)	$\pm .0007$	$\pm .0004$
E-scale (correlated)	$\pm .0006$	$\pm .0012$
$\int L dt$		
Integrated Luminosity	$4.015 \text{pb}^{-1}$	$1.315 \text{pb}^{-1}$
$\sigma \cdot B$		
$\sigma \cdot B(W \rightarrow \tau\nu)$	$2.04 \pm .22 \pm .18 \text{nb}$	$2.08 \pm .40 \pm .26$
$\sigma \cdot B(W \rightarrow \tau\nu)$ combined	$2.05 \pm .27 \text{nb}$	
$\sigma \cdot B(W \rightarrow e\nu)$	$2.19 \pm .04 \text{ (stat)} \pm .11 \text{ (syst.) nb}$	
$g_\tau/g_e$		
$g_\tau/g_e$	$.97 \pm .07$	

by normalizing the QCD dijet sample to the  $W \rightarrow \tau\nu$  sample in the background multiplicity region, and extrapolating into the signal multiplicity region. The normalization is weighted to account for small differences in the  $E_T$  spectra between the two samples; this weighting




FIG. 1.  $N_{\text{track}}$  for tau clusters with  $N_{\text{isolation}} = 0$ , for the  $W \rightarrow \tau\nu$  data sample (squares) and the QCD background sample (triangles): (a) Missing- $E_T$  trigger sample; (b) tau trigger sample.

changes the number of background events by 10%.

The normalized multiplicity for clusters from the QCD dijet sample with  $N_{\text{isolation}}=0$  is also shown in Fig. 1. The number of background events in the signal region from QCD jets is  $63 \pm 3(\text{stat}) \pm 8(\text{syst})$  for the missing- $E_T$  trigger, and  $26 \pm 2(\text{stat}) \pm 4(\text{syst})$  for the tau trigger. The statistical error follows from the number of events used to calculate the background normalization factor. The systematic error is taken from the variation in the size of the background for various ranges of  $E_T$  in the QCD dijet events.

There are also backgrounds to the  $W \rightarrow \tau\nu$  sample from  $Z \rightarrow \tau\tau$  decays with one lost tau, and from  $W \rightarrow e\nu$  decays with a misidentified electron.  $Z \rightarrow \tau\tau$  events generated by ISAJET [11] and simulated with the CDF simulation indicate that there are  $7 \pm 2$  and  $4 \pm 1$   $Z \rightarrow \tau\tau$  events as background to the two  $W \rightarrow \tau\nu$  samples, respectively. From the known efficiencies of the electron selection cuts [10], we estimate  $5 \pm 1$   $W \rightarrow e\nu$  events are background to the missing- $E_T$  trigger sample only. After we subtract the background from the number of events in the signal region, there are  $132 \pm 14 \pm 8$  events from the missing- $E_T$  trigger, and  $47 \pm 9 \pm 4$  from the tau trigger.

The  $W \rightarrow \tau\nu$  detection efficiency is measured with the CDF  $W \rightarrow e\nu$  data [10]. The energy deposition and the track from the electron are removed and replaced by a simulated tau, which is allowed to decay using the branching ratios measured by the CELLO Collaboration [12]. The decay products are simulated using the CDF simulation, and the simulated data are merged with the remainder of the original  $W \rightarrow e\nu$  event. The use of real data ensures that the global characteristics of the event are modeled correctly. The simulation of the calorimeter response has been tuned to an accuracy of 5% with a sample of isolated charged tracks and with test-beam charged pions at 7 and 10 GeV [8].

The probability to detect  $W \rightarrow \tau\nu$  events can be factored into a part which includes those kinematic and geometrical requirements common to both the  $W \rightarrow \tau\nu$  and  $W \rightarrow e\nu$  analyses, and a part which applies to just the  $W \rightarrow \tau\nu$  sample. The relevant acceptance cuts applied to both  $W \rightarrow \tau\nu$  and  $W \rightarrow e\nu$  events are  $P_T(\text{lepton}) \geq 20$  GeV/c,  $|\eta|(\text{lepton}) \leq 1.0$ , and  $P_T(\nu) \geq 20$  GeV. The acceptance for these requirements has been calculated to be  $A=0.396 \pm 0.016$  [10]. The systematic error in this quantity, largely due to the  $P_T$  distribution of the  $W$  boson and the choice of structure functions, will cancel in the ratio  $\sigma B(W \rightarrow \tau\nu)/\sigma B(W \rightarrow e\nu)$ .

The second factor, the tau efficiency, includes all the selection cuts made to collect the  $W \rightarrow \tau\nu$  sample, after the above acceptance cuts have been applied. We find that 6.4% and 6.8% of events which pass the acceptance cuts, for the missing- $E_T$  and tau samples, respectively, pass the  $W \rightarrow \tau\nu$  selection cuts and have  $N_{\text{track}}$  and  $N_{\text{isolation}}$  in the signal region. Two small corrections have been included in the efficiency to account for the differing fiducial regions used in the  $W \rightarrow e\nu$  and  $W \rightarrow \tau\nu$  analyses

and the presence of background in the  $W \rightarrow e\nu$  based Monte Carlo simulation. The total acceptance for  $W \rightarrow \tau\nu$  events, including the branching ratio of the tau into hadrons, is  $A(W \rightarrow \tau\nu)=0.0161 \pm 0.0010$  for the missing- $E_T$  sample, and  $A(W \rightarrow \tau\nu)=0.0172 \pm 0.0016$  for the tau sample.

The factors involved in the calculation of the acceptance and its uncertainty are summarized in Table I. The statistical uncertainty comes from the 2664  $W \rightarrow e\nu$  events used as input for the Monte Carlo simulation. We take the uncertainty in the knowledge of the energy scale for hadrons in the calorimeter to be  $\pm 5\%$ , which leads to an uncertainty in the efficiency for  $W \rightarrow \tau\nu$  events to satisfy the  $E_T$  and  $E_T$  thresholds. There is also an uncertainty from the knowledge of the tau branching ratios into the various exclusive modes. The acceptance varies significantly by mode, with modes with greater numbers of pions, especially neutral pions, having higher acceptance. Uncertainties in the branching ratios are taken from Ref. [12], and are used to estimate the change in the efficiency. Finally the Monte Carlo simulation is used to assess the systematic uncertainty in the tau sample due to the efficiency of the trigger  $N_{\text{towers}}$  cut and the  $P_T(\tau)$  cut.

In Fig. 2 we compare the Monte Carlo prediction for the track multiplicity, assuming lepton universality, against the measured track multiplicity. The agreement between data and Monte Carlo simulation is excellent. Other kinematic variables such as  $E_T(\tau)$  and  $P_T(\tau)$  also show very good agreement between data and Monte Carlo simulation.

Combining the calculated acceptances, the number of




FIG. 2.  $N_{\text{track}}$  for tau clusters with  $N_{\text{isolation}}=0$ , for the  $W \rightarrow \tau\nu$  data sample with background subtracted (squares) and the Monte Carlo prediction (histogram): (a) Missing- $E_T$  trigger sample; (b) tau trigger sample.

$W \rightarrow \tau \nu$  events after a background subtraction, and the integrated luminosity we calculate  $\sigma_B(W \rightarrow \tau \nu)$ . To measure the ratio of the  $W$ - $\tau$  and  $W$ - $e$  couplings, the systematic uncertainty from the integrated luminosity and the acceptance has been omitted for both electron and tau  $\sigma_B$ , as it cancels in the ratio. We find  $\sigma_B = 2.04 \pm 0.22(\text{stat}) \pm 0.18(\text{syst})$  nb from the missing- $E_T$  sample, and  $\sigma_B = 2.08 \pm 0.40(\text{stat}) \pm 0.26(\text{syst})$  nb from the tau sample. The results from the two data samples are combined to yield a value of  $\sigma_B(W \rightarrow \tau \nu) = 2.05 \pm 0.27$  nb. The correlations in the statistical and systematic errors between the two samples have been included. When this is combined with the CDF measurement  $\sigma_B(W \rightarrow e \nu) = 2.19 \pm 0.04 \pm 0.11$  nb [10], the ratio of weak coupling constants is found to be  $g_\tau/g_e = 0.97 \pm 0.07$ , in agreement with the hypothesis of lepton universality.

We thank the Fermilab Accelerator Division and the staffs at our individual institutions for their exceptional performance. This work was supported by the DOE, the NSF, Istituto Nazionale di Fisica Nucleare, the Ministry of Science, Culture and Education of Japan, and the A. P. Sloan Foundation.

<sup>(a)</sup>Visitor.

- [1] S. L. Glashow, Nucl. Phys. **22**, 579 (1961); S. Weinberg, Phys. Rev. Lett. **19**, 1264 (1967); A. Salam, in *Elementary Particle Physics*, edited by N. Svarthom (Almqvist and Wiksells, Stockholm, 1968), p. 367; A. DeRujula, in

*Weak and Electromagnetic Interactions at High Energy*, Proceedings of the Les Houches Summer School, Session XXIX, edited by R. Balian and C. H. L. Smith (North-Holland, Amsterdam, 1976), p. 569.

- [2] Particle Data Group, J. J. Hernandez *et al.*, Phys. Lett. B **239**, 1 (1990).  
 [3] UA1 Collaboration, C. Albajar *et al.*, Z. Phys. C **44**, 15 (1989); UA2 Collaboration, J. Alitti *et al.*, CERN Technical Report No. CERN-PPE/91-69 (to be published).  
 [4] ALEPH, DELPHI, L3, and OPAL Collaborations, CERN Technical Report No. CERN-PPE/91-232 (to be published).  
 [5] CDF Collaboration, F. Abe *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **271**, 387 (1988).  
 [6] Missing transverse energy is defined as  $E_T \equiv ((\sum_i E_T(i) \times \sin\phi(i))^2 + [\sum_i E_T(i) \cos\phi(i)]^2)^{1/2}$  where the sum is over calorimeter towers.  
 [7] D. Amidei *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **269**, 51 (1988).  
 [8] For details see Aaron Roodman, Ph.D. thesis, University of Chicago, 1991 (unpublished).  
 [9] CDF Collaboration, F. Abe *et al.*, Phys. Rev. D **43**, 664 (1990).  
 [10] CDF Collaboration, F. Abe *et al.*, Phys. Rev. D **44**, 29 (1991).  
 [11] F. Paige and S. D. Protopopescu, ISAJET Monte Carlo version 6.21, Brookhaven National Laboratory Technical Report No. BNL 38034, 1986 (unpublished).  
 [12] We use the CELLO results because the sum of the exclusive tau branching ratios is 1.0 and their likelihood technique based on  $\pi^\pm$  and  $\pi^0$  content is well suited for our measurement. See CELLO Collaboration, H. J. Behrend *et al.*, Z. Phys. C **46**, 537 (1990).