

Are There Oscillations in the Baryon-to-Meson Ratio?

Jean-René Cudell^(a) and Keith R. Dienes^(b)

Department of Physics, McGill University, 3600 University Street, Montréal, Québec, Canada H3A-2T8

(Received 13 March 1992)

All available data indicate a surplus of baryon states over meson states for energies greater than about 1.5 GeV. Since hadron-scale string theory suggests that their numbers should become equal with increasing energy, it has recently been proposed that there must exist exotic mesons with masses just above 1.7 GeV in order to fill the deficit. We demonstrate that a stringlike picture is actually consistent with the present numbers of baryon and meson states, and in fact predicts regular oscillations in their ratio. This suggests a different role for new hadronic states.

PACS numbers: 12.40.Lk, 11.17.+y

In a recent work, Freund and Rosner [1] have examined the separate densities of observed meson and baryon states as functions of their masses. They find that the integrated number of baryon states is less than that of meson states for masses less than about 1.7 GeV, but then greatly surpasses the meson number at higher energies. Since hadron-scale string theories are successful in modeling not only the hadronic Regge trajectories but also the exponential (Hagedorn) growth [2] in the total hadronic density, Freund and Rosner point out that such theories may also serve as the basis for understanding the relation *between* the separate meson and baryon densities. This is possible in part due to a recent result of Kutasov and Seiberg [3], which states that the numbers of bosonic and fermionic states in a nonsupersymmetric tachyon-free string theory must approach each other as increasingly massive states are included. On the basis of this theoretical result, Freund and Rosner predict that there must exist a number of mesons yet to be discovered with masses above 1.7 GeV (in order to match the rise in baryon number); furthermore, since the presently observed baryon/meson ratio is consistent with quark-model calculations which include only conventional mesons and baryons [4] (i.e., states with $q\bar{q}$ and qqq quark configurations, respectively), they additionally speculate that these new mesons are likely to be exotic (with quark content $q^{p+1}\bar{q}^{p+1}$, $p \geq 1$). This then implies the existence of exotic baryons (with configurations $q^{p+3}\bar{q}^p$, $p \geq 1$), and one is led to imagine a tower of exotic hadronic states with higher and higher masses.

In this Letter we first present a more refined analysis of the existing data and then examine more precisely the role a hadron-scale string theory might play in predicting the densities of baryon and meson states. In particular, while the result of Kutasov and Seiberg can be expected to hold in the *asymptotic* region (mass $M \rightarrow \infty$), we find that for energies in the GeV range a naive hadron-scale string picture implies that the ratio between the numbers of baryon and meson states should in fact *oscillate* around unity, with mesons favored first, then baryons, then mesons again. The amplitude of this oscillation falls to zero as the mass increases (in accordance with the Kutasov-Seiberg result), but we find that for masses

below 2 GeV, the oscillation is still within its first cycle and can thus accommodate both the apparent surplus of lower-energy mesons as well as the surplus of higher-energy baryons. While there is therefore no apparent need for exotic mesons in the mass range Freund and Rosner had in mind ($1.7 \leq M \leq 2$ GeV), this oscillating ratio suggests an entirely different scenario for exotic hadrons: Each repeating cycle of the oscillation may correspond to the threshold for the next-order exotic mesons and baryons. Other scenarios (e.g., involving glueballs and hybrid quark-gluon states) are possible as well.

Let us now be more specific, and first outline some of the basic results of string theory (including that of Kutasov and Seiberg) which will be relevant for our discussion. Strings are one-dimensional extended objects whose different vibrational and rotational configurations correspond to different spacetime particles or states; in general the mass of such a state is given by

$$m = (n/\alpha')^{1/2}, \quad n \in \mathbf{Z}, \quad (1)$$

where α' is a constant characterizing the energy scale of the theory and where n is related to the number of vibrational mode excitations necessary for producing the state. Since the Lorentz spin J of such a state must satisfy $J \leq n + a_0$ where a_0 is a constant, we have the general result

$$J \leq \alpha' m^2 + a_0, \quad (2)$$

which identifies the constant α' as the traditional Regge slope. If the particular string theory contains both bosonic and fermionic states, we may denote their numbers at each level n as B_n and F_n , respectively; note that these are the numbers of *states* or field-theoretic degrees of freedom, and not the number of particles (e.g., spin or isospin multiplets). Another well-known prediction of string theory, then, is the asymptotic exponential growth of these numbers as functions of n :

$$B_n, F_n \sim a n^{-b} e^{c\sqrt{n}} \quad \text{as } n \rightarrow \infty, \quad (3)$$

where the positive constants a , b , and c are theory-specific parameters. Equations (2) and (3) apply in general to all string-type theories. More recently, however, Kutasov

and Seiberg have obtained a result [3] which applies to those string theories (or more generally, to those two-dimensional conformal field theories) which are free of physical tachyons and which have modular-invariant one-loop (toroidal) partition functions. Specifically, if we define $B(N) \equiv \sum_{n=0}^N B_n$ and $F(N) \equiv \sum_{n=0}^N F_n$, then Kutasov and Seiberg claim that

$$\lim_{N \rightarrow \infty} [B(N) - F(N)] = 0, \tag{4}$$

which in turn implies the weaker constraint

$$\lim_{N \rightarrow \infty} [F(N)/B(N)] = 1. \tag{5}$$

We shall require only this weaker form of the Kutasov-Seiberg result; indeed, the stronger version in Eq. (4) may not be entirely correct [5].

The extent to which such a string theory can be taken as a theory of hadrons is far from clear, and therefore in this Letter we shall confine ourselves to only those issues which follow from direct comparisons with the above generic results. Specifically, we shall assume [1] that one can model hadronic physics as a GeV-scale string theory giving rise to Eqs. (2), (3), and (5), with bosonic states identified as meson degrees of freedom and fermionic states as baryon degrees of freedom; furthermore, we shall consider only those generic aspects of string theory which affect the *relative* numbers of these states (i.e., their ratio) or their separate *patterns* of growth. Any other features, such as the specific absolute sizes of $B(N)$ and $F(N)$ or the mapping between particular string configurations and particular hadronic states, are likely to be highly model dependent.

We have computed the numbers and densities of experimentally observed meson and baryon states as functions of their masses. We have included those states containing only the three light quarks (u, d, s) both for reasons of experimental statistics [1] and more fundamentally because hadrons composed of heavy quarks do not lie on linear Regge trajectories as a string picture would dictate [Eq. (2)]. We differ from Ref. [1], however, in recognizing that although states in string theory are typically of zero width, most of the hadronic states or resonances are quite broad. Therefore, we have taken the hadronic density of states to be a sum of normalized Breit-Wigner distributions:

$$\frac{dN}{dm} = \frac{1}{2\pi} \sum_i W_i \frac{\Gamma_i}{(m - M_i)^2 + \Gamma_i^2/4}, \tag{6}$$

where M_i and Γ_i are respectively the masses and widths of the observed states [6], and where W_i are their multiplicities [i.e., the number of *states* per resonance, or $(2I+1)(2J+1)$ for a charge self-conjugate state of spin J and isospin I , and twice that otherwise]. In Fig. 1 we have plotted the total hadronic density of states as a function of m , and it is clear that this density experiences the exponential (Hagedorn-like) growth suggested in Eq. (3)




FIG. 1. Total density of observed hadronic states as function of mass, along with best fit to Hagedorn form of Ref. [2].

with Hagedorn temperature [2] $T_H \equiv [c(\alpha')^{1/2}]^{-1} \approx 250$ MeV, at least for masses up to 2 GeV. Barring unexpected physics, the failure of the curve in Fig. 1 to maintain this growth beyond 2 GeV is likely to be a reflection of current experimental limitations. Thus, we shall henceforth limit our attention to the experimental data below 2 GeV.

In Fig. 2 we have plotted the separate numbers (or integrated densities) of baryon and meson states with masses $m \leq M$ as functions of M . In order to facilitate a comparison with Eq. (5), we have also plotted their *ratio* as the shaded region in Fig. 3: This shaded region indicates the uncertainty in the ratio function due to the hadronic widths, with the upper border of the region corresponding to the Breit-Wigner densities in Eq. (6) and the lower border corresponding to the zero-width case. Ei-




FIG. 2. Total numbers of observed baryons (solid line) and mesons (dashed line) with masses $\leq M$, as functions of M .




FIG. 3. Shaded region: observed ratio of numbers of baryons and mesons, as discussed in text. Solid line: ratio function from a typical string model.

ther way, several features are immediately apparent, among them the pronounced surplus of mesons below 1.5 GeV and the pronounced surplus of baryons above this energy; indeed, this ratio shows no sign of a plateau near unity. This figure thus clearly indicates that it is hardly compelling to interpret this mass region as the region of onset of Kutasov-Seiberg asymptotic behavior. It is in fact straightforward to estimate the string level n in Eq. (1) to which a mass of 1.5 GeV corresponds: Taking the measured value of the hadronic Regge slope $\alpha' \approx 0.9$ (GeV) $^{-2}$, we obtain $n \approx 2$. Indeed, the entire region < 2 GeV corresponds only to string levels $n \leq 4$. Thus, even though these low-lying levels experience the asymptotic growth in Eq. (3), they clearly need not manifest the asymptotic behavior predicted in Eq. (5); indeed, the latter asymptotic behavior occurs only at higher energies.

Therefore, in order to determine the characteristics of the *approach* towards asymptotic behavior, we have calculated the ratio functions $R(N) = F(N)/B(N)$ predicted by a variety of different string theories (or string “models”) of the sort to which Eq. (5) should apply. While certain features of this function vary greatly and are highly model dependent, others—such as the exponential increase in the level degeneracies [Eq. (3)] or the existence of a Kutasov-Seiberg limit [Eq. (5)]—indeed appear to be generic. In particular, we find an important third universal feature [5]: As N increases, we find that the function $R(N)$ oscillates *around* unity, with the amplitude of this oscillation decreasing with increasing N . This “damped” oscillation, periodic in $n = \alpha' M^2$, is of course consistent with the Kutasov-Seiberg result in Eq. (5). Such an oscillation between bosonic and fermionic states is a consequence (and in fact the signature) of an underlying string symmetry known as modular invariance, and the wavelength λ of this oscillation is determined only by the energy scale of the theory [5], $\lambda = 4/\alpha'$.

The amplitude, on the other hand, is somewhat model dependent, and in fact vanishes in the case of supersymmetry: Indeed, the only way to break supersymmetry while preserving modular invariance is to do so in this regular oscillatory manner [5]. In Fig. 3 we have superimposed the results of a calculation based on a typical nonsupersymmetric string model, plotting $R(N)$ vs $M \equiv (N/\alpha')^{1/2}$.

In the mass range $M \leq 2$ GeV, the behavior of the string ratio in Fig. 3 is certainly consistent with the observed ratio: This oscillation typically begins with $R < 1$ (at $N=0$), first crosses $R=1$ at $N=2$ (corresponding to $M \approx 1.5$ GeV), and then increases beyond 1 as M approaches 2 GeV. Thus we see that the sign of the oscillation, as well as the position of the first node, are consistent with the data, and a surplus of mesons below 1.5 GeV as well as a surplus of baryons above 1.5 GeV are easily accommodated. Thus, on the basis of a comparison between these two figures in the $M \leq 2$ GeV range, we find that we need *not* claim a deficit of meson states with masses just above 1.5 GeV.

It will be interesting, however, to see whether the *entire* string-theoretic oscillation is ultimately realized at higher energies. While such an oscillation between bosonic and fermionic states has not been observed experimentally, we have seen in Fig. 1 that many hadronic states with energies above 2 GeV must be missing if Hagedorn-like growth is to be maintained in that region. That many such states are missing is also expected from an SU(3) picture as well as from conventional Regge-trajectory arguments. Such an oscillation, therefore, remains entirely possible.

It is important to bear in mind that we have focused on only the generic features predicted by a generic string-type theory, and one would need to further refine a particular string picture in order to expect a more quantitative agreement between the observed and predicted ratio functions. For example, the string theories we have examined here are intrinsically noninteracting: All of their states (or particles) have zero width, and can populate only the discrete energy levels indicated in Eq. (1). This is the origin of the sharp changes in the string ratio function in Fig. 3, and a more fully developed string theory incorporating particle interactions would undoubtedly yield a smoother, more continuous ratio function. Furthermore, dynamical considerations are also at the root of the relatively small size of the experimentally observed ratio function at masses $M \leq 1$ GeV: The lowest-lying mesons (i.e., the pions) have masses protected by a nearly unbroken chiral symmetry, while the masses of the lowest-lying baryons (i.e., the proton and neutron) are entirely unprotected and consequently much greater. This is in contrast to noninteracting string theories, which generally contain both bosons *and* fermions at the (exactly) massless level. A fully interacting string theory, therefore, should be expected to yield a closer agreement between the ratio functions, especially in the lower-mass

region. On the other hand, the *oscillations* in the ratio function are of a more universal nature, and although interactions can be expected to make them smooth, they should remain quite pronounced in the region $M < 4$ GeV where their amplitudes are large.

Given that string theories generically lead to such oscillations, and given that we cannot soon expect to observe *all* existing states in the several-GeV region, it is natural to try to predict how these oscillations might arise within the context of a more traditional quark-gluon picture. While the string theories themselves unambiguously predict which string vibrational-rotational configurations are ultimately responsible for producing these oscillations [5], one must specify or choose a particular mapping between these configurations and the various quark-gluon states in order to interpret these oscillations in terms of selected groups of baryons and mesons. The results are then highly model dependent. Therefore, rather than advocate a particular string-to-hadron mapping, we will simply propose two possible resulting scenarios which naturally extend the ideas of Ref. [1].

One natural scheme which might lead to such a regular, periodic meson-baryon oscillation involves exotic hadrons—i.e., mesons with quark structure $(q\bar{q})^{p+1}$ and baryons with quark structure $q^{p+3}\bar{q}^p$ for $p \geq 1$. The special cases with $p=0$ of course correspond to the ordinary mesons and baryons which respectively dominate the two halves of the first cycle of the oscillation. It is thus natural to speculate that such a repeating pattern of oscillations is the result of regularly spaced thresholds for the p th exotic hadrons, implying alternating mass regions in which either the p th exotic mesons or baryons dominate:

$$\begin{aligned} (q\bar{q})^{p+1} \text{ mesons: } & (p+1/4)\lambda \leq M^2 \leq (p+1/2)\lambda, \\ q^{p+3}\bar{q}^p \text{ baryons: } & (p+3/4)\lambda \leq M^2 \leq (p+1)\lambda, \end{aligned} \quad (7)$$

where $\lambda = 4/\alpha' \approx 4.4$ (GeV)². Such an ordering of thresholds is in fact consistent with alternative analyses [1,7]. Another scenario involves not only glueballs but hadron-gluon “hybrids,” for such states—if color neutral—are in principle also present in a quark-gluon theory. While glueballs are necessarily bosonic, hybrid states can contribute to both bosonic and fermionic degrees of freedom depending on their quark content. In this scenario, then, each subsequent cycle of our oscillation corresponds to the crossing of the threshold for the next-order hybrid hadrons (i.e., hadrons with one addi-

tional gluonic insertion), with the wavelength $\lambda = 4/\alpha'$ of our oscillation representing the mass shift resulting from such gluonic insertions. Thus, this picture too can naturally explain the regularity of the string-predicted oscillation. Note, however, that *any* such picture necessarily implies the existence of exponentially increasing numbers of fundamentally new hadronic states at each of the mass regions listed in Eq. (7)—starting with, in particular, several hundred between 2 and 2.3 GeV.

In summary, then, we find that a generic hadron-scale string theory is consistent with the observed ratio of baryon and meson states; in particular, agreement with string theory does not require the existence of “missing mesons” (ordinary or exotic) in the mass region just above 1.5 GeV. On the other hand, we find that string theory and modular invariance predict a fermion/boson ratio which *oscillates* around unity as the mass increases, with the amplitude of these oscillations steadily decreasing. Such a picture therefore lends itself to a variety of interpretations involving exotic and/or hybrid hadrons, with each cycle of this oscillation corresponding to the thresholds for the next-order mesons and baryons. It will be interesting to see whether such pictures can be realized in more traditional (e.g., statistical or potential) quark models as well.

We are pleased to thank our colleagues at McGill, especially C. S. Lam and B. Margolis, for many fruitful discussions. This work was supported in part by NSERC (Canada) and les fonds FCAR (Québec).

(a)Electronic address: cudell@hep.physics.mcgill.ca.

(b)Electronic address: dien@hep.physics.mcgill.ca.

- [1] P. G. O. Freund and J. L. Rosner, Phys. Rev. Lett. **68**, 765 (1992).
- [2] R. Hagedorn, Nuovo Cimento **56A**, 1027 (1968).
- [3] D. Kutasov and N. Seiberg, Nucl. Phys. **B358**, 600 (1991).
- [4] S. Godfrey and N. Isgur, Phys. Rev. D **32**, 189 (1985); S. Capstick and N. Isgur, Phys. Rev. D **34**, 2809 (1986).
- [5] K. R. Dienes, McGill University Report No. McGill/92-08 (to be published).
- [6] Particle Data Group, J. J. Hernández *et al.*, Phys. Lett. B **239**, 1 (1990).
- [7] J. R. Cudell, K. R. Dienes, and B. Margolis (to be published).