

numerical solutions in terms of the single parameter  $\lambda$ . For  $\lambda = -0.20$  they calculate  $\nu_R = 3.9$ ,  $\Gamma = 0.6$ , and  $a_{S0} = 2.0$ ; J. W. Moffat and B. H. Bransden (private communication).

<sup>10</sup>Y. Nambu, Phys. Rev. 106, 1366 (1957).

<sup>11</sup>R. M. Sternheimer and S. J. Lindenbaum, Phys.

Rev. 109, 1723 (1958); J. G. Rushbrooke and D. Radojčić, Phys. Rev. Letters 5, 567 (1960).

<sup>12</sup>M. Stearns and M. B. Stearns, Phys. Rev. 103, 1534 (1956).

<sup>13</sup>G. Goldhaber, S. Goldhaber, W. Lee, and A. Pais, Phys. Rev. 120, 300 (1960).

---

E R R A T A

---

EVIDENCE FOR A  $\pi$ - $\pi$  RESONANCE IN THE  $I=1$ ,  $J=1$  STATE. A. R. Erwin, R. March, W. D. Walker, and E. West [Phys. Rev. Letters 6, 628 (1961)].

Footnote 3 should contain an additional reference to G. F. Chew and S. Mandelstam, Phys. Rev. 119, 467 (1960). At the end of the text, there should be an additional footnote: <sup>8</sup>It is possible that the effects of large S-wave  $\pi$ - $\pi$  scattering have been observed by A. Abashian, N. E. Booth, and K. M. Crowe, Phys. Rev. Letters 5, 258 (1960).

PROTON-PROTON INTERACTION. H. Feshbach, E. Lomon, and A. Tubis [Phys. Rev. Letters 6, 635 (1961)].

The last two terms in the square bracket of Eq. (1) for  $V_4(r)$  should be

$$-\vec{\sigma}^1 \cdot \vec{\sigma}^2 R_2(\mu r) - S_{12} R_3(\mu r),$$

instead of the same terms with positive signs.

The first two terms in the first square bracket of the expression for  $R_1(x)$  just below Eq. (1) should read:

$$\left( \frac{12}{x^2} + \frac{23}{x^4} \right) K_1(2x) \text{ instead of } \frac{12}{x^2} + \frac{23}{x^4} K_1(2x).$$

The first line of p. 636 should have  $\xi = 0$  instead of  $\xi = 1$ , and the second line should have  $\xi = 1$  instead of  $\xi = 0$ .