

**Enhanced Leading Production of D^\pm and $D^{*\pm}$ in 250
GeV π^\pm -Nucleon Interactions
[Phys. Rev. Lett. 72, 812 (1994)]**

G. A. Alves, S. Amato, J. C. Anjos, J. A. Appel, J. Astorga, T. Bernard, S. B. Bracker, L. M. Cremaldi, C. L. Darling, R. L. Dixon, D. Errede, C. Gay, D. R. Green, R. Jedicke, P. E. Karchin, S. Kwan, L. H. Lueking, J. R. T. de Mello Neto, J. Metheny, R. H. Milburn, J. M. de Miranda, H. da Motta Filho, A. Napier, D. Passmore, A. Rafatian, A. C. dos Reis, W. R. Ross, A. F. S. Santoro, M. Sheaff, M. H. G. Souza, W. J. Spalding, C. Stoughton, M. E. Streetman, D. J. Summers, S. F. Takach, A. Wallace,
and Z. Wu

(Fermilab E769 Collaboration)

The fit parameters reported in the paragraph above Fig. 1 should read $\chi^2 = 20.2$, upper tail probability 0.3%, and 99.7% confidence level. In the next paragraph, the fit parameters for the combined data should read $\chi^2 = 51.2$ and C.L. = $(1-8.4) \times 10^{-9}$. In the paragraph under Fig. 2, the fit parameters should read $A = 0.059 \pm 0.030$ (also in the Fig. 2 caption) with $\chi^2 = 5.9$ and upper tail probability 75%.