

**Multiple  $Mn^{2+}$ -Spin-Flip Raman Scattering at High Fields via Magnetic Polaron States in Semimagnetic Quantum Wells**  
**[Phys. Rev. Lett. 74, 2567 (1995)]**

J. Stühler, G. Schaack, M. Dahl, A. Waag, G. Landwehr, K. V. Kavokin, and I. A. Merkulov

The chemical composition of our samples has been mistakenly typeset, Mg replaced by Mn, in these positions:

Abstract line 2 should read as follows: ... in a quantum well structure with semimagnetic  $Cd_{0.98}Mn_{0.02}Te$  wells and nonmagnetic  $Cd_{0.76}Mg_{0.24}Te$  barriers ...

Page 2567, column 2, line 19/20 should read as follows: semimagnetic wells of  $Cd_{0.98}Mn_{0.02}Te$  and nonmagnetic barriers of  $Cd_{0.76}Mg_{0.24}Te$  ...

Page 2567, column 2, line 25 should read as follows: (wells:  $Cd_{0.985}Mn_{0.015}Te$ , barriers:  $Cd_{0.62}Mg_{0.38}Te$ ) ...

In the caption of Fig. 1, the first sentence should read as follows: Raman spectra  $-z(\sigma, \sigma)z$  of a single quantum well (18 Å) of  $Cd_{0.76}Mg_{0.24}Te/Cd_{0.98}Mn_{0.02}Te$  for various magnetic fields ...

Also on page 2569, column 1, line 30, the formula for the exchange field should read as follows:  $\vec{B}_{\text{exch}}(\vec{R}_i) = (3g\mu_B)^{-1}\beta J|\Psi_{HH}(\vec{R}_i)|^2$ .