

Search for W Boson Pair Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV

S. Abachi,¹² B. Abbott,³³ M. Abolins,²³ B. S. Acharya,⁴⁰ I. Adam,¹⁰ D. L. Adams,³⁴ M. Adams,¹⁵ S. Ahn,¹² H. Aihara,²⁰ J. Alitti,³⁶ G. Álvarez,¹⁶ G. A. Alves,⁸ E. Amidi,²⁷ N. Amos,²² E. W. Anderson,¹⁷ S. H. Aronson,³ R. Astur,³⁸ R. E. Avery,²⁹ A. Baden,²¹ V. Balamurali,³⁰ J. Balderston,¹⁴ B. Baldin,¹² J. Bantly,⁴ J. F. Bartlett,¹² K. Bazizi,⁷ J. Bendich,²⁰ S. B. Beri,³¹ I. Bertram,³⁴ V. A. Bezzubov,³² P. C. Bhat,¹² V. Bhatnagar,³¹ M. Bhattacharjee,¹¹ A. Bischoff,⁷ N. Biswas,³⁰ G. Blazey,¹² S. Blessing,¹³ A. Boehnlein,¹² N. I. Bojko,³² F. Borchering,¹² J. Borders,³⁵ C. Boswell,⁷ A. Brandt,¹² R. Brock,²³ A. Bross,¹² D. Buchholz,²⁹ V. S. Burtovoi,³² J. M. Butler,¹² D. Casey,³⁵ H. Castilla-Valdez,⁹ D. Chakraborty,³⁸ S.-M. Chang,²⁷ S. V. Chekulaev,³² L.-P. Chen,²⁰ W. Chen,³⁸ L. Chevalier,³⁶ S. Chopra,³¹ B. C. Choudhary,⁷ J. H. Christenson,¹² M. Chung,¹⁵ D. Claes,³⁸ A. R. Clark,²⁰ W. G. Cobau,²¹ J. Cochran,⁷ W. E. Cooper,¹² C. Cretsinger,³⁵ D. Cullen-Vidal,⁴ M. Cummings,¹⁴ D. Cutts,⁴ O. I. Dahl,²⁰ K. De,⁴¹ M. Demarteau,¹² R. Demina,²⁷ K. Denisenko,¹² N. Denisenko,¹² D. Denisov,¹² S. P. Denisov,³² W. Dharmaratna,¹³ H. T. Diehl,¹² M. Diesburg,¹² G. Di Loreto,²³ R. Dixon,¹² P. Draper,⁴¹ J. Drinkard,⁶ Y. Ducros,³⁶ S. R. Dugad,⁴⁰ S. Durston-Johnson,³⁵ D. Edmunds,²³ A. O. Efimov,³² J. Ellison,⁷ V. D. Elvira,^{12,*} R. Engelmann,³⁸ S. Eno,²¹ G. Eppley,³⁴ P. Ermolov,²⁴ O. V. Eroshin,³² V. N. Evdokimov,³² S. Fahey,²³ T. Fahland,⁴ M. Fatyga,³ M. K. Fatyga,³⁵ J. Featherly,³ S. Feher,³⁸ D. Fein,² T. Ferbel,³⁵ G. Finocchiaro,³⁸ H. E. Fisk,¹² Yu. Fisyak,²⁴ E. Flattum,²³ G. E. Forden,² M. Fortner,²⁸ K. C. Frame,²³ P. Franzini,¹⁰ S. Fredriksen,³⁹ S. Fuess,¹² A. N. Galjaev,³² E. Gallas,⁴¹ C. S. Gao,^{12,†} S. Gao,^{12,†} T. L. Geld,²³ R. J. Genik II,²³ K. Genser,¹² C. E. Gerber,^{12,‡} B. Gibbard,³ V. Glebov,³⁵ S. Glenn,⁵ B. Gobbi,²⁹ M. Goforth,¹³ A. Goldschmidt,²⁰ B. Gomez,¹ P. I. Goncharov,³² H. Gordon,³ L. T. Goss,⁴² N. Graf,³ P. D. Grannis,³⁸ D. R. Green,¹² J. Green,²⁸ H. Greenlee,¹² G. Griffin,⁶ N. Grossman,¹² P. Grudberg,²⁰ S. Grünendahl,³⁵ J. A. Guida,³⁸ J. M. Guida,³ W. Guryan,³ S. N. Gurzhev,³² Y. E. Gutnikov,³² N. J. Hadley,²¹ H. Haggerty,¹² S. Hagopian,¹³ V. Hagopian,¹³ K. S. Hahn,³⁵ R. E. Hall,⁶ S. Hansen,¹² R. Hatcher,²³ J. M. Hauptman,¹⁷ D. Hedin,²⁸ A. P. Heinson,⁷ U. Heintz,¹² R. Hernández-Montoya,⁹ T. Heuring,¹³ R. Hirosky,¹³ J. D. Hobbs,¹² B. Hoeneisen,^{1,§} J. S. Hoftun,⁴ F. Hsieh,²² Ting Hu,³⁸ Tong Hu,¹⁶ T. Huehn,⁷ S. Igarashi,¹² A. S. Ito,¹² E. James,² J. Jaques,³⁰ S. A. Jerger,²³ J. Z.-Y. Jiang,³⁸ T. Joffe-Minor,²⁹ H. Johari,²⁷ K. Johns,² M. Johnson,¹² H. Johnstad,³⁹ A. Jonckheere,¹² M. Jones,¹⁴ H. Jöstlein,¹² S. Y. Jun,²⁹ C. K. Jung,³⁸ S. Kahn,³ J. S. Kang,¹⁸ R. Kehoe,³⁰ M. Kelly,³⁰ A. Kernan,⁷ L. Kerth,²⁰ C. L. Kim,¹⁸ S. K. Kim,³⁷ A. Klatchko,¹³ B. Klima,¹² B. I. Klochkov,³² C. Klopfenstein,³⁸ V. I. Klyukhin,³² V. I. Kochetkov,³² J. M. Kohli,³¹ D. Koltick,³³ A. V. Kostitskiy,³² J. Kotcher,³ J. Kourlas,²⁶ A. V. Kozelov,³² E. A. Kozlovski,³² M. R. Krishnaswamy,⁴⁰ S. Krzywdzinski,¹² S. Kunori,²¹ S. Lami,³⁸ G. Landsberg,³⁸ R. E. Lanou,⁴ J.-F. Lebrat,³⁶ A. Leflat,²⁴ H. Li,³⁸ J. Li,⁴¹ Y. K. Li,²⁹ Q. Z. Li-Demarteau,¹² J. G. R. Lima,⁸ D. Lincoln,²² S. L. Linn,¹³ J. Linnemann,²³ R. Lipton,¹² Y. C. Liu,²⁹ F. Lobkowicz,³⁵ S. C. Loken,²⁰ S. Lökös,³⁸ L. Lueking,¹² A. L. Lyon,²¹ A. K. A. Maciel,⁸ R. J. Madaras,²⁰ R. Madden,¹³ I. V. Mandrichenko,³² Ph. Mangeot,³⁶ S. Mani,⁵ B. Mansoulié,³⁶ H. S. Mao,^{12,†} S. Margulies,¹⁵ R. Markeloff,²⁸ L. Markosky,² T. Marshall,¹⁶ M. I. Martin,¹² M. Marx,³⁸ B. May,²⁹ A. A. Mayorov,³² R. McCarthy,³⁸ T. McKibben,¹⁵ J. McKinley,²³ H. L. Melanson,¹² J. R. T. de Mello Neto,⁸ K. W. Merritt,¹² H. Miettinen,³⁴ A. Milder,² C. Milner,³⁹ A. Mincer,²⁶ J. M. de Miranda,⁸ C. S. Mishra,¹² M. Mohammadi-Baarmand,³⁸ N. Mokhov,¹² N. K. Mondal,⁴⁰ H. E. Montgomery,¹² P. Mooney,¹ M. Mudan,²⁶ C. Murphy,¹⁶ C. T. Murphy,¹² F. Nang,⁴ M. Narain,¹² V. S. Narasimham,⁴⁰ A. Narayanan,² H. A. Neal,²² J. P. Negret,¹ E. Neis,²² P. Nemethy,²⁶ D. Nešić,⁴ D. Norman,⁴² L. Oesch,²² V. Oguri,⁸ E. Oltman,²⁰ N. Oshima,¹² D. Owen,²³ P. Padley,³⁴ M. Pang,¹⁷ A. Para,¹² C. H. Park,¹² Y. M. Park,¹⁹ R. Partridge,⁴ N. Parua,⁴⁰ M. Paterno,³⁵ J. Perkins,⁴¹ A. Peryshkin,¹² M. Peters,¹⁴ H. Piekarczyk,¹³ Y. Pischalnikov,³³ A. Pluquet,³⁶ V. M. Podstavkov,³² B. G. Pope,²³ H. B. Prosper,¹³ S. Protopopescu,³ D. Pušeljić,²⁰ J. Qian,²² P. Z. Quintas,¹² R. Raja,¹² S. Rajagopalan,³⁸ O. Ramirez,¹⁵ M. V. S. Rao,⁴⁰ P. A. Rapidis,¹² L. Rasmussen,³⁸ A. L. Read,¹² S. Reucroft,²⁷ M. Rijssenbeek,³⁸ T. Rockwell,²³ N. A. Roe,²⁰ J. M. R. Roldan,¹ P. Rubinov,³⁸ R. Ruchti,³⁰ S. Rusin,²⁴ J. Rutherford,² A. Santoro,⁸ L. Sawyer,⁴¹ R. D. Schamberger,³⁸ H. Schellman,²⁹ D. Schmid,³⁹ J. Sculli,²⁶ E. Shabalina,²⁴ C. Shaffer,¹³ H. C. Shankar,⁴⁰ R. K. Shivpuri,¹¹ M. Shupe,² J. B. Singh,³¹ V. Sirotenko,²⁸ W. Smart,¹² A. Smith,² R. P. Smith,¹² R. Snihur,²⁹ G. R. Snow,²⁵ S. Snyder,³⁸ J. Solomon,¹⁵ P. M. Sood,³¹ M. Sosebee,⁴¹ M. Souza,⁸ A. L. Spadafora,²⁰ R. W. Stephens,⁴¹ M. L. Stevenson,²⁰ D. Stewart,²² F. Stocker,³⁹ D. A. Stoianova,³² D. Stoker,⁶ K. Streets,²⁶ M. Strovink,²⁰ A. Taketani,¹² P. Tamburello,²¹ J. Tarazi,⁶ M. Tartaglia,¹² T. L. Taylor,²⁹ J. Teiger,³⁶ J. Thompson,²¹ T. G. Trippe,²⁰ P. M. Tuts,¹⁰ N. Varelas,²³ E. W. Varnes,²⁰ P. R. G. Virador,²⁰ D. Vititoe,² A. A. Volkov,³² A. P. Vorobiev,³² H. D. Wahl,¹³ J. Wang,^{12,†} L. Z. Wang,^{12,†} J. Warchol,³⁰ M. Wayne,³⁰ H. Weerts,²³ W. A. Wenzel,²⁰ A. White,⁴¹ J. T. White,⁴²

J. A. Wightman,¹⁷ J. Wilcox,²⁷ S. Willis,²⁸ S. J. Wimpenny,⁷ J. V. D. Wirjawan,⁴² Z. Wolf,³⁹ J. Womersley,¹² E. Won,³⁵ D. R. Wood,¹² H. Xu,⁴ R. Yamada,¹² P. Yamin,³ C. Yanagisawa,³⁸ J. Yang,²⁶ T. Yasuda,²⁷ C. Yoshikawa,¹⁴ S. Youssef,¹³ J. Yu,³⁵ Y. Yu,³⁷ Y. Zhang,^{12,†} Y. H. Zhou,^{12,†} Q. Zhu,²⁶ Y. S. Zhu,^{12,†} Z. H. Zhu,³⁵ D. Zieminska,¹⁶ A. Zieminski,¹⁶ A. Zinchenko,¹⁷ and A. Zylberstejn³⁶

(D0 Collaboration)

- ¹Universidad de los Andes, Bogota, Colombia
²University of Arizona, Tucson, Arizona 85721
³Brookhaven National Laboratory, Upton, New York 11973
⁴Brown University, Providence, Rhode Island 02912
⁵University of California, Davis, California 95616
⁶University of California, Irvine, California 92717
⁷University of California, Riverside, California 92521
⁸LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil
⁹Centro de Investigación y de Estudios Avanzados, Mexico City, Mexico
¹⁰Columbia University, New York, New York 10027
¹¹Delhi University, Delhi, India 110007
¹²Fermi National Accelerator Laboratory, Batavia, Illinois 60510
¹³Florida State University, Tallahassee, Florida 32306
¹⁴University of Hawaii, Honolulu, Hawaii 96822
¹⁵University of Illinois, Chicago, Illinois 60680
¹⁶Indiana University, Bloomington, Indiana 47405
¹⁷Iowa State University, Ames, Iowa 50011
¹⁸Korea University, Seoul, Korea
¹⁹Kyungsung University, Pusan, Korea
²⁰Lawrence Berkeley Laboratory, Berkeley, California 94720
²¹University of Maryland, College Park, Maryland 20742
²²University of Michigan, Ann Arbor, Michigan 48109
²³Michigan State University, East Lansing, Michigan 48824
²⁴Moscow State University, Moscow, Russia
²⁵University of Nebraska, Lincoln, Nebraska 68588
²⁶New York University, New York, New York 10003
²⁷Northeastern University, Boston, Massachusetts 02115
²⁸Northern Illinois University, DeKalb, Illinois 60115
²⁹Northwestern University, Evanston, Illinois 60208
³⁰University of Notre Dame, Notre Dame, Indiana 46556
³¹University of Panjab, Chandigarh 16-00-14, India
³²Institute for High Energy Physics, 142-284 Protvino, Russia
³³Purdue University, West Lafayette, Indiana 47907
³⁴Rice University, Houston, Texas 77251
³⁵University of Rochester, Rochester, New York 14627
³⁶Commissariat à l'Energie Atomique, Département d'Astrophysique, Physique des Particules, Physique Nucléaire et d'Instrumentation Associée/Service de Physique des Particules, CE-Saclay, France
³⁷Seoul National University, Seoul, Korea
³⁸State University of New York, Stony Brook, New York 11794
³⁹SSC Laboratory, Dallas, Texas 75237
⁴⁰Tata Institute of Fundamental Research, Colaba, Bombay 400005, India
⁴¹University of Texas, Arlington, Texas 76019
⁴²Texas A&M University, College Station, Texas 77843

(Received 10 March 1995)

The results of a search for W boson pair production in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV with subsequent decay to dilepton ($e\mu$, ee , and $\mu\mu$) channels are presented. One event is observed with an expected background of 0.56 ± 0.13 events with an integrated luminosity of approximately 14 pb^{-1} . Assuming equal strengths for the WWZ and $WW\gamma$ gauge boson coupling parameters κ and λ , limits on the CP -conserving anomalous coupling constants are $-2.6 < \Delta\kappa < 2.8$ and $-2.1 < \lambda < 2.1$ at the 95% confidence level.

PACS numbers: 14.70.Fm, 13.40.Em, 13.40.Gp, 13.85.Qk

The standard model (SM) of electroweak interactions makes precise predictions for the gauge boson self-couplings due to the non-Abelian gauge symmetry of $SU(2)_L \otimes U(1)_Y$. The $WW\gamma$ coupling has been studied using the cross section and photon transverse energy spectrum of $W\gamma$ events at UA2 [1], CDF [2], and D0 [3]. However, the WWZ trilinear coupling has not been previously tested. The W boson pair production process provides a direct test of both the $WW\gamma$ and WWZ couplings [4].

The leading-order SM diagrams for W boson pair production in $p\bar{p}$ collisions are u - and t -channel quark exchange as well as s -channel production with either a photon or a Z boson as the mediating particle. The latter process contains the $WW\gamma$ and WWZ trilinear couplings. The SM predicts that these couplings are $g_{WW\gamma} = -e$ and $g_{WWZ} = -e \cot\theta_W$ and that unitarity violation due to the u - and t -channel amplitudes (which depend on the well-known couplings between the W boson and quarks) is prevented by cancellations provided by the s -channel amplitudes. Thus, W boson pair production provides a test of the SM gauge structure.

A formalism has been developed to describe the $WW\gamma$ and WWZ interactions for models beyond the SM [5]. The general effective Lorentz invariant Lagrangian for the electroweak gauge couplings, after imposing C , P , and CP symmetry, contains six dimensionless coupling parameters: g_1^V , κ_V , and λ_V , where $V = \gamma$ or Z . g_1^Z is assumed to be equal to g_1^γ , which is restricted to unity by electromagnetic gauge invariance. The terms in the effective Lagrangian have a normalization which makes the $V = Z$ terms larger than the $V = \gamma$ terms by $\cot\theta_W$. The effective Lagrangian can be reduced to the SM Lagrangian by setting $\kappa_V = 1$ ($\Delta\kappa_V \equiv \kappa_V - 1 = 0$) and $\lambda_V = 0$. Throughout this Letter, it is assumed that $\kappa_\gamma = \kappa_Z$ and $\lambda_\gamma = \lambda_Z$. The coupling parameters are related to the magnetic dipole moments (μ_W) and electric quadrupole moments (Q_W^e) of the W boson: $\mu_W = (e/2M_W)(1 + \kappa + \lambda)$ and $Q_W^e = -(e/M_W^2)(\kappa - \lambda)$, where e and M_W are the charge and the mass of the W boson [6].

The effective Lagrangian leads to a W boson pair production cross section which grows with \hat{s} , the square of the invariant mass of the WW system, for non-SM values of the couplings. In order to avoid unitarity violation, the anomalous couplings are parametrized as form factors with a scale, Λ [e.g., $\Delta\kappa/(1 + \hat{s}/\Lambda^2)^2$]. By requiring that tree-level unitarity is satisfied, a constraint $\Lambda \leq \{6.88/[(\kappa - 1)^2 + 2\lambda^2]\}^{1/4}$ TeV is obtained [4]. Limits on the coupling parameters κ and λ are obtained by comparing the measured cross section for W boson pair production to the predicted non-SM values; the cross section increases with κ and λ above the SM prediction of 9.5 pb [7].

In this Letter the results of a search for $p\bar{p}(\sqrt{s} = 1.8 \text{ TeV}) \rightarrow WW + X \rightarrow l\bar{l}'\bar{\nu}\nu' + X$, where the leptons include muons and electrons, are presented. The data sam-

ple corresponds to an integrated luminosity of approximately 14 pb^{-1} collected with the D0 detector during the 1992–1993 Tevatron collider run at Fermilab.

The D0 detector [8] consists of three major components: the calorimeter, tracking, and muon systems. A hermetic, compensating, uranium-liquid argon sampling calorimeter with fine transverse and longitudinal segmentation in projective towers measures energy out to $|\eta| \sim 4.0$, where η is the pseudorapidity. The energy resolution for electrons and photons is $15\%/\sqrt{E(\text{GeV})}$. The resolution for the transverse component of missing energy, $\cancel{E}_T^{\text{cal}}$, is $1.1 \text{ GeV} + 0.02(\sum E_T)$, where $\sum E_T$ is the scalar sum of transverse energy, E_T , in GeV, deposited in the calorimeter. The central and forward drift chambers are used to identify charged tracks for $|\eta| \leq 3.2$. There is no central magnetic field. Muons are identified and their momentum measured with three layers of proportional drift tubes, one inside and two outside of the magnetized iron toroids, providing coverage for $|\eta| \leq 3.3$. The muon momentum resolution, determined from $J/\psi \rightarrow \mu\mu$ and $Z \rightarrow \mu\mu$ events, is $\sigma(1/p) = 0.18(p - 2)/p^2 \oplus 0.008$ (p in GeV/ c). The p_T of identified muons is used to correct $\cancel{E}_T^{\text{cal}}$ to form the missing transverse energy, \cancel{E}_T .

Muons are required to be isolated, to have energy deposition in the calorimeter corresponding to at least that of a minimum ionizing particle, and to have $|\eta| \leq 1.7$. For the $\mu\mu$ channel, cosmic rays are rejected by requiring that the muons have timing consistent with the beam crossing. Electrons are identified through the longitudinal and transverse shape of isolated energy clusters in the calorimeter and by the detection of a matching track in the drift chambers. Electrons are required to be within a fiducial region of $|\eta| \leq 2.5$. A criterion on ionization (dE/dx), measured in the drift chambers, is imposed to reduce backgrounds from photon conversions and hadronic showers with large electromagnetic content.

The event samples come from triggers with dilepton signatures. The $e\mu$ sample is selected from events passing the trigger requirement of an electromagnetic cluster with $E_T \geq 7 \text{ GeV}$ and a muon with $p_T \geq 5 \text{ GeV}/c$. The ee candidates are required to have two isolated electromagnetic clusters, each with $E_T \geq 10 \text{ GeV}$. The $\mu\mu$ candidates are selected from events where at least one muon is identified with $p_T \geq 5 \text{ GeV}/c$ at the trigger level.

In the offline selection for the $e\mu$ channel, a muon with $p_T \geq 15 \text{ GeV}/c$ and an electron with $E_T \geq 20 \text{ GeV}$ are required. Both \cancel{E}_T and $\cancel{E}_T^{\text{cal}}$ are required to be $\geq 20 \text{ GeV}$. In order to suppress $Z \rightarrow \tau\bar{\tau}$ and $b\bar{b}$ backgrounds, it is required that $20^\circ \leq \Delta\phi(p_T^\mu, \cancel{E}_T) \leq 160^\circ$ if $\cancel{E}_T \leq 50 \text{ GeV}$, where $\Delta\phi(p_T^\mu, \cancel{E}_T)$ is the angle in the transverse plane between the muon and \cancel{E}_T . One event survives these selection cuts in a data sample corresponding to an integrated luminosity of $13.9 \pm 0.8 \text{ pb}^{-1}$.

For the ee channel, two electrons are required, each with $E_T \geq 20 \text{ GeV}$. The \cancel{E}_T is required to be $\geq 20 \text{ GeV}$.

The Z boson background is reduced by removing events where the dielectron invariant mass is between 77 and 105 GeV/c^2 . It is required that $20^\circ \leq \Delta\phi(p_T^e, \cancel{E}_T) \leq 160^\circ$ for the lower energy electron if $\cancel{E}_T \leq 50$ GeV . This selection suppresses $Z \rightarrow ee$ as well as $\tau\tau$. The integrated luminosity in this channel is 14.3 ± 0.8 pb^{-1} . One event survives these selection requirements.

For the $\mu\mu$ channel, two muons are required, one with $p_T \geq 20$ GeV/c and another with $p_T \geq 15$ GeV/c . In order to remove Z boson events, it is required that the \cancel{E}_T projected on the line bisecting the muon tracks in the transverse plane be greater than 30 GeV . This selection requirement is less sensitive to the momentum resolution of the muons than is a dimuon invariant mass cut. It is required that $\Delta\phi(p_T^\mu, \cancel{E}_T) \leq 170^\circ$ for the higher p_T muon. No events survive these selection requirements in a data sample corresponding to an integrated luminosity of 12.2 ± 0.7 pb^{-1} .

Finally, in order to suppress background from $t\bar{t}$ production, the vector sum of the E_T from hadrons, \vec{E}_T^{had} , defined as $-(\vec{E}_T^{l1} + \vec{E}_T^{l2} + \vec{\cancel{E}}_T)$ is required to be less than 40 GeV in magnitude for all channels. Figure 1 shows a Monte Carlo simulation of E_T^{had} for ~ 20 fb^{-1} of SM WW and $t\bar{t}$ events. For WW events, nonzero values of E_T^{had} are due to gluon radiation and detector resolution. For $t\bar{t}$ events, the most significant contribution is the b -quark jets from the t -quark decays. This selection reduces the background from $t\bar{t}$ production by a factor of 4 for a t -quark mass of 160 GeV/c^2 and is slightly more effective for a more massive t quark. The efficiency of this selection criterion for SM W boson pair production events is $0.95_{-0.04}^{+0.01}$ and decreases slightly with increasing W boson pair invariant mass. The surviving ee candidate passes this selection requirement but the $e\mu$ candidate [9] is rejected.




FIG. 1. E_T^{had} for Monte Carlo WW and $t\bar{t}$ events with $M_{\text{top}} = 160$ GeV/c^2 ($\int L dt \sim 20$ fb^{-1}). Events with $E_T^{\text{had}} \geq 40$ GeV were rejected.

The detection efficiency for SM W boson pair production events is determined using the PYTHIA [10] event generator followed by a detailed GEANT [11] simulation of the D0 detector. Muon trigger and electron identification efficiencies are derived from the data. The overall detection efficiency for SM $WW \rightarrow e\mu$ is 0.092 ± 0.010 . For the ee channel the efficiency is 0.094 ± 0.008 . For the $\mu\mu$ channel it is 0.033 ± 0.003 . For the three channels combined, the expected number of events for SM W boson pair production, based on a cross section of 9.5 pb [7], is 0.47 ± 0.07 . The efficiency is larger for anomalous couplings because of the increase in the W boson P_T . The Monte Carlo program of Ref. [4] followed by a fast detector simulation [12] is used to estimate the detection efficiency for W boson pair production as a function of the coupling parameters λ and κ .

The backgrounds due to Z boson, Drell-Yan dilepton, $W\gamma$, and $t\bar{t}$ events are estimated using the PYTHIA and ISAJET [13] Monte Carlo event generators followed by the GEANT detector simulation. The backgrounds from $b\bar{b}$, $c\bar{c}$, multijet, and W + jet events, where a jet is misidentified as an electron, are estimated using the data. The $t\bar{t}$ cross section estimates are from calculations of Laenen, Smith, and van Neerven [14]. The $t\bar{t}$ background is averaged for $M_{\text{top}} = 160, 170, \text{ and } 180$ GeV/c^2 . The background estimates are summarized in Table I.

The 95% confidence level upper limit on the W boson pair production cross section is estimated based on one signal event including a subtraction of the expected background of 0.56 ± 0.13 events. The branching ratio $W \rightarrow l\bar{\nu} = 0.108 \pm 0.004$ [15] is assumed. Poisson-distributed numbers of events are convoluted with Gaussian uncertainties on the detection efficiencies, background and luminosity. For SM W boson pair production, the upper limit for the cross section is 87 pb at the 95% confidence level. From the observed limit, as a function of λ and κ , and the theoretical prediction of the W boson pair production cross section, the 95% confidence level limits on the coupling parameters shown in Fig. 2 (solid line) are obtained. Also shown in Fig. 2 (dotted line) is the contour of the unitarity constraint on the coupling

TABLE I. Summary of backgrounds to $WW \rightarrow ee$, $WW \rightarrow e\mu$, and $WW \rightarrow \mu\mu$. The units are the expected number of background events in the data sample. The uncertainties include both statistical and systematic contributions.

Background	$e\mu$	ee	$\mu\mu$
$Z \rightarrow ee$ or $\mu\mu$		0.02 ± 0.01	0.068 ± 0.026
$Z \rightarrow \tau\tau$	0.11 ± 0.05	$< 10^{-3}$	$< 10^{-3}$
Drell-Yan dileptons		$< 10^{-3}$	$< 10^{-3}$
$W\gamma$	0.04 ± 0.03	0.02 ± 0.01	
QCD	0.07 ± 0.07	0.15 ± 0.08	$< 10^{-3}$
$t\bar{t}$	0.04 ± 0.02	0.03 ± 0.01	0.009 ± 0.003
Total	0.26 ± 0.10	0.22 ± 0.08	0.077 ± 0.026




FIG. 2. 95% CL limits on the CP -conserving anomalous couplings λ and $\Delta\kappa$, assuming that $\lambda_\gamma = \lambda_Z$ and $\kappa_\gamma = \kappa_Z$. The dotted contour is the unitarity limit for the form factor scale $\Lambda = 900$ GeV which was used to set the coupling limits.

limits for the form factor scale $\Lambda = 900$ GeV. This value of Λ is chosen so that the observed coupling limits lie within this ellipse. Because of the $\cot\theta_W$ difference in the normalization of the anomalous couplings, the limits on λ and κ are particularly sensitive to the hitherto unmeasured κ_Z and λ_Z . The limits on the CP -conserving anomalous coupling parameters are $-2.6 < \Delta\kappa < 2.8$ ($\lambda = 0$) and $-2.1 < \lambda < 2.1$ ($\Delta\kappa = 0$).

The coupling limits are insensitive to the decrease in the expected $t\bar{t}$ background that would occur if the top quark is much more massive than $160\text{--}180$ GeV/ c^2 . If the top background is negligible, the 95% confidence level upper limit for SM W boson pair production is 89 pb.

In conclusion, a search for $WW \rightarrow$ dileptons in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV is made. In approximately 14 pb $^{-1}$ of data, one event is found with an expected background of 0.56 ± 0.13 events. From the standard model, 0.46 ± 0.08 events are expected. For the SM W boson pair production, the upper limit for the cross section is 87 pb at the 95% confidence level. The limits on the CP -conserving anomalous coupling parameters are $-2.6 < \Delta\kappa < 2.8$ ($\lambda = 0$) and $-2.1 < \lambda < 2.1$ ($\Delta\kappa = 0$) at the 95% confidence level where κ_γ and λ_γ are assumed to equal κ_Z and λ_Z , respectively. The limits on λ and $\Delta\kappa$ exhibit almost no correlation, in contrast to limits from Refs. [1–3]. The maximum form factor scale accessible for this experiment is $\Lambda = 900$ GeV.

We thank U. Baur for providing us with much helpful advice and D. Zeppenfeld for the WW Monte Carlo generator and useful instructions. We thank the Fermi-

lab Accelerator, Computing and Research Divisions, and the support staffs at the collaborating institutions for their contributions to the success of this experiment. We also acknowledge support provided by the U.S. Department of Energy, the U.S. National Science Foundation, the Commissariat à l’Energie Atomique in France, the Ministry for Atomic Energy and the Ministry of Science and Technology Policy in Russia, CNPq in Brazil, the Departments of Atomic Energy and Science and Education in India, Colciencias in Colombia, CONACyT in Mexico, and the Ministry of Education, Research Foundation and KOSEF in Korea.

*Visitor from CONICET, Argentina.

†Visitor from IHEP, Beijing, China.

‡Visitor from the Universidad de Buenos Aires, Argentina.

§Visitor from Univ. San Francisco de Quito, Ecuador.

- [1] UA2 Collaboration, J. Alitti *et al.*, Phys. Lett. B **277**, 194 (1992).
- [2] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **74**, 1936 (1995).
- [3] D0 Collaboration, S. Abachi *et al.*, “Measurement of the $WW\gamma$ gauge boson couplings in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV” (to be published).
- [4] K. Hagiwara, J. Woodside, and D. Zeppenfeld, Phys. Rev. D **41**, 2113 (1990).
- [5] K. Hagiwara, R. D. Peccei, D. Zeppenfeld, and K. Hikasa, Nucl. Phys. **B282**, 253 (1987).
- [6] K. Kim and Y-S. Tsai, Phys. Rev. D **7**, 3710 (1973).
- [7] J. Ohnemus, Phys. Rev. D **44**, 1403 (1991).
- [8] D0 Collaboration, S. Abachi *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **338**, 185 (1994).
- [9] D0 Collaboration, S. Abachi *et al.*, Phys. Rev. Lett. **72**, 2138 (1994). The kinematic properties of this $t\bar{t}$ candidate are discussed in detail here.
- [10] T. Sjöstrand, “PYTHIA 5.6 and Jetset 7.3 Physics and Manual,” Report No. CERN-TH.6488/92, 1992 (unpublished).
- [11] F. Carminati *et al.*, “GEANT Users Guide,” CERN Program Library, 1991 (unpublished).
- [12] H. Johari, Ph.D. thesis, Northeastern University, 1995 (unpublished).
- [13] F. Paige and S. Protopopescu, BNL Report No. BNL-38034, 1986 (unpublished), release V6.49.
- [14] E. Laenen, J. Smith, and W. L. van Neerven, Phys. Lett. B **321**, 254 (1994). For the $t\bar{t}$ background, the central value estimate of the cross section is used.
- [15] Particle Data Group, L. Montanet *et al.*, Phys. Rev. D **50**, 1173 (1994). The weighted average of the $W \rightarrow e\nu$ and $W \rightarrow \mu\nu$ branching fraction data is used.