

Novel Universal Correlations in Invariant Random-Matrix Models

E. Kanzieper and V. Freilikher

The Jack and Pearl Resnick Institute of Advanced Technology, Department of Physics, Bar-Ilan University, Ramat-Gan 52900, Israel

(Received 16 December 1996)

We show that eigenvalue correlations in unitary-invariant ensembles of large random matrices satisfy novel universal laws that only depend on a multicriticality of the bulk density of states near the soft edge of the spectrum. Our consideration is based on the previously unknown observation that the genuine density of states and the n -point correlation function are completely determined by the Dyson's density analytically continued onto the entire real axis. [S0031-9007(97)03184-0]

PACS numbers: 05.45.+b, 02.10.Sp, 05.40.+j

Random matrices have been introduced in a physical context since the works by Wigner [1] and Dyson [2]. Initially proposed as an effective phenomenological model for description of the higher excitations in nuclei [3], the invariant ensembles of large random matrices found numerous applications in very diverse fields of physics such as two-dimensional quantum gravity [4], quantum chromodynamics [5], quantum chaos [6], and mesoscopic physics [7]. Apparently, this ubiquity owes its origin to the very idea of the construction of the invariant one-matrix model [8], which only reflects the fundamental symmetry (orthogonal, unitary, or symplectic) of the underlying physical system/phenomenon but discards its (irrelevant) microscopic details. Since the symmetry constraints follow from first principles, even a rather crude matrix model allows identification of universal features which persist for a variety of systems with the same symmetry. This circumstance emphasizes the importance of the study of universality intrinsic to random matrices.

The simplest invariant random-matrix model is defined by the probability density

$$P[\mathbf{H}] = \frac{1}{Z_N} \exp\{-\beta \text{Tr} V[\mathbf{H}]\} \quad (1)$$

of the entries H_{ij} of the $N \times N$ random matrix \mathbf{H} , where the function $V[\mathbf{H}]$ referred to as the "confinement potential" must ensure existence of the partition function

Z_N , $N \gg 1$. The symmetry parameter β is equal to the number of independent elements of the nondiagonal entries of matrix \mathbf{H} . In the following we restrict our consideration to the unitary invariant, $U(N)$, matrix model with $\beta = 2$.

Nowadays it is widely believed that $U(N)$ invariant ensembles of large random matrices with rather strong level confinement may exhibit *three* different types of locally universal eigenlevel correlations which are characterized by the appropriately scaled two-point kernels.

(i) *Bulk scaling limit* is associated with a spectrum range where the confinement potential is well behaved, and density of levels can approximately be taken as a constant. It has been proven in Refs. [9–11] that for rather strong confinement potentials [12] the two-point kernel follows the *universal sine law*

$$K_{\text{bulk}}(s, s') = \frac{\sin[\pi(s - s')]}{\pi(s - s')}. \quad (2)$$

Here the scaling variable s is measured in the units of the mean level spacing: $s = \varepsilon/\Delta_N$.

(ii) *Origin scaling limit* deals with that part of the spectrum where the confinement potential displays a logarithmic singularity: $V(\varepsilon) \rightarrow V(\varepsilon) - \alpha \ln |\varepsilon|$. In the vicinity of the singularity $\varepsilon = 0$, the two-point kernel satisfies the *universal Bessel law* [13],

$$K_{\text{orig}}(s, s') = \frac{\pi}{2} \sqrt{ss'} \frac{J_{\alpha+1/2}(\pi s) J_{\alpha-1/2}(\pi s') - J_{\alpha-1/2}(\pi s) J_{\alpha+1/2}(\pi s')}{s - s'}. \quad (3)$$

Here s is scaled by the level spacing near the origin, $s = \varepsilon/\Delta_N(0)$.

(iii) *Soft-edge scaling limit*, relevant to the tail of eigenvalue support where crossover occurs from a nonzero density of states to a vanishing one [14], has been only investigated for Gaussian unitary ensemble [15], and quite recently for $U(N)$ invariant ensembles of large random matrices associated with quartic and sextic confinement potentials [16]. It has been found that in the soft-edge scaling limit, for all these ensembles the two-point ker-

nels follow the *Airy law*

$$K_{\text{soft}}(s, s') = \frac{\text{Ai}(s)\text{Ai}'(s') - \text{Ai}(s')\text{Ai}'(s)}{s - s'}. \quad (4)$$

Here $s \propto N^{2/3}(\varepsilon/D_N - 1)$ with D_N being the end point of the spectrum. Whereas universality in the spectrum bulk and near its origin has rigorously been proven for a wide class of strong symmetric confinement potentials, the supposed universality of the Airy kernel has not been proven.

Our aim here is to demonstrate that the problem of eigenvalue correlations in the random-matrix ensembles with nonsingular confinement potentials can be treated in a unified way by means of the new universal second-order differential equation for the wave functions of fictitious noninteracting fermions naturally appearing in

the method of orthogonal polynomials [8]. This allows us to prove that the Airy correlations, Eq. (4), being universal for a wide class of matrix models Eq. (1), are indeed a particular case of more general novel universal correlations which are represented by the scaled m th multicritical two-point kernel

$$K_{\text{soft}}^{(m)}(s, s') = \frac{G(s|\nu^*)G'(s'|\nu^*)s^{3/2-\nu^*} - G(s'|\nu^*)G'(s|\nu^*)(s')^{3/2-\nu^*}}{s - s'}, \quad (5)$$

where the function G is expressed through the Bessel functions as

$$G(s|\nu^*) = \frac{1}{2\sqrt{\nu^*}} \left[\sin\left(\frac{\pi}{4\nu^*}\right) + (-1)^{\nu^*-3/2} \right]^{-1/2} \times \begin{cases} s^{(1/2)(\nu^*-1/2)} [I_{(-1/2)(1-1/2\nu^*)}(\frac{s\nu^*}{\nu^*}) - I_{(1/2)(1-1/2\nu^*)}(\frac{s\nu^*}{\nu^*})], & s > 0, \\ |s|^{(1/2)(\nu^*-1/2)} [J_{(-1/2)(1-1/2\nu^*)}(\frac{|s|\nu^*}{\nu^*}) + (-1)^{\nu^*-3/2} J_{(1/2)(1-1/2\nu^*)}(\frac{|s|\nu^*}{\nu^*})], & s < 0, \end{cases} \quad (6)$$

and parameter ν^* is determined by the even critical index $m = 0, 2, 4, \text{etc.}$, of the matrix model,

$$\nu^* = m + \frac{3}{2}. \quad (7)$$

Note that the critical index m is completely determined by the type of singularity of the Dyson's density of states near the soft edge [14]: $\nu_D(\varepsilon) \propto (1 - \varepsilon^2/D_N^2)^{m+1/2}$.

Equations (5)–(7) together with Eqs. (22) and (23) below are the main results of the paper. Although we concentrate our attention on the problem of eigenvalue correlations near the soft edge, the treatment we present here is quite general being relevant to an arbitrary spectrum range.

Within the orthogonal polynomial technique the two-point kernel $K_N(\varepsilon, \varepsilon')$ determining the n -point correlation function R_n for eigenvalue spectrum of large random matrices, $R_n(\varepsilon_1, \dots, \varepsilon_n) = \det[K_N(\varepsilon_i, \varepsilon_j)]_{i,j=1\dots n}$, can be written through the fictitious “wave functions” $\psi_n(\varepsilon)$ as

$$K_N(\varepsilon, \varepsilon') = c_N \frac{\psi_N(\varepsilon')\psi_{N-1}(\varepsilon) - \psi_N(\varepsilon)\psi_{N-1}(\varepsilon')}{\varepsilon' - \varepsilon}. \quad (8)$$

Here c_N is the recurrence coefficient entering the three-term recurrence equation

$$\varepsilon P_{n-1} = c_n P_n + c_{n-1} P_{n-2} \quad (9)$$

for polynomials P_n orthogonal on the entire real axis,

$$\int d\alpha(\varepsilon) P_n(\varepsilon) P_m(\varepsilon) = \delta_{nm}, \quad (10)$$

and the wave function $\psi_n(\varepsilon) = P_n(\varepsilon) \exp\{-V(\varepsilon)\}$. The measure $d\alpha(\varepsilon) = \exp\{-2V(\varepsilon)\}d\varepsilon$ is completely determined by symmetric confinement potential

$$V(\varepsilon) = \sum_{k=1}^p \frac{d_k}{2k} \varepsilon^{2k} \quad (11)$$

with $d_p > 0$. The signs of the rest d_k 's can be arbitrary but they should lead to an eigenvalue density supported on a single connected interval $(-D_N, +D_N)$.

To study the eigenvalue correlations in the random-matrix ensemble with confinement potential Eq. (11) we

note that a three-term recurrence equation for orthogonal polynomials $P_n(\varepsilon)$ can be mapped onto a second-order differential equation for these orthogonal polynomials and/or corresponding wave functions $\psi_n(\varepsilon)$. This was already observed for the first time by Shohat in 1930 [17]. Considerably later Shohat's idea was developed by Bonan and Clark [18]. The simple and elegant method proposed in Refs. [17,18] turns out to be a very general and powerful one for the analysis of spectral properties possessed by large random matrices.

To map Eq. (9) onto a second-order differential equation for ψ_n , we note the following identity:

$$\frac{dP_n}{d\varepsilon} = A_n(\varepsilon)P_{n-1} - B_n(\varepsilon)P_n, \quad (12)$$

where the functions $A_n(\varepsilon)$ and $B_n(\varepsilon)$ can be found from considerations below. Since $dP_n/d\varepsilon$ is a polynomial of the degree $n - 1$, it can be represented [19] through the Fourier expansion in terms of the kernel $Q_n(t, \varepsilon) = \sum_{k=0}^{n-1} P_k(t)P_k(\varepsilon)$ as follows:

$$\frac{dP_n}{d\varepsilon} = \int d\alpha(t) \frac{dP_n}{dt} Q_n(t, \varepsilon). \quad (13)$$

Integrating by parts we obtain that

$$\frac{dP_n}{d\varepsilon} = 2 \int d\alpha(t) Q_n(t, \varepsilon) \left(\frac{dV}{dt} - \frac{dV}{d\varepsilon} \right) P_n(t). \quad (14)$$

Now, making use of the Christoffel-Darboux theorem [19], we conclude that unknown functions A_n and B_n in Eq. (12) are

$$A_n(\varepsilon) = 2c_n \int d\alpha(t) \frac{V'(t) - V'(\varepsilon)}{t - \varepsilon} P_n^2(t), \quad (15)$$

$$B_n(\varepsilon) = 2c_n \int d\alpha(t) \frac{V'(t) - V'(\varepsilon)}{t - \varepsilon} P_n(t)P_{n-1}(t). \quad (16)$$

At first glance representations Eqs. (15) and (16) are rather useless because they involve the same orthogonal polynomials which enter Eq. (12). Nevertheless these

expressions do allow us to get the functions A_n and B_n in closed forms directly related to the confinement potential and to the end point of eigenvalue spectrum. Restricting our following considerations to large indices $n = N \gg 1$, we reduce Eq. (9) to the asymptotic form $\varepsilon P_N = c_N(P_{N+1} + P_{N-1})$, whence

$$\varepsilon^\lambda P_N = \sum_{j=0}^{\lambda} \binom{\lambda}{j} c_N^\lambda P_{N+2j-\lambda}, \quad \lambda \geq 0. \quad (17)$$

Substituting $V(\varepsilon)$ given by Eq. (11) into Eq. (15) yields

$$A_N(\varepsilon) = 2c_N \sum_{k=1}^p \sum_{\lambda=1}^{2k-1} d_k \varepsilon^{\lambda-1} \int d\alpha(t) P_N^2(t) t^{2k-\lambda-1}. \quad (18)$$

Then, taking into account Eq. (17) as well as the orthogonality of P_n , we arrive at the expression for $A_N(\varepsilon)$ defined for arbitrary ε ,

$$A_N(\varepsilon) = \frac{\pi \nu_D(\varepsilon)}{\sqrt{1 - \varepsilon^2/D_N^2}}, \quad (19)$$

where

$$\nu_D(\varepsilon) = \frac{2}{\pi^2} \mathcal{P} \int_0^{D_N} \frac{t dt}{t^2 - \varepsilon^2} \frac{dV}{dt} \sqrt{\frac{1 - \varepsilon^2/D_N^2}{1 - t^2/D_N^2}} \quad (20)$$

is the *Dyson's density continued onto the entire real axis*, so that ε can lie both inside and outside of an eigenvalue support. The spectrum end point $D_N = 2c_N$ is the positive root of the integral equation

$$N = \frac{2}{\pi} \int_0^{D_N} \frac{dV}{dt} \frac{t dt}{\sqrt{D_N^2 - t^2}}, \quad (21)$$

following from normalization of Dyson's density.

Combining Eqs. (12) and (19), and using asymptotic identity $B_N = \varepsilon A_N/D_N - dV/d\varepsilon$, which is a consequence of Eqs. (15) and (16), it is a straightforward step to obtain the following remarkable asymptotic differential equation:

$$\psi_N'' - \left[\frac{d}{d\varepsilon} \ln \left(\frac{\pi \nu_D(\varepsilon)}{\sqrt{1 - \varepsilon^2/D_N^2}} \right) \right] \psi_N' + \pi^2 \nu_D^2(\varepsilon) \psi_N = 0, \quad (22)$$

which together with relationship

$$\psi_N' = \frac{\pi \nu_D(\varepsilon)}{\sqrt{1 - \varepsilon^2/D_N^2}} \left(\psi_{N-1} - \frac{\varepsilon}{D_N} \psi_N \right) \quad (23)$$

provides a general basis for the study of eigenvalue correlations in an *arbitrary spectral range*.

An interesting property of these equations is that they do not contain the confinement potential explicitly, but only involve the Dyson's density ν_D and spectrum end point D_N . Moreover, it turns out that the *knowledge of Dyson's density* (that coincides with the real density of states only in the spectrum bulk) *is sufficient to determine the genuine density of states, as well as the n -*

point correlation function, everywhere. We also note that Eq. (22) can be derived in a different way for monotonic confinement potentials increasing at least as fast as $|\varepsilon|$ at infinity. This suggests that differential equation Eq. (22) should hold generally and not only for confinement potentials having the polynomial form Eq. (11).

Up to this point our derivation was quite general without any respect to the soft edge of eigenvalue support. We now focus our attention on the eigenvalue correlations near the soft edge $\varepsilon = D_N$. It is known [14] that by tuning coefficients d_k which enter V , one can obtain a bulk (Dyson's) density of states which possesses a singularity of the type

$$\nu_D(\varepsilon) = \left(1 - \frac{\varepsilon^2}{D_N^2} \right)^{m+1/2} \mathcal{R}_N \left(\frac{\varepsilon}{D_N} \right), \quad (24)$$

with $m = 0, 2, 4$, etc., and \mathcal{R}_N being a well-behaved function with $\mathcal{R}_N(1) \neq 0$. [Odd indices m are inconsistent with our choice that the leading coefficient d_p , entering confinement potential $V(\varepsilon)$, be positive in order to keep a convergence of integral for partition function Z_N in Eq. (1).] Such an m th multicriticality can be achieved by many means, and the corresponding plethora of multicritical potentials $V^{(m)}$ is given by the equation

$$\frac{dV^{(m)}(\varepsilon)}{d\varepsilon} = \mathcal{P} \int_{-D_N}^{+D_N} \frac{dt}{\varepsilon - t} \left(1 - \frac{t^2}{D_N^2} \right)^{m+1/2} \times \mathcal{R}_N \left(\frac{t}{D_N} \right). \quad (25)$$

So-called minimal multicritical potentials which correspond to $\mathcal{R}_N = \text{const}$ can be found in Refs. [4,14].

Below we intend to demonstrate that as long as multicriticality of order m is reached, the eigenvalue correlations in the vicinity of the soft edge become universal, and are independent of the particular potential chosen. The order m of the multicriticality is the only parameter which governs spectral correlations in the soft-edge scaling limit.

Let us move the spectrum origin to its end point D_N , making the replacement

$$\varepsilon_s = D_N \left[1 + \frac{1}{2} s \left(\frac{2}{\pi D_N \mathcal{R}_N(1)} \right)^{1/\nu^*} \right], \quad (26)$$

that defines the m th *soft-edge scaling limit* provided $s \ll [D_N \mathcal{R}_N(1)]^{1/\nu^*} \propto N^{1/\nu^*}$. It is straightforward to show from Eqs. (22) and (23) that the function $\hat{\psi}_N(s) = \psi_N(\varepsilon_s - D_N)$ obeys the differential equation

$$\hat{\psi}_N''(s) - \frac{(\nu^* - 3/2)}{s} \hat{\psi}_N'(s) - s^{2(\nu^*-1)} \hat{\psi}_N(s) = 0, \quad (27)$$

and that the following relation takes place:

$$\hat{\psi}_{N-1}(s) = \hat{\psi}_N(s) + (-1)^{\nu^*-3/2} \left(\frac{2}{\pi D_N \mathcal{R}_N(1)} \right)^{1/2\nu^*} \times s^{(3/2)-\nu^*} \hat{\psi}_N'(s). \quad (28)$$

The solution to Eq. (27) which decreases at $s \rightarrow +\infty$ (that is, at far tails of the density of states) is given (up to an arbitrary factor λ_N) by the function $G(s|\nu^*)$, Eq. (6). The factor λ_N can be found by fitting [16] the density of states $K_N(\varepsilon_s, \varepsilon_{s'})$, Eq. (8), to the bulk density of states, Eq. (24), near the soft edge provided $1 \ll s \ll N^{1/\nu^*}$. Then, making use of Eqs. (6), (8), and (28), we easily obtain that in the m th soft-edge scaling limit, Eq. (26), the two-point kernel

$$K_{\text{soft}}^{(m)}(s, s') = \lim_{N \rightarrow \infty} K_N(\varepsilon_s, \varepsilon_{s'}) \frac{d\varepsilon_s}{ds} \quad (29)$$

is determined by Eq. (5). In the particular case of $m = 0$ that is inherent in random-matrix ensembles with mono-

tonic confinement potential, the function G coincides with the Airy function, $G(s|\frac{3}{2}) = \text{Ai}(s)$, and the Airy correlations, Eq. (4), are recovered.

It follows from Eqs. (5) and (27) that the density of states in the same scaling limit

$$\nu_{\text{soft}}^{(m)}(s) = \left(\frac{d}{ds} G(s|\nu^*) \right)^2 s^{3/2-\nu^*} - [G(s|\nu^*)]^2 s^{\nu^*-1/2} \quad (30)$$

is also universal. The large- $|s|$ behavior of $\nu_{\text{soft}}^{(m)}$ can be deduced from the known asymptotic expansions of the Bessel functions,

$$\nu_{\text{soft}}^{(m)}(s) = \begin{cases} \frac{|s|^{\nu^*-1}}{\pi} + \frac{(-1)^{\nu^*-1/2}}{4\pi|s|} \cos(2|s|^{\nu^*}/\nu^*), & s \rightarrow -\infty, \\ \frac{\exp(-2s^{\nu^*}/\nu^*)}{4\pi s} \frac{\cos^2(\pi/4\nu^*)}{\sin(\pi/4\nu^*) + (-1)^{\nu^*-3/2}}, & s \rightarrow +\infty. \end{cases} \quad (31)$$

Note that the leading order behavior as $s \rightarrow -\infty$ is consistent with the $|s|^{\nu^*-1}$ singularity of the bulk density of states, Eq. (24).

To conclude, in this Letter we presented a general formalism for a treatment of eigenlevel correlations in spectra of $U(N)$ invariant ensembles of large random matrices with strong level confinement. An important ingredient of our analysis is the second-order differential equation which connects the Dyson's density with a fictitious "wave function" ψ_N which is needed for calculations of eigenvalue correlations within the framework of orthogonal polynomial technique. This consideration is relevant to an arbitrary energy range. We have applied this formalism to examine the eigenlevel correlations near the end point of single spectrum support. It has been shown that in the soft-edge scaling limit there are novel universal eigenlevel correlations which only depend on the even multicritical index of a matrix model. In a particular case $m = 0$, corresponding to monotonic confinement potentials, universality of the Airy correlations is recovered.

One of the authors (E. K.) acknowledges the support of the Levy Eshkol Fellowship from the Ministry of Science of Israel.

-
- [1] E. P. Wigner, *Ann. Math.* **53**, 36 (1951).
 [2] F. J. Dyson, *J. Math. Phys. (N.Y.)* **3**, 140 (1962); **3**, 157 (1962); **3**, 166 (1962).
 [3] T. A. Brody, J. Flores, J. B. French, P. A. Mello, A. Pandey, and S. S. M. Wong, *Rev. Mod. Phys.* **53**, 385 (1981).

-
- [4] D. J. Gross and A. A. Migdal, *Phys. Rev. Lett.* **64**, 127 (1990); P. Di Francesco, P. Ginsparg, and J. Zinn-Justin, *Phys. Rep.* **254**, 1 (1995).
 [5] J. J. M. Verbaarschot and I. Zahed, *Phys. Rev. Lett.* **73**, 2288 (1994).
 [6] M. C. Gutzwiller, *Chaos in Classical and Quantum Mechanics* (Springer-Verlag, New York, 1990).
 [7] See relevant papers in *Mesoscopic Phenomena in Solids*, edited by B. L. Altshuler, P. A. Lee, and R. A. Webb (North-Holland, Amsterdam, 1991).
 [8] M. L. Mehta, *Random Matrices* (Academic Press, Boston, 1991).
 [9] E. Brézin and A. Zee, *Nucl. Phys.* **B402**, 613 (1993).
 [10] G. Hackenbroich and H. A. Weidenmüller, *Phys. Rev. Lett.* **74**, 4118 (1995).
 [11] V. Freilikher, E. Kanzieper, and I. Yurkevich, *Phys. Rev. E* **54**, 210 (1996); **53**, 2200 (1996).
 [12] Namely, $V(\varepsilon)$ should increase at least as fast as $|\varepsilon|$ at infinity; see Refs. [10,11].
 [13] G. Akemann, P. H. Damgaard, U. Magnea, and S. Nishigaki, *Nucl. Phys.* **B487**, 721 (1997). Note that in this proof the confinement potential is stronger than quadratic, and parameter α is positive integer.
 [14] M. J. Bowick and E. Brézin, *Phys. Lett. B* **268**, 21 (1991).
 [15] P. J. Forrester, *Nucl. Phys.* **B402**, 709 (1993).
 [16] E. Kanzieper and V. Freilikher, *Phys. Rev. E* **55**, 3712 (1997).
 [17] J. Shohat, *C. R. Acad. Sci. Paris Ser. I Math.* **191**, 989 (1930); *Duke Math. J.* **5**, 401 (1939).
 [18] S. S. Bonan and D. S. Clark, *J. Approx. Theory* **46**, 408 (1986); **63**, 210 (1990).
 [19] G. Szegő, *Orthogonal Polynomials* (American Mathematical Society, Providence, 1967), Vol. XXIII.