

Limits on WWZ and $WW\gamma$ Couplings from $p\bar{p} \rightarrow e\nu jjX$ Events at $\sqrt{s} = 1.8$ TeV

B. Abbott,²⁸ M. Abolins,²⁵ B. S. Acharya,⁴³ I. Adam,¹² D. L. Adams,³⁷ M. Adams,¹⁷ S. Ahn,¹⁴ H. Aihara,²² G. A. Alves,¹⁰ E. Amidi,²⁹ N. Amos,²⁴ E. W. Anderson,¹⁹ R. Astur,⁴² M. M. Baarmand,⁴² A. Baden,²³ V. Balamurali,³² J. Balderston,¹⁶ B. Baldin,¹⁴ S. Banerjee,⁴³ J. Bantly,⁵ J. F. Bartlett,¹⁴ K. Bazizi,³⁹ A. Belyaev,²⁶ S. B. Beri,³⁴ I. Bertram,³¹ V. A. Bezzubov,³⁵ P. C. Bhat,¹⁴ V. Bhatnagar,³⁴ M. Bhattacharjee,¹³ N. Biswas,³² G. Blazey,³⁰ S. Blessing,¹⁵ P. Bloom,⁷ A. Boehnlein,¹⁴ N. I. Bojko,³⁵ F. Borcherding,¹⁴ J. Borders,³⁹ C. Boswell,⁹ A. Brandt,¹⁴ R. Brock,²⁵ A. Bross,¹⁴ D. Buchholz,³¹ V. S. Burtovoi,³⁵ J. M. Butler,³ W. Carvalho,¹⁰ D. Casey,³⁹ Z. Casilum,⁴² H. Castilla-Valdez,¹¹ D. Chakraborty,⁴² S.-M. Chang,²⁹ S. V. Chekulaev,³⁵ L.-P. Chen,²² W. Chen,⁴² S. Choi,⁴¹ S. Chopra,²⁴ B. C. Choudhary,⁹ J. H. Christenson,¹⁴ M. Chung,¹⁷ D. Claes,²⁷ A. R. Clark,²² W. G. Cobau,²³ J. Cochran,⁹ W. E. Cooper,¹⁴ C. Cretsinger,³⁹ D. Cullen-Vidal,⁵ M. A. C. Cummings,¹⁶ D. Cutts,⁵ O. I. Dahl,²² K. Davis,² K. De,⁴⁴ K. Del Signore,²⁴ M. Demarteau,¹⁴ D. Denisov,¹⁴ S. P. Denisov,³⁵ H. T. Diehl,¹⁴ M. Diesburg,¹⁴ G. Di Loreto,²⁵ P. Draper,⁴⁴ Y. Ducros,⁴⁰ L. V. Dudko,²⁶ S. R. Dugad,⁴³ D. Edmunds,²⁵ J. Ellison,⁹ V. D. Elvira,⁴² R. Engelmann,⁴² S. Eno,²³ G. Eppley,³⁷ P. Ermolov,²⁶ O. V. Eroshin,³⁵ V. N. Evdokimov,³⁵ T. Fahland,⁸ M. Fatyga,⁴ M. K. Fatyga,³⁹ J. Featherly,⁴ S. Feher,¹⁴ D. Fein,² T. Ferbel,³⁹ G. Finocchiaro,⁴² H. E. Fisk,¹⁴ Y. Fisyak,⁷ E. Flattum,¹⁴ G. E. Forden,² M. Fortner,³⁰ K. C. Frame,²⁵ S. Fuess,¹⁴ E. Gallas,⁴⁴ A. N. Galyaev,³⁵ P. Gartung,⁹ T. L. Geld,²⁵ R. J. Genik II,²⁵ K. Genser,¹⁴ C. E. Gerber,¹⁴ B. Gibbard,⁴ S. Glenn,⁷ B. Gobbi,³¹ M. Goforth,¹⁵ A. Goldschmidt,²² B. Gómez,¹ G. Gómez,²³ P. I. Goncharov,³⁵ J. L. González Solís,¹¹ H. Gordon,⁴ L. T. Goss,⁴⁵ K. Gounder,⁹ A. Goussiou,⁴² N. Graf,⁴ P. D. Grannis,⁴² D. R. Green,¹⁴ J. Green,³⁰ H. Greenlee,¹⁴ G. Grim,⁷ S. Grinstein,⁶ N. Grossman,¹⁴ P. Grudberg,²² S. Grünendahl,³⁹ G. Guglielmo,³³ J. A. Guida,² J. M. Guida,⁵ A. Gupta,⁴³ S. N. Gurzhiev,³⁵ P. Gutierrez,³³ Y. E. Gutnikov,³⁵ N. J. Hadley,²³ H. Haggerty,¹⁴ S. Hagopian,¹⁵ V. Hagopian,¹⁵ K. S. Hahn,³⁹ R. E. Hall,⁸ S. Hansen,¹⁴ J. M. Hauptman,¹⁹ D. Hedin,³⁰ A. P. Heinson,⁹ U. Heintz,¹⁴ R. Hernández-Montoya,¹¹ T. Heuring,¹⁵ R. Hirosky,¹⁵ J. D. Hobbs,¹⁴ B. Hoeneisen,^{1,*} J. S. Hoftun,⁵ F. Hsieh,²⁴ Ting Hu,⁴² Tong Hu,¹⁸ T. Huehn,⁹ A. S. Ito,¹⁴ E. James,² J. Jaques,³² S. A. Jerger,²⁵ R. Jesik,¹⁸ J. Z.-Y. Jiang,⁴² T. Joffe-Minor,³¹ K. Johns,² M. Johnson,¹⁴ A. Jonckheere,¹⁴ M. Jones,¹⁶ H. Jöstlein,¹⁴ S. Y. Jun,³¹ C. K. Jung,⁴² S. Kahn,⁴ G. Kalbfleisch,³³ J. S. Kang,²⁰ R. Kehoe,³² M. L. Kelly,³² C. L. Kim,²⁰ S. K. Kim,⁴¹ A. Klatchko,¹⁵ B. Klima,¹⁴ C. Klopfenstein,⁷ V. I. Klyukhin,³⁵ V. I. Kochetkov,³⁵ J. M. Kohli,³⁴ D. Koltick,³⁶ A. V. Kostritskiy,³⁵ J. Kotcher,⁴ A. V. Kotwal,¹² J. Kourlas,²⁸ A. V. Kozelov,³⁵ E. A. Kozlovski,³⁵ J. Krane,²⁷ M. R. Krishnaswamy,⁴³ S. Krzywdzinski,¹⁴ S. Kunori,²³ S. Lami,⁴² H. Lan,^{14,†} R. Lander,⁷ F. Landry,²⁵ G. Landsberg,¹⁴ B. Lauer,¹⁹ A. Leflat,²⁶ H. Li,⁴² J. Li,⁴⁴ Q. Z. Li-Demarteau,¹⁴ J. G. R. Lima,³⁸ D. Lincoln,²⁴ S. L. Linn,¹⁵ J. Linnemann,²⁵ R. Lipton,¹⁴ Q. Liu,^{14,*} Y. C. Liu,³¹ F. Lobkowicz,³⁹ S. C. Loken,²² S. Lökös,⁴² L. Lueking,¹⁴ A. L. Lyon,²³ A. K. A. Maciel,¹⁰ R. J. Madaras,²² R. Madden,¹⁵ L. Magaña-Mendoza,¹¹ S. Mani,⁷ H. S. Mao,^{14,†} R. Markeloff,³⁰ L. Markosky,² T. Marshall,¹⁸ M. I. Martin,¹⁴ K. M. Mauritz,¹⁹ B. May,³¹ A. A. Mayorov,³⁵ R. McCarthy,⁴² J. McDonald,¹⁵ T. McKibben,¹⁷ J. McKinley,²⁵ T. McMahon,³³ H. L. Melanson,¹⁴ M. Merkin,²⁶ K. W. Merritt,¹⁴ H. Miettinen,³⁷ A. Mincer,²⁸ J. M. de Miranda,¹⁰ C. S. Mishra,¹⁴ N. Mokhov,¹⁴ N. K. Mondal,⁴³ H. E. Montgomery,¹⁴ P. Mooney,¹ H. da Motta,¹⁰ C. Murphy,¹⁷ F. Nang,² M. Narain,¹⁴ V. S. Narasimhan,⁴³ A. Narayanan,² H. A. Neal,²⁴ J. P. Negret,¹ P. Nemethy,²⁸ M. Nicola,¹⁰ D. Norman,⁴⁵ L. Oesch,²⁴ V. Oguri,³⁸ E. Oltman,²² N. Oshima,¹⁴ D. Owen,²⁵ P. Padley,³⁷ M. Pang,¹⁹ A. Para,¹⁴ Y. M. Park,²¹ R. Partridge,⁵ N. Parua,⁴³ M. Paterno,³⁹ J. Perkins,⁴⁴ M. Peters,¹⁶ R. Piegaiia,⁶ H. Piekarz,¹⁵ Y. Pischalnikov,³⁶ V. M. Podstavkov,³⁵ B. G. Pope,²⁵ H. B. Prosper,¹⁵ S. Protopopescu,⁴ J. Qian,²⁴ P. Z. Quintas,¹⁴ R. Raja,¹⁴ S. Rajagopalan,⁴ O. Ramirez,¹⁷ L. Rasmussen,⁴² S. Reucroft,²⁹ M. Rijssenbeek,⁴² T. Rockwell,²⁵ N. A. Roe,²² P. Rubinov,³¹ R. Ruchti,³² J. Rutherford,² A. Sánchez-Hernández,¹¹ A. Santoro,¹⁰ L. Sawyer,⁴⁴ R. D. Schamberger,⁴² H. Schellman,³¹ J. Sculli,²⁸ E. Shabalina,²⁶ C. Shaffer,¹⁵ H. C. Shankar,⁴³ R. K. Shivpuri,¹³ M. Shupe,² H. Singh,⁹ J. B. Singh,³⁴ V. Sirotenko,³⁰ W. Smart,¹⁴ A. Smith,² R. P. Smith,¹⁴ R. Snihur,³¹ G. R. Snow,²⁷ J. Snow,³³ S. Snyder,⁴ J. Solomon,¹⁷ P. M. Sood,³⁴ M. Sosebee,⁴⁴ N. Sotnikova,²⁶ M. Souza,¹⁰ A. L. Spadafora,²² R. W. Stephens,⁴⁴ M. L. Stevenson,²² D. Stewart,²⁴ D. A. Stoianova,³⁵ D. Stoker,⁸ M. Strauss,³³ K. Streets,²⁸ M. Strovink,²² A. Sznajder,¹⁰ P. Tamburello,²³ J. Tarazi,⁸ M. Tartaglia,¹⁴ T. L. T. Thomas,³¹ J. Thompson,²³ T. G. Trippe,²² P. M. Tuts,¹² N. Varelas,²⁵ E. W. Varnes,²² D. Vititoe,² A. A. Volkov,³⁵ A. P. Vorobiev,³⁵ H. D. Wahl,¹⁵ G. Wang,¹⁵ J. Warchol,³² G. Watts,⁵ M. Wayne,³² H. Weerts,²⁵ A. White,⁴⁴ J. T. White,⁴⁵ J. A. Wightman,¹⁹ S. Willis,³⁰ S. J. Wimpenny,⁹ J. V. D. Wirjawan,⁴⁵ J. Womersley,¹⁴ E. Won,³⁹ D. R. Wood,²⁹ H. Xu,⁵ R. Yamada,¹⁴ P. Yamin,⁴ C. Yanagisawa,⁴² J. Yang,²⁸ T. Yasuda,²⁹ P. Yepes,³⁷ C. Yoshikawa,¹⁶ S. Youssef,¹⁵ J. Yu,¹⁴ Y. Yu,⁴¹ Z. H. Zhu,³⁹ D. Zieminska,¹⁸ A. Zieminski,¹⁸ E. G. Zverev,²⁶ and A. Zylberstejn⁴⁰

(D0 Collaboration)

- ¹Universidad de los Andes, Bogotá, Colombia
²University of Arizona, Tucson, Arizona 85721
³Boston University, Boston, Massachusetts 02215
⁴Brookhaven National Laboratory, Upton, New York 11973
⁵Brown University, Providence, Rhode Island 02912
⁶Universidad de Buenos Aires, Buenos Aires, Argentina
⁷University of California, Davis, California 95616
⁸University of California, Irvine, California 92697
⁹University of California, Riverside, California 92521
¹⁰LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil
¹¹CINVESTAV, Mexico City, Mexico
¹²Columbia University, New York, New York 10027
¹³Delhi University, Delhi, India 110007
¹⁴Fermi National Accelerator Laboratory, Batavia, Illinois 60510
¹⁵Florida State University, Tallahassee, Florida 32306
¹⁶University of Hawaii, Honolulu, Hawaii 96822
¹⁷University of Illinois at Chicago, Chicago, Illinois 60607
¹⁸Indiana University, Bloomington, Indiana 47405
¹⁹Iowa State University, Ames, Iowa 50011
²⁰Korea University, Seoul, Korea
²¹Kyungshung University, Pusan, Korea
²²Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720
²³University of Maryland, College Park, Maryland 20742
²⁴University of Michigan, Ann Arbor, Michigan 48109
²⁵Michigan State University, East Lansing, Michigan 48824
²⁶Moscow State University, Moscow, Russia
²⁷University of Nebraska, Lincoln, Nebraska 68588
²⁸New York University, New York, New York 10003
²⁹Northeastern University, Boston, Massachusetts 02115
³⁰Northern Illinois University, DeKalb, Illinois 60115
³¹Northwestern University, Evanston, Illinois 60208
³²University of Notre Dame, Notre Dame, Indiana 46556
³³University of Oklahoma, Norman, Oklahoma 73019
³⁴University of Panjab, Chandigarh 16-00-14, India
³⁵Institute for High Energy Physics, 142-284 Protvino, Russia
³⁶Purdue University, West Lafayette, Indiana 47907
³⁷Rice University, Houston, Texas 77005
³⁸Universidade Estadual do Rio de Janeiro, Rio de Janeiro, Brazil
³⁹University of Rochester, Rochester, New York 14627
⁴⁰CEA, DAPNIA/Service de Physique des Particules, CE-SACLAY, Gif-sur-Yvette, France
⁴¹Seoul National University, Seoul, Korea
⁴²State University of New York, Stony Brook, New York 11794
⁴³Tata Institute of Fundamental Research, Colaba, Mumbai 400005, India
⁴⁴University of Texas, Arlington, Texas 76019
⁴⁵Texas A&M University, College Station, Texas 77843

(Received 15 May 1997)

We present limits on anomalous WWZ and $WW\gamma$ couplings from a search for WW and WZ production in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV. We use $p\bar{p} \rightarrow e\nu jjX$ events recorded with the D0 detector at the Fermilab Tevatron Collider during the 1992–1995 run. The data sample corresponds to an integrated luminosity of 96.0 ± 5.1 pb⁻¹. Assuming identical WWZ and $WW\gamma$ coupling parameters, the 95% C.L. limits on the CP -conserving couplings are $-0.33 < \lambda < 0.36$ ($\Delta\kappa = 0$) and $-0.43 < \Delta\kappa < 0.59$ ($\lambda = 0$), for a form factor scale $\Lambda = 2.0$ TeV. Limits based on other assumptions are also presented. [S0031-9007(97)03883-0]

PACS numbers: 13.85.Rm, 12.15.Ji, 13.40.Gp, 14.70.Fm

The vector boson trilinear couplings predicted by the non-Abelian gauge symmetry of the standard model (SM) can be measured directly in pair production processes such as $q\bar{q} \rightarrow W^+W^-$, $W^\pm\gamma$, $Z\gamma$, and $W^\pm Z$. Devia-

tions from the SM couplings would signal new physics. Studies of such effects have been reported by the UA2 [1], CDF [2], D0 [3–6], and LEP [7] Collaborations. In this Letter we report on the measurement of WWV

couplings (where $V = \gamma$ or Z) using the diboson production processes $p\bar{p} \rightarrow WWX \rightarrow evjjX$ and $p\bar{p} \rightarrow WZX \rightarrow evjjX$, where j represents a jet.

The Lorentz invariant Lagrangian which describes the $WW\gamma$ and WWZ interactions has 14 independent coupling parameters [8], seven describing the $WW\gamma$ vertex and seven for the WWZ vertex. Assuming electromagnetic gauge invariance and CP conservation, the number of parameters is reduced to five: g_1^Z , κ_Z , κ_γ , λ_Z , and λ_γ . In the SM at tree level, the coupling parameters have the values $\Delta g_1^Z (\equiv g_1^Z - 1) = 0$, $\Delta \kappa_Z (\equiv \kappa_Z - 1) = \Delta \kappa_\gamma (\equiv \kappa_\gamma - 1) = 0$, $\lambda_Z = \lambda_\gamma = 0$. The SM cross sections for $p\bar{p} \rightarrow W^+W^-X$ and $p\bar{p} \rightarrow W^\pm ZX$ production at the Tevatron, at $\sqrt{s} = 1.8$ TeV, are 9.5 and 2.7 pb [9], respectively.

Non-SM values of the coupling parameters would result in an increase of the production cross section, especially for large values of the transverse momentum of the W boson (p_T^W). Since tree level unitarity restricts the WWV couplings to their SM values at asymptotically high energies, each of the couplings must be modified by a form factor, e.g., $\lambda_Z(\hat{s}) = \lambda_Z/(1 + \hat{s}/\Lambda^2)^2$, where \hat{s} is the square of the invariant mass of the WW or WZ system and Λ is the form-factor scale. We have used $\Lambda = 1.0, 1.5$, and 2.0 TeV.

The analysis reported here uses $p\bar{p} \rightarrow evjjX$ events recorded with the D0 detector during the 1992–1993 and 1993–1995 Fermilab Tevatron Collider runs at $\sqrt{s} = 1.8$ TeV, corresponding to a total integrated luminosity of 96.0 ± 5.1 pb $^{-1}$. The D0 detector and data collection system are described elsewhere [10]. The basic elements of the trigger and reconstruction algorithms for jets, electrons, and neutrinos are given in Ref. [6]. The analysis of $evjjX$ events from the 1992–1993 Tevatron Collider run (13.7 ± 0.7 pb $^{-1}$) was reported previously [5]. This Letter focuses on the analysis of the 1993–1995 data set (82.3 ± 4.4 pb $^{-1}$), using a method slightly different from that of Ref. [5], and gives the combined result from both analyses. Further details are available in Ref. [11].

The data sample was obtained with a trigger which required an isolated electromagnetic (EM) calorimeter cluster with transverse energy $E_T > 20$ GeV and missing transverse energy $\cancel{E}_T > 15$ GeV. The off-line event selection required that the EM cluster have $|\eta| < 1.1$ in the central calorimeter or $1.5 < |\eta| < 2.5$ in an end calorimeter, where η is the pseudorapidity. Electrons were identified by requiring that the EM cluster pass the shower profile and tracking information criteria, as described in our earlier analysis [5]. The presence of a neutrino was inferred from the \cancel{E}_T , calculated from the vector sum of the E_T measured in each calorimeter tower. Jets were reconstructed using a cone algorithm with radius $\mathcal{R} \equiv \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} = 0.5$. To remove spurious jets due to detector effects, this analysis used the same quality cuts as were used in Ref. [12]. Jets were required to be within $|\eta| < 2.5$. The jet energies were

corrected for effects of jet energy scale calibration, out-of-cone showering, energy from the underlying event [13], and energy loss due to out-of-cone gluon radiation.

The WW/WZ candidates were selected by searching for events containing an isolated electron with high E_T^e (>25 GeV), large \cancel{E}_T (>25 GeV), and at least two high E_T^j jets (>20 GeV). The transverse mass of the electron and neutrino system was required to be consistent with a $W \rightarrow e\nu$ decay [$M_T^{e\nu} = [2E_T^e\cancel{E}_T(1 - \cos\phi^{e\nu})/c^4]^{1/2} > 40$ GeV/ c^2 , where $\phi^{e\nu}$ is the azimuthal angle between the electron and \cancel{E}_T vector]. The invariant mass (m^{jj}) of the two jet system (the largest invariant mass if there were more than two jets with $E_T^j > 20$ GeV in the event) was required to be $50 < m^{jj} < 110$ GeV/ c^2 , as expected for a $W \rightarrow jj$ or $Z \rightarrow jj$ decay. Monte Carlo studies showed that the dijet invariant mass resolution for signal events is 16 GeV/ c^2 . The scalar transverse momenta of the two gauge bosons were required to be the same within 40 GeV/ c , as is expected for WW/WZ production.

There are two major sources of background to $WW/WZ \rightarrow evjj$ production: (i) $W + \geq 2$ jets with $W \rightarrow e\nu$, and (ii) QCD multijet events where one of the jets is misidentified as an electron and there is significant \cancel{E}_T in the event due to mismeasurement. Other backgrounds, such as $t\bar{t}$ production with subsequent decay to $W^+bW^-\bar{b}$ followed by $W \rightarrow e\nu$, WW or WZ production with $W \rightarrow \tau\nu$ followed by $\tau \rightarrow e\nu\bar{\nu}$, $ZX \rightarrow eeX$, where one electron is mismeasured or not identified, and $ZX \rightarrow \tau\tau X$ with $\tau \rightarrow e\nu\bar{\nu}$, are negligible.

The $W + \geq 2$ jets background was estimated using the VECBOS [14] event generator, with $Q^2 = (p_T^j)^2$, followed by parton fragmentation using the HERWIG [15] package and a detailed GEANT [16] based simulation of the detector. Normalization of the $W + \geq 2$ jets background was determined by comparing the number of events expected from the VECBOS estimate to the number of candidate events outside the dijet mass window, after the multijet backgrounds had been subtracted. The systematic uncertainties in this background are due to the normalization and to the jet energy scale correction. The multijet background was estimated following the same procedure used in our previous analysis [5]. The background sample, which consisted of data events containing a jet satisfying the electron trigger selection but failing the electron identification, was normalized to the signal sample in the region $\cancel{E}_T < 15$ GeV where the actual WW/WZ contribution is negligible. The number of background events was then determined from this scaled background sample with the rest of the selection criteria applied [11]. The backgrounds from $t\bar{t} \rightarrow W^+bW^-\bar{b}$ and other minor sources were estimated using the ISAJET event generator [17] followed by detector simulation. Table I summarizes the background estimates and the total number of events seen. The number of observed events was consistent with the background estimates which dominate the SM WW/WZ signal.

TABLE I. Summary of signal and backgrounds.

	1992–1993	1993–1995
Luminosity	13.7 pb ⁻¹	82.3 pb ⁻¹
Backgrounds		
W + ≥2 jets	62.2 ± 13.0	279.5 ± 36.0
QCD multijet	12.2 ± 2.6	104.3 ± 12.3
<i>t</i> \bar{t} → <i>e</i> <i>v</i> <i>jj</i> X	0.9 ± 0.1	3.7 ± 1.3
Total background	75.3 ± 13.3	387.5 ± 38.1
Data	84	399
SM WW + WZ prediction	3.2 ± 0.6	17.5 ± 3.0

The trigger and off-line electron identification efficiencies were estimated using $Z \rightarrow ee$ events. The trigger efficiency was $(98.1 \pm 1.9)\%$ [4]. The electron identification efficiencies were found to be $(74.5 \pm 1.1)\%$ in the central calorimeter and $(61.9 \pm 1.1)\%$ in the end calorimeters. We studied the $W \rightarrow jj$ efficiency for the jet cone size $\mathcal{R} = 0.5$ using the ISAJET and PYTHIA [18] event generators followed by detector simulation. The selection criteria were applied to these samples and it was found that the efficiency was $\approx 50\%$ for $p_T^W < 250$ GeV/*c* and that this decreased significantly for $p_T^W > 250$ GeV/*c* due to merging of the two jets into one. The efficiencies obtained from ISAJET were used to estimate the detection efficiencies of the WW(WZ) processes since they gave more conservative results.

The overall event selection efficiency was calculated using the leading order event generator of Ref. [19] to generate four momenta for WW and WZ processes as a function of the coupling parameter values. A fast detector simulation was used to take into account the detector resolutions and efficiencies described above. Higher order QCD effects were approximated by a *K*-factor = $1 + \frac{8}{9}\pi\alpha_s = 1.34$ [9] and a smearing of the transverse momentum of the diboson system according to the experimentally determined p_T^Z spectrum from the inclusive $Z \rightarrow ee$ sample. The total selection efficiencies for the detection of SM WW and WZ events were estimated to be $[14.7 \pm 02(\text{stat}) \pm 1.2(\text{syst})]\%$ and $[14.6 \pm 0.4(\text{stat}) \pm 1.1(\text{syst})]\%$, respectively. The systematic uncertainty (8%) includes electron trigger and selection efficiencies (1%), \cancel{E}_T smearing and p_T of the WW/WZ diboson system (5%), difference between the ISAJET and PYTHIA programs for $W \rightarrow jj$ efficiency parametrization (5%), statistical uncertainty for $W \rightarrow jj$ efficiency parametrization (2%), and jet energy scale (3%).

The expected signal for WW plus WZ production with SM couplings is 20.7 ± 3.2 events based on the total integrated luminosity of 96.0 pb⁻¹. The uncertainty is the sum in quadrature of the uncertainties in the selection efficiency, luminosity, and higher order QCD corrections to the expected signal (14%) [9]. Figure 1 shows the $p_T^{e\nu}$ distributions for candidate events from 1993–1995 data, total background estimate plus SM expectations, and SM expectations for WW and WZ production, after all selection criteria have been applied. There is no




FIG. 1. p_T distributions of the $e\nu$ system for the 1993–1995 data set. The points with error bars represent the data. The solid histogram is the total background estimate plus the SM Monte Carlo predictions of WW and WZ production (shown as shaded histogram). The inset shows the predicted $d\sigma/dp_T^W$, folded with the detection efficiencies, for SM WW γ and WWZ couplings (lower curve), and for SM WW γ and the indicated anomalous WWZ couplings (upper curve).

clear difference between the observed $p_T^{e\nu}$ spectrum and that expected from background plus SM WW and WZ prediction.

Using the detection efficiencies for SM WW and WZ production and the background subtracted signal, and assuming the SM ratio of cross sections for WW and WZ production, we can set an upper limit at the 95% C.L. on the cross section $\sigma(p\bar{p} \rightarrow W^+W^-X)$ of 76 pb.

Since we observed no excess of high $p_T^{e\nu}$ events, large deviations from the SM trilinear coupling values are excluded. Limits on the anomalous coupling parameters were set by performing a binned likelihood fit to the observed $p_T^{e\nu}$ spectrum with the Monte Carlo signal prediction plus the estimated background. Unequal width bins were used to evenly distribute the observed events,




FIG. 2. Limits on CP -conserving anomalous couplings parameters: (a) $\Delta\kappa \equiv \Delta\kappa_\gamma = \Delta\kappa_Z$, $\lambda \equiv \lambda_\gamma = \lambda_Z$, (b) HISZ relations, and (c),(d) SM WW γ couplings. The inner and middle curves represent 95% C.L. one and two degree-of-freedom exclusion contours, respectively. The outermost curves show *S* matrix unitarity bounds. $\Lambda = 1.5$ TeV is used for all four cases. The SM predictions is $\Delta\kappa = 0$, $\lambda = 0$, $\Delta g_1^Z = 0$.

TABLE II. Axis limits at the 95% C.L. with various assumptions and three different Λ values.

Couplings/ Λ (TeV)		1.0	1.5	2.0
(i)	$\lambda_\gamma = \lambda_Z$	-0.42, 0.45	-0.36, 0.39	-0.33, 0.36
	$\Delta\kappa_\gamma = \Delta\kappa_Z$	-0.55, 0.79	-0.47, 0.63	-0.43, 0.59
(ii)	λ_γ (HISZ)	-0.42, 0.45	-0.36, 0.38	-0.34, 0.36
	$\Delta\kappa_\gamma$ (HISZ)	-0.69, 1.04	-0.56, 0.85	-0.53, 0.78
(iii)	λ_Z (SM $WW\gamma$)	-0.47, 0.51	-0.40, 0.43	-0.37, 0.40
	$\Delta\kappa_Z$ (SM $WW\gamma$)	-0.74, 0.99	-0.60, 0.79	-0.54, 0.72
	Δg_1^Z (SM $WW\gamma$)	-0.75, 1.06	-0.64, 0.89	-0.60, 0.81
(iv)	λ_γ (SM WWZ)	-1.28, 1.33	-1.21, 1.25	
	$\Delta\kappa_\gamma$ (SM WWZ)	-1.60, 2.03	-1.38, 1.70	

especially those at the end of the spectrum. In each $p_T^{e\nu}$ bin for a given set of anomalous coupling parameters, we calculated the probability for the sum of the background estimate and Monte Carlo WW/WZ prediction to fluctuate to the observed number of events. The limits on the anomalous coupling parameters are from a combined likelihood fit to both data sets. The uncertainties in the background estimates, efficiencies, integrated luminosity, and higher order QCD corrections to the signal were convoluted with Gaussian distributions into the likelihood function. Uncertainties common to both analyses, e.g., theoretical uncertainties, were convoluted only once.

In Fig. 2, bounds on four pairs of coupling parameters are shown using $\Lambda = 1.5$ TeV. In each case all other couplings are fixed to their SM values. The one and two degree-of-freedom 95% C.L. contour limits (corresponding to likelihood function values 1.92 and 3.00 units below the maximum, respectively) are shown as the inner curves, along with the S matrix unitarity limits, shown as the outermost curves, which are obtained by evaluating all (i.e., WW , $W\gamma$, and WZ) processes. Figure 2(a) shows the contour limits when coupling parameters for $WW\gamma$ are assumed to be equal to those for WWZ . Figure 2(b) shows contour limits assuming Hagiwara-Ishihara-Szalapski-Zeppenfeld (HISZ) relations [20]. In Figs. 2(c) and 2(d) SM $WW\gamma$ couplings are assumed and the coupling limits for WWZ are shown.

When SM $WW\gamma$ couplings are assumed, the U(1) point ($\kappa_Z = 0$, $\lambda_Z = 0$, $g_1^Z = 0$) is excluded at the 99% C.L. This is direct evidence for the existence of the WWZ couplings.

Table II lists the 95% C.L. axis limits for three different values of Λ and four assumptions: (i) $\Delta\kappa \equiv \Delta\kappa_\gamma = \Delta\kappa_Z$, $\lambda \equiv \lambda_\gamma = \lambda_Z$, (ii) HISZ relations, (iii) SM $WW\gamma$ couplings, and (iv) SM WWZ couplings. The results with the SM $WW\gamma$ assumption are unique to WW/WZ production since the WWZ couplings are not accessible with $W\gamma$ production. The results indicate that this analysis is more sensitive to WWZ couplings than to $WW\gamma$ ones as expected from the larger overall couplings for WWZ than $WW\gamma$ [8]. The dependence of the coupling parameter limits on Λ is clearly seen. As Λ is increased, the experimental limits tighten while the limits from S -matrix

unitarity tighten more rapidly. When the unitarity limits are tighter than the experimental limits, as occurs for the SM WWZ couplings with $\Lambda = 2.0$ TeV, the experimental limits no longer are relevant.

In conclusion, we have presented limits on anomalous WWZ and $WW\gamma$ coupling parameters which are the most stringent to date. They are significantly tighter than those from the analyses of the 1992–1993 data set [2,5], and significantly better on $\Delta\kappa$ (but comparable on λ) to those measured using $W\gamma$ production with the 1992–1995 data set [4].

We thank U. Baur for useful discussions and D. Zeppenfeld for providing us with the WW and WZ Monte Carlo generators. We thank the staffs at Fermilab and collaborating institutions for their contributions to this work, and acknowledge support from the Department of Energy and National Science Foundation (U.S.A.), Commissariat à l’Energie Atomique (France), State Committee for Science and Technology and Ministry for Atomic Energy (Russia), CNPq (Brazil), Departments of Atomic Energy and Science and Education (India), Colciencias (Colombia), CONACyT (Mexico), Ministry of Education and KOSEF (Korea), CONICET and UBACyT (Argentina), and the A. P. Sloan Foundation.

*Visitor from Universidad San Francisco de Quito, Quito, Ecuador.

†Visitor from IHEP, Beijing, China.

- [1] UA2 Collaboration, J. Alitti *et al.*, Phys. Lett. B **277**, 194 (1992).
- [2] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **74**, 1936 (1995); **74**, 1941 (1995); **75**, 1017 (1995); Fermilab-Pub-96/311-E (to be published).
- [3] D0 Collaboration, S. Abachi *et al.*, Phys. Rev. Lett. **75**, 1023 (1995); **75**, 1028 (1995); **75**, 1034 (1995); **78**, 3640 (1997).
- [4] D0 Collaboration, S. Abachi *et al.*, Phys. Rev. Lett. **78**, 3634 (1997).
- [5] D0 Collaboration, S. Abachi *et al.*, Phys. Rev. Lett. **77**, 3303 (1996).
- [6] D0 Collaboration, S. Abachi *et al.*, Fermilab-Pub-97/088-E, hep-ex/9704004 (to be published).
- [7] OPAL Collaboration, K. Ackerstaff *et al.*, CERN-PPE/97-004 (to be published); DELPHI Collaboration,

- P. Abreu *et al.*, CERN-PPE/97-009 (to be published); L3 Collaboration, M. Acciarri *et al.*, CERN-PPE/97-014 (to be published).
- [8] K. Hagiwara, R. D. Peccei, D. Zeppenfeld, and K. Hikasa, Nucl. Phys. **B282**, 253 (1987).
- [9] J. Ohnemus, Phys. Rev. D **44**, 1403 (1991); **44**, 3477 (1991).
- [10] D0 Collaboration, S. Abachi *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **338**, 185 (1994).
- [11] A. Sánchez-Hernández, Ph.D. dissertation, CINVESTAV, Mexico City, Mexico, 1997 (unpublished).
- [12] D0 Collaboration, S. Abachi *et al.*, Phys. Rev. Lett. **75**, 618 (1995).
- [13] D0 Collaboration, R. Kehoe *et al.*, in *Proceedings of the 6th International Conference on Calorimetry in High Energy Physics, Frascati, 1996*, Frascati Physics Series Vol. VI (Frascati, Bologna, 1997), pp. 349–358.
- [14] F. A. Berends *et al.*, Nucl. Phys. **B357**, 32 (1991). We used version 3.0.
- [15] G. Marchesini *et al.*, Comput. Phys. Commun. **67**, 465 (1992). We used version 5.7.
- [16] F. Carminati *et al.*, *GEANT Users Guide*, CERN Program Library Long Writeup WS013, 1993 (unpublished).
- [17] F. Paige and S. Protopopescu, BNL Report BNL38034, 1986 (unpublished). We used version 7.22.
- [18] T. Sjöstrand, Comput. Phys. Commun. **82**, 74 (1994).
- [19] K. Hagiwara, J. Woodside, and D. Zeppenfeld, Phys. Rev. D **41**, 2113 (1990); D. Zeppenfeld (private communication).
- [20] K. Hagiwara, S. Ishihara, R. Szalapski, and D. Zeppenfeld, Phys. Rev. D **48**, 2182 (1993); Phys. Lett. B **283**, 353 (1992). They parametrize the $WW\gamma$ couplings in terms of the $WW\gamma$ couplings: $\Delta\kappa_Z = \Delta\kappa_\gamma(1 - \tan^2\theta_w)/2$, $\Delta g_1^Z = \Delta\kappa_\gamma/2 \cos^2\theta_w$, and $\lambda_Z = \lambda_\gamma$.