

## Gamma Rays and Neutrinos from the Crab Nebula Produced by Pulsar Accelerated Nuclei

W. Bednarek\* and R. J. Protheroe

*Department of Physics and Mathematical Physics, University of Adelaide, Adelaide, SA 5005, Australia*  
(Received 7 April 1997)

We investigate the consequences of the acceleration of heavy nuclei (e.g., iron nuclei) by the Crab pulsar. Accelerated nuclei can photodisintegrate in collisions with soft photons produced in the pulsar's outer gap, injecting energetic neutrons which decay either inside or outside the Crab nebula. The protons from neutron decay inside the nebula accumulate, producing gamma rays and neutrinos in collisions with the matter in the nebula. Neutrons decaying outside contribute to the galactic cosmic rays. We compute the expected fluxes of gamma rays and neutrinos, and find that our model could account for the observed emission at high energies and may be tested by searching for high energy neutrinos with future neutrino telescopes currently in the design stage. [S0031-9007(97)04179-3]

PACS numbers: 98.70.Sa, 95.85.Ry, 97.60.Gb, 98.70.Rz

The Crab nebula is a well established  $\gamma$ -ray source with a spectrum extending up to at least TeV energies [1]. Its emission below a few tens of MeV is interpreted as synchrotron emission by  $\sim 10^{15-16}$  eV electrons in the Crab nebula magnetic field of strength  $\sim 3 \times 10^{-4}$  G [2]. These models interpret the emission at higher energies as being due this same population of electrons by inverse Compton scattering (ICS) of synchrotron photons (SSC models). The TeV  $\gamma$  rays might also originate near the light cylinder as a result of ICS of infrared photons from another outer gap [3]. The possibility of hadronic processes has also been considered by Cheng *et al.* [4] who assumed that relativistic protons accelerated in the Crab pulsar outer gap interact with the matter inside the nebula, and noted that this may result in TeV  $\gamma$  rays.

In this Letter, we analyze the consequences of acceleration of heavy nuclei in the pulsar magnetosphere as a possible mechanism of energetic radiation from the Crab nebula. The importance of photodisintegration of nuclei is determined by the reciprocal mean free path which we calculate using cross sections of Karakuła and Tkaczyk [5] and show in Fig. 1 for nuclei during propagation in the radiation field supposed to be present in the outer gap of the Crab pulsar. Ho [6] has shown that the total luminosity and spectrum of the outer gap radiation is not strongly dependent on  $P$ . For  $P = 33$  ms we adopt the photon number density given by Eq. 7.1 in Paper II of Cheng *et al.* [7], and for other periods we scale this by  $r_{LC}^{-3}$ . We assume this radiation is produced by cascading due to electrons and positrons accelerated in the gap, and so is highly anisotropic and directed across the gap (in both directions). The dimension of the outer gap is of the order of the radius of the light cylinder,  $l_{gap} \approx r_{LC} = c/\Omega$ . For the case of the Crab pulsar,  $r_{LC} \approx 1.5 \times 10^8$  cm, and it is evident from Fig. 1 that multiple photodisintegrations of primary Fe nuclei will occur provided they can be accelerated to Lorentz factors above  $\sim 10^3$ . We have also computed the mean interaction length for pion photoproduction using the cross section given by Stecker [8], and the energy loss length for pair production using the approximations

given by Chodorowski *et al.* [9]. The results are shown in Fig. 1, and it is clear that these latter processes are not important loss mechanisms compared to photodisintegration.

We assume that Fe nuclei can escape from the polar cap surface of the Crab pulsar, and move along magnetic field lines to enter the outer gap where they can be accelerated in the outer gap potential as in the model of Cheng *et al.* [7] and Ho [6]. The nuclei accelerated in the outer gap will interact with photons either producing secondary  $e^\pm$  pairs (with negligible loss of energy) or extracting a nucleon. The pairs could eventually saturate the electric field of the gap, although this is far from certain. We shall assume that for some reason the field is not shorted. Possibilities include: (a) Nuclei, having much larger Larmor radii and much lower synchrotron and curvature losses than electrons, more easily drift across the magnetic field lines away from the region of the gap where they had produced pairs; (b) the secondary  $e^\pm$  pairs may




FIG. 1. Reciprocal mean free paths for photodisintegration of nuclei with mass numbers,  $A = 56, 40, 32, 24, 16, 8,$  and  $4$  (full curves from the top) in the radiation field of the Crab pulsar's outer gap as a function of the Lorentz factor of the nuclei. The reciprocal mean free path for pion photoproduction (dotted curve) and the energy loss length for  $e^\pm$  pair production (dashed curve) by iron nuclei are also shown.

not be able to saturate the electric field in the outer gap which is close to the light cylinder since the Goldreich and Julian [10] density tends to infinity where the gap approaches the light cylinder.

In order to obtain the energy spectrum of neutrons extracted from Fe nuclei,  $N_n(\gamma_n)$ , we simulate their acceleration and propagation through the outer gap using a Monte Carlo method. To obtain the rate of injection of neutrons per unit energy we multiply  $N_n(\gamma_n)$  by the number of Fe nuclei injected per unit time,  $\dot{N}_{\text{Fe}}$ , which can be simply related to the total power output of the pulsar  $L_{\text{Crab}}(B, P)$  [11],

$$\dot{N}_{\text{Fe}} = \xi L_{\text{Crab}}(B, P) / Z\Phi(B, P), \quad (1)$$

where  $\xi$  is the parameter describing the fraction of the total power taken by relativistic nuclei accelerated in the outer gap,  $Z = 26$  is the atomic number of Fe,  $B$  is the surface magnetic field,  $P$  is the pulsar's period, and

$$\Phi(B, P) \approx 5 \times 10^{16} \left( \frac{B}{4 \times 10^{12} \text{ G}} \right) \left( \frac{P}{1 \text{ s}} \right)^{4/3} \text{ V} \quad (2)$$

is the potential difference across the outer gap [4]. We assume  $B = 4 \times 10^{12} \text{ G}$ .

Soon after the supernova explosion, when the nebula was relatively small, nearly all energetic neutrons would be expected to decay outside the nebula. However, at early times we must take account of collisions with matter. The optical depth may be estimated from  $\tau_{nH} \approx \sigma_{pp} n_H r \approx 8.6 \times 10^{14} M_1 v_8^{-2} t^{-2}$ , where  $M = M_1 M_\odot$  is the mass ejected during the Crab supernova explosion in units of solar masses,  $r = vt$ ,  $n_H = M / (4/3 \pi r^3 m_p)$  is the number density of target nuclei, and  $v = 10^8 v_8 \text{ cm s}^{-1}$  is the expansion velocity of the nebula. Note that  $\tau_{nH} = 1$  at  $t \approx 0.93 M_1^{1/2} v_8^{-1} \text{ yr}$ , and that in this paper we consider only interactions in the ejecta and have neglected swept-up interstellar matter.

We estimate the evolution of the Crab pulsar's period from birth to the present time taking account of magnetic dipole radiation energy losses and gravitational energy losses for an ellipticity of  $3 \times 10^{-4}$  [12]. Magnetic dipole losses determine the pulsar period at present, but the initial period is determined largely by gravitational losses and is probably shorter than  $\sim 10 \text{ ms}$ . Hence, we consider two initial periods, 5 and 10 ms.

The spectrum of protons from neutron decay outside the Crab nebula is given by

$$N_p^{\text{out}}(\gamma_p, t_{\text{CN}}) = \int_0^{t_{\text{CN}}} dt \dot{N}_{\text{Fe}}(t) N_n(\gamma_p, t) \times e^{-\tau_{nH}(t)} e^{-vt_{\text{CN}}/c\gamma_p\tau_n}, \quad (3)$$

where  $\tau_n \approx 900 \text{ s}$  is the neutron decay time, and we make the approximation that the Lorentz factor of protons is equal to that of parent neutrons,  $\gamma_p \approx \gamma_n$ .

For the spectrum of protons injected inside the Crab nebula we must take account of proton-proton collisions and adiabatic energy losses due to the expansion of the nebula. The Lorentz factor of these protons at time  $t$  after the explosion such that their present Lorentz factor is  $\gamma_p$  is given by

$$\gamma_p(t) \approx \gamma_p \frac{(t + t_{\text{CN}})}{2tK\tau_{pp}(t)}, \quad (4)$$

where  $\tau_{pp}(t)$  is the optical depth for collision of protons with matter between  $t$  and  $t_{\text{CN}}$ , and is given by  $\tau_{pp}(t) \approx 1.3 \times 10^{17} M_1 v_8^{-3} (t^{-2} - t_{\text{CN}}^{-2})$ , and  $K \approx 0.5$  is the inelasticity coefficient in proton-proton collisions.

Since we are interested in protons interacting inside the nebula at the present time, we must also include those neutrons which decayed at locations outside the nebula at time  $t$  but which will be inside the nebula at time  $t_{\text{CN}}$ . Taking account of all these effects, we arrive at the formula below for the proton spectrum inside the nebula at time  $t_{\text{CN}}$ ,

$$N_p^{\text{in}}(\gamma_p, t_{\text{CN}}) = \gamma_p^{-1} \int_0^{t_{\text{CN}}} dt \dot{N}_{\text{Fe}}(t) e^{-\tau_{nH}(t)} \left\{ N_n[\gamma_p(t), t] \gamma_p(t) \{ 1 - \exp[-vt/c\gamma_p(t)\tau_n] \} + \int_t^{t_{\text{CN}}} dt' N_n[\gamma_p(t'), t] \gamma_p(t') \frac{v \exp[-vt'/c\gamma_p(t')\tau_n]}{c\gamma_p(t')\tau_n} \right\}. \quad (5)$$

The first term gives the contribution from neutrons decaying initially inside the nebula while the second term gives the contribution from neutrons decaying at points initially outside the nebula which will be inside the nebula at time  $t_{\text{CN}}$ . Using the minimum diffusion coefficient we estimate the typical diffusion distance during time  $t_{\text{CN}}$  for protons with  $\gamma_p = 10^5$  in a magnetic field of the order of  $B = 5 \times 10^{-6} \text{ G}$  to be comparable to the radius of the Crab nebula. Hence, in Eq. (5) we have assumed that protons from neutrons decaying inside the nebula will remain inside the nebula.

The diffusion distances discussed above also suggest that even protons from neutrons decaying initially outside the nebula may eventually be captured by the nebula as it expands, and this is what we have assumed in Eq. (5) (second term). However, it is possible that energetic protons from neutrons decaying outside the nebula may have exerted sufficient pressure on the surrounding gas such that it expanded rapidly together with the energetic proton gas, reducing the probability of capture of energetic protons by the nebula. If this occurred, it would have had the effect of reducing the  $\gamma$ -ray and neutrino fluxes, and we




FIG. 2. Spectra of protons from the decay of neutrons injected by the Crab pulsar inside and outside the Crab nebula assuming  $M_1 = 3$  for the following parameters:  $P_0 = 10$  ms, and Crab nebula radius  $r_{\text{CN}} = 1$  pc (dot-dashed curve) and 2 pc (dashed curve), and for  $P_0 = 5$  ms and  $r_{\text{CN}} = 1$  pc (full curve). The dotted curve shows the effect for  $P_0 = 5$  ms and  $r_{\text{CN}} = 1$  pc of no capture by the nebula of  $p$ 's from  $n$ 's decaying outside it.

shall investigate this possibility in an approximate way by neglecting the second term in Eq. (5).

We compute separately the spectra of protons from neutrons decaying inside and outside the Crab nebula. In Fig. 2 proton spectra are shown for two initial periods, 5 and 10 ms, and present nebula radii of 1 and 2 pc. Protons from neutrons decaying outside the present nebula radius will typically have high energies and diffuse rapidly, thus escaping to become galactic cosmic rays. In the case of  $r_{\text{CN}} = 1$  pc and  $P_0 = 5$  ms we also show the result for protons inside the nebula for the case where there is no capture of energetic protons during nebula expansion [i.e., neglecting the second term in Eq. (5)].

The protons which have accumulated inside the Crab nebula since the pulsar was formed (i.e., with spectra labeled “inside” in Fig. 2) can produce observable fluxes of  $\gamma$  rays and neutrinos. We compute the expected  $\gamma$ -ray spectra for five different models, taking various initial pulsar periods, present sizes, and masses of the Crab nebula, and whether or not there is capture of energetic protons as a result of nebula expansion. The fluxes may possibly be enhanced if protons are efficiently trapped by the dense filaments as suggested by Atoyan and Aharonian [13]. Filaments with density  $\sim 500 \text{ cm}^{-3}$  are present in the Crab nebula [14]. Therefore we introduce an effective density experienced by the protons inside nebula,  $n_H^{\text{eff}} = \mu n_H$ , where  $n_H$  is defined above, and the parameter  $\mu$  takes into account the possible effects of proton trapping by the filaments. The  $\gamma$ -ray spectra are computed in terms of a scaling model [15], and the photon fluxes expected at the Earth are shown in Fig. 3 for  $\xi\mu = 1$ , and are compared with observations of the Crab nebula above 0.2 TeV. Results are shown for models I to V having  $P_0$ ,  $r_{\text{CN}}$ , and  $M_1$  as specified in Table I, and for a distance to




FIG. 3. Spectra of  $\gamma$  rays from interactions of protons with matter inside the Crab nebula for the different proton spectra shown in Fig. 2 and for the different models of the nebula (I–V) considered in the text. Observations: Whipple Observatory [16] (dotted line and error box); THEMISTOCLE [17] (+); and CANGAROO [18] (solid line and error box). Upper limits from various experiments mentioned in Ref. [19]: T—Tibet, H—HEGRA, C—CYGNUS, and U—CASA-MIA.

the Crab nebula of 1830 pc [14]. Model V is the same as model I except that there is no capture of energetic protons by the nebula as it expands, and as expected the fluxes are lower than for model I. Comparison with the Whipple observations at 10 TeV allows us to place constraints on the free parameters of the model, and upper limits on  $\xi\mu$  which are given in Table I as  $(\xi\mu)_\gamma$ . Note that it is usually argued for the standard pulsar model that the rate of injection of Fe nuclei into the pulsar magnetosphere should not cause the charge density to exceed the Goldreich and Julian density [10]. In the case of no additional currents flowing through the magnetosphere, this condition constrains the value of  $\xi$  to  $\xi \approx 0.1$  for the Crab pulsar with period 33 ms. However, in the presence of additional currents the value of  $\xi$  can be higher.

The question of the importance of hadronic interactions in the Crab nebula can be settled by the detection of a neutrino signal from the Crab. In Fig. 4 we show the neutrino spectrum produced in collisions of protons with matter inside the Crab nebula and compare this with the atmospheric neutrino background flux within  $1^\circ$  of the source direction [20]. It is clear that neutrino detectors with good angular resolution should be able to detect

TABLE I. Model parameters and limits on  $\xi\mu$ .

Model	I	II	III	IV	V
$P_0$ (ms)	5	10	10	10	5
$r_{\text{CN}}$ (pc)	1	2	2	1	1
$M_1$	3	3	10	3	3
Nebula captures $p$ 's	yes	yes	yes	yes	no
$(\xi\mu)_\gamma$ at 10 TeV	0.63	6.9	2.0	1.0	3.2
$(\xi\mu)_\nu$ at 10 TeV	0.22	1.8	0.54	0.29	0.87




FIG. 4. Spectra of neutrinos ( $\nu_\mu + \bar{\nu}_\mu$ ) expected for the different models (I–V) considered in the text. The atmospheric neutrino background [20] within  $1^\circ$  of the source is indicated by the hatched band which shows the variation with angle: horizontal (upper bound), vertical (lower bound).

neutrinos at 10 TeV from the Crab nebula if  $\xi\mu$  is greater than  $(\xi\mu)_\nu$  given in Table I. Note that in all cases  $(\xi\mu)_\nu < (\xi\mu)_\gamma$  and so the possibility of  $\nu$  detection is allowed by the existing  $\gamma$ -ray observations.

In the present paper we have not considered those nuclei which had survived propagation through the outer gap radiation field and were injected into the inner nebula region following magnetic field lines. These nuclei (and protons) are expected to accumulate in the inner part of the Crab nebula where the magnetic fields are  $\sim 2 \times 10^{-3}$  G [21]. The diffusion distance during  $t_{CN}$  is  $\sim 0.3$  pc for  $\gamma_A = 10^6$ , and so most nuclei will probably be confined to the inner part of the Crab nebula where there is no evidence of dense matter [14] (other than the neutron star). Hence, we may be justified in neglecting the contribution of such nuclei to the  $\gamma$ -ray and neutrino production by hadronic collisions in comparison with the contribution from interactions of protons from neutron decay. However, some fraction of the energy of these nuclei could be transferred to very high energy electrons by resonant scattering [22]. Such electrons might then produce additional  $\gamma$  rays by the SSC process as in the models discussed in Ref. [2]. Hence, we suggest that the complex Crab nebula spectrum may, in fact, be formed as a result of both processes, i.e., SSC and as a result of pulsar acceleration of nuclei discussed here.

Another consequence of our model is that particles (nuclei and protons from neutron decay) will also be injected by the Crab pulsar and other pulsars into the galactic cosmic rays; this may affect the cosmic ray spectrum and composition. It is worth noting that the energy distribution of the protons from decay of neutrons outside the Crab nebula peaks at  $\sim 10^{15}$  eV (see Fig. 2), which is close to the energy of the knee in the cosmic ray spectrum. This is particularly interesting as the possibility of a single source

of cosmic rays in the range  $10^{15}$  to  $10^{16}$  eV has been discussed recently by Erlykin and Wolfendale [23].

W. B. thanks the University of Adelaide for hospitality during his visit. This research is supported by a grant from the Australian Research Council.

\*Permanent address: University of Łódź, 90-236 Łódź, ul. Pomorska 149/153, Poland.

- [1] P. L. Nolan *et al.*, *Astrophys. J.* **409**, 697 (1993); R. P. Much *et al.*, *Astron. Astrophys.* **299**, 435 (1995); O. C. de Jager *et al.*, *Astrophys. J.* **457**, 253 (1996); T. C. Weekes *et al.*, *Astrophys. J.* **342**, 379 (1989); G. Vacanti *et al.*, *Astrophys. J.* **337**, 467 (1991).
- [2] R. J. Gould, *Phys. Rev. Lett.* **15**, 577 (1965); G. H. Rieke and T. C. Weekes, *Astrophys. J.* **155**, 429 (1969); J. E. Grindlay and J. A. Hoffman, *Astrophys. Lett.* **8**, 209 (1971); W. Kundt and E. Krotschek, *Astron. Astrophys.* **83**, 1 (1980); O. C. de Jager and A. K. Harding, *Astrophys. J.* **396**, 161 (1992); F. Aharonian and A. Atoyan, *Astropart. Phys.* **3**, 275 (1995).
- [3] W. M. Kwok, K. S. Cheng, and M. M. Lau, *Astrophys. J.* **379**, 653 (1991); W. M. Cheung and K. S. Cheng, *Astrophys. J.* **413**, 694 (1993).
- [4] K. S. Cheng, T. Cheung, M. M. Lau, K. N. Yu, and W. M. Kwok, *J. Phys. G* **16**, 1115 (1990).
- [5] S. Karakuła and W. Tkaczyk, *Astropart. Phys.* **1**, 229 (1993).
- [6] C. Ho, *Astrophys. J.* **342**, 396 (1989).
- [7] K. S. Cheng, C. Ho, and M. Ruderman, *Astrophys. J.* **300**, 500 (1986); **300**, 522 (1986).
- [8] F. W. Stecker, *Phys. Rev. Lett.* **21**, 1016 (1968).
- [9] M. J. Chodorowski, A. A. Zdziarski, and M. Sikora, *Astrophys. J.* **400**, 181 (1992).
- [10] P. Goldreich and W. H. Julian, *Astrophys. J.* **157**, 869 (1969).
- [11] R. N. Manchester and J. H. Taylor, *Pulsars* (Freeman, San Francisco, 1977).
- [12] S. L. Shapiro and S. L. Teukolsky, *Black Holes, White Dwarfs and Neutron Stars* (Wiley, New York, 1983).
- [13] A. M. Atoyan and F. A. Aharonian, *Mon. Not. R. Astron. Soc.* **278**, 525 (1996).
- [14] K. Davidson and R. A. Fesen, *Annu. Rev. Astron. Astrophys.* **23**, 119 (1985).
- [15] A. H. Hillas, *Proc. 17th ICRC (Paris)* **8**, 193 (1981).
- [16] D. A. Lewis *et al.*, *Proc. 23rd ICRC (Dublin)* **1**, 279 (1993).
- [17] THEMISTOCLE Collaboration, *Proc. 24th ICRC (Rome)* **2**, 315 (1995).
- [18] T. Tanimori *et al.* (to be published).
- [19] M. Amenomori *et al.*, *Proc. 24th ICRC (Rome)* **2**, 346 (1995).
- [20] P. Lipari, *Astropart. Phys.* **1**, 195 (1993).
- [21] J. J. Hester *et al.*, *Astrophys. J.* **448**, 240 (1995).
- [22] M. Hoshino, J. Arons, Y. A. Gallant, and A. B. Langdon, *Astrophys. J.* **390**, 454 (1992); Y. A. Gallant and J. Arons, *Astrophys. J.* **435**, 230 (1994).
- [23] A. D. Erlykin and A. W. Wolfendale, *J. Phys. G* **8**, 979 (1997).