

Search for Flavor-Changing Neutral Current Decays of the Top Quark in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV

F. Abe,¹⁷ H. Akimoto,³⁶ A. Akopian,³¹ M. G. Albrow,⁷ S. R. Amendolia,²⁷ D. Amidei,²⁰ J. Antos,³³ S. Aota,³⁶ G. Apollinari,³¹ T. Asakawa,³⁶ W. Ashmanskas,¹⁸ M. Atac,⁷ F. Azfar,²⁶ P. Azzi-Bacchetta,²⁵ N. Bacchetta,²⁵ W. Badgett,²⁰ S. Bagdasarov,³¹ M. W. Bailey,²² J. Bao,³⁹ P. de Barbaro,³⁰ A. Barbaro-Galtieri,¹⁸ V. E. Barnes,²⁹ B. A. Barnett,¹⁵ M. Barone,⁹ E. Barzi,⁹ G. Bauer,¹⁹ T. Baumann,¹¹ F. Bedeschi,²⁷ S. Behrends,³ S. Belforte,²⁷ G. Bellettini,²⁷ J. Bellinger,³⁸ D. Benjamin,³⁵ J. Benlloch,¹⁹ J. Bensinger,³ D. Benton,²⁶ A. Beretvas,⁷ J. P. Berge,⁷ J. Berryhill,⁵ S. Bertolucci,⁹ S. Bettelli,²⁷ B. Bevensee,²⁶ A. Bhatti,³¹ K. Biery,⁷ M. Binkley,⁷ D. Bisello,²⁵ R. E. Blair,¹ C. Blocker,³ A. Bodek,³⁰ W. Bokhari,¹⁹ V. Bolognesi,² G. Bolla,²⁹ D. Bortoletto,²⁹ J. Boudreau,²⁸ L. Breccia,² C. Bromberg,²¹ N. Bruner,²² E. Buckley-Geer,⁷ H. S. Budd,³⁰ K. Burkett,²⁰ G. Busetto,²⁵ A. Byon-Wagner,⁷ K. L. Byrum,¹ J. Cammerata,¹⁵ C. Campagnari,⁷ M. Campbell,²⁰ A. Caner,²⁷ W. Carithers,¹⁸ D. Carlsmith,³⁸ A. Castro,²⁵ D. Cauz,²⁷ Y. Cen,³⁰ F. Cervelli,²⁷ P. S. Chang,³³ P. T. Chang,³³ H. Y. Chao,³³ J. Chapman,²⁰ M.-T. Cheng,³³ M. Chertok,³⁴ G. Chiarelli,²⁷ T. Chikamatsu,³⁶ C. N. Chiou,³³ L. Christofek,¹³ S. Cihangir,⁷ A. G. Clark,¹⁰ M. Cobal,²⁷ E. Cocca,²⁷ M. Contreras,⁵ J. Conway,³² J. Cooper,⁷ M. Cordelli,⁹ C. Couyoumtzelis,¹⁰ R. L. Coxe,¹¹ D. Crane,¹ D. Cronin-Hennessy,⁶ R. Culbertson,⁵ T. Daniels,¹⁹ F. DeJongh,⁷ S. Delchamps,⁷ S. Dell'Agnello,²⁷ M. Dell'Orso,²⁷ R. Demina,⁷ L. Demortier,³¹ M. Deninno,² P. F. Derwent,⁷ T. Devlin,³² J. R. Dittmann,⁶ S. Donati,²⁷ J. Done,³⁴ T. Dorigo,²⁵ A. Dunn,²⁰ N. Eddy,²⁰ K. Einsweiler,¹⁸ J. E. Elias,⁷ R. Ely,¹⁸ E. Engels, Jr.,²⁸ D. Errede,¹³ S. Errede,¹³ Q. Fan,³⁰ G. Feild,³⁹ C. Ferretti,²⁷ I. Fiori,² B. Flaughner,⁷ G. W. Foster,⁷ M. Franklin,¹¹ M. Frautschi,³⁵ J. Freeman,⁷ J. Friedman,¹⁹ H. Frisch,⁵ Y. Fukui,¹⁷ S. Funaki,³⁶ S. Galeotti,²⁷ M. Gallinaro,²⁶ O. Ganel,³⁵ M. Garcia-Sciveres,¹⁸ A. F. Garfinkel,²⁹ C. Gay,¹¹ S. Geer,⁷ D. W. Gerdes,¹⁵ P. Giannetti,²⁷ N. Giokaris,³¹ P. Giromini,⁹ G. Giusti,²⁷ L. Gladney,²⁶ D. Glenzinski,¹⁵ M. Gold,²² J. Gonzalez,²⁶ A. Gordon,¹¹ A. T. Goshaw,⁶ Y. Gotra,²⁵ K. Goulianos,³¹ H. Grassmann,²⁷ L. Groer,³² C. Grosso-Pilcher,⁵ G. Guillian,²⁰ R. S. Guo,³³ C. Haber,¹⁸ E. Hafen,¹⁹ S. R. Hahn,⁷ R. Hamilton,¹¹ R. Handler,³⁸ R. M. Hans,³⁹ F. Happacher,⁹ K. Hara,³⁶ A. D. Hardman,²⁹ B. Harral,²⁶ R. M. Harris,⁷ S. A. Hauger,⁶ J. Hauser,⁴ C. Hawk,³² E. Hayashi,³⁶ J. Heinrich,²⁶ B. Hinrichsen,¹⁴ K. D. Hoffman,²⁹ M. Hohlmann,⁵ C. Holck,²⁶ R. Hollebeek,²⁶ L. Holloway,¹³ S. Hong,²⁰ G. Houk,²⁶ P. Hu,²⁸ B. T. Huffman,²⁸ R. Hughes,²³ J. Huston,²¹ J. Huth,¹¹ J. Hysten,⁷ H. Ikeda,³⁶ M. Incagli,²⁷ J. Incandela,⁷ G. Introzzi,²⁷ J. Iwai,³⁶ Y. Iwata,¹² H. Jensen,⁷ U. Joshi,⁷ R. W. Kadel,¹⁸ E. Kajfasz,²⁵ H. Kambara,¹⁰ T. Kamon,³⁴ T. Kaneko,³⁶ K. Karr,³⁷ H. Kasha,³⁹ Y. Kato,²⁴ T. A. Keaffaber,²⁹ K. Kelley,¹⁹ R. D. Kennedy,⁷ R. Kephart,⁷ P. Kesten,¹⁸ D. Kestenbaum,¹¹ H. Keutelian,⁷ F. Keyvan,⁴ B. Kharadia,¹³ B. J. Kim,³⁰ D. H. Kim,^{7,*} H. S. Kim,¹⁴ S. B. Kim,²⁰ S. H. Kim,³⁶ Y. K. Kim,¹⁸ L. Kirsch,³ P. Koehn,²³ K. Kondo,³⁶ J. Konigsberg,⁸ S. Kopp,⁵ K. Kordas,¹⁴ A. Korytov,⁸ W. Koska,⁷ E. Kovacs,^{7,*} W. Kowald,⁶ M. Krasberg,²⁰ J. Kroll,⁷ M. Kruse,³⁰ T. Kuwabara,³⁶ S. E. Kuhlmann,¹ E. Kuns,³² A. T. Laasanen,²⁹ S. Lami,²⁷ S. Lammel,⁷ J. I. Lamoureux,³ M. Lancaster,¹⁸ M. Lanzoni,²⁷ G. Latino,²⁷ T. LeCompte,¹ S. Leone,²⁷ J. D. Lewis,⁷ P. Limon,⁷ M. Lindgren,⁴ T. M. Liss,¹³ J. B. Liu,³⁰ Y. C. Liu,³³ N. Lockyer,²⁶ O. Long,²⁶ C. Loomis,³² M. Loreti,²⁵ J. Lu,³⁴ D. Lucchesi,²⁷ P. Lukens,⁷ S. Lusin,³⁸ J. Lys,¹⁸ K. Maeshima,⁷ A. Maghakian,³¹ P. Maksimovic,¹⁹ M. Mangano,²⁷ J. Mansour,²¹ M. Mariotti,²⁵ J. P. Marriner,⁷ A. Martin,³⁹ J. A. J. Matthews,²² R. Mattingly,¹⁹ P. Mazzanti,² P. McIntyre,³⁴ P. Melese,³¹ A. Menzione,²⁷ E. Meschi,²⁷ S. Metzler,²⁶ C. Miao,²⁰ T. Miao,⁷ G. Michail,¹¹ R. Miller,²¹ H. Minato,³⁶ S. Miscetti,⁹ M. Mishina,¹⁷ H. Mitsushio,³⁶ T. Miyamoto,³⁶ S. Miyashita,³⁶ N. Moggi,²⁷ Y. Morita,¹⁷ A. Mukherjee,⁷ T. Muller,¹⁶ P. Murat,²⁷ H. Nakada,³⁶ I. Nakano,³⁶ C. Nelson,⁷ D. Neuberger,¹⁶ C. Newman-Holmes,⁷ C.-Y. P. Ngan,¹⁹ M. Ninomiya,³⁶ L. Nodulman,¹ S. H. Oh,⁶ K. E. Ohl,³⁹ T. Ohmoto,¹² T. Ohsugi,¹² R. Oishi,³⁶ M. Okabe,³⁶ T. Okusawa,²⁴ R. Oliveira,²⁶ J. Olsen,³⁸ C. Pagliarone,²⁷ R. Paoletti,²⁷ V. Papadimitriou,³⁵ S. P. Pappas,³⁹ N. Parashar,²⁷ S. Park,⁷ A. Parri,⁹ J. Patrick,⁷ G. Pauletta,²⁷ M. Paulini,¹⁸ A. Perazzo,²⁷ L. Pescara,²⁵ M. D. Peters,¹⁸ T. J. Phillips,⁶ G. Piacentino,²⁷ M. Pillai,³⁰ K. T. Pitts,⁷ R. Plunkett,⁷ L. Pondrom,³⁸ J. Proudfoot,¹ F. Ptohos,¹¹ G. Punzi,²⁷ K. Ragan,¹⁴ D. Reher,¹⁸ A. Ribon,²⁵ F. Rimondi,² L. Ristori,²⁷ W. J. Robertson,⁶ T. Rodrigo,²⁷ S. Rolli,³⁷ J. Romano,⁵ L. Rosenson,¹⁹ R. Roser,¹³ T. Saab,¹⁴ W. K. Sakumoto,³⁰ D. Saltzberg,⁵ A. Sansoni,⁹ L. Santi,²⁷ H. Sato,³⁶ P. Schlabach,⁷ E. E. Schmidt,⁷ M. P. Schmidt,³⁹ A. Scribano,²⁷ S. Segler,⁷ S. Seidel,²² Y. Seiya,³⁶ F. Semeria,² G. Sganos,¹⁴ M. D. Shapiro,¹⁸ N. M. Shaw,²⁹ Q. Shen,²⁹ P. F. Shepard,²⁸ M. Shimojima,³⁶ M. Shochet,⁵ J. Siegrist,¹⁸ A. Sill,³⁵ P. Sinervo,¹⁴ P. Singh,²⁸ J. Skarha,¹⁵ K. Sliwa,³⁷ F. D. Snider,¹⁵ T. Song,²⁰ J. Spalding,⁷ T. Speer,¹⁰ P. Sphicas,¹⁹ F. Spinella,²⁷ M. Spiropulu,¹¹ L. Spiegel,⁷ L. Stanco,²⁵ J. Steele,³⁸ A. Stefanini,²⁷ K. Strahl,¹⁴ J. Strait,⁷ R. Ströhmer,^{7,*} D. Stuart,⁷ G. Sullivan,⁵ K. Sumorok,¹⁹ J. Suzuki,³⁶ T. Takada,³⁶ T. Takahashi,²⁴ T. Takano,³⁶

K. Takikawa,³⁶ N. Tamura,¹² B. Tannenbaum,²² F. Tartarelli,²⁷ W. Taylor,¹⁴ P. K. Teng,³³ Y. Teramoto,²⁴ S. Tether,¹⁹ D. Theriot,⁷ T. L. Thomas,²² R. Thun,²⁰ R. Thurman-Keup,¹ M. Timko,³⁷ P. Tipton,³⁰ A. Titov,³¹ S. Tkaczyk,⁷ D. Toback,⁵ K. Tollefson,³⁰ A. Tollestrup,⁷ H. Toyoda,²⁴ W. Trischuk,¹⁴ J. F. de Troconiz,¹¹ S. Truitt,²⁰ J. Tseng,¹⁹ N. Turini,²⁷ T. Uchida,³⁶ N. Uemura,³⁶ F. Ukegawa,²⁶ G. Unal,²⁶ J. Valls,^{7,*} S. C. van den Brink,²⁸ S. Vajcik III,²⁰ G. Velev,²⁷ R. Vidal,⁷ R. Vilar,^{7,*} M. Vondracek,¹³ D. Vucinic,¹⁹ R. G. Wagner,¹ R. L. Wagner,⁷ J. Wahl,⁵ N. B. Wallace,²⁷ A. M. Walsh,³² C. Wang,⁶ C. H. Wang,³³ J. Wang,⁵ M. J. Wang,³³ Q. F. Wang,³¹ A. Warburton,¹⁴ T. Watts,³² R. Webb,³⁴ C. Wei,⁶ H. Wenzel,¹⁶ W. C. Wester III,⁷ A. B. Wicklund,¹ E. Wicklund,⁷ R. Wilkinson,²⁶ H. H. Williams,²⁶ P. Wilson,⁵ B. L. Winer,²³ D. Winn,²⁰ D. Wolinski,²⁰ J. Wolinski,²¹ S. Worm,²² X. Wu,¹⁰ J. Wyss,²⁵ A. Yagil,⁷ W. Yao,¹⁸ K. Yasuoka,³⁶ Y. Ye,¹⁴ G. P. Yeh,⁷ P. Yeh,³³ M. Yin,⁶ J. Yoh,⁷ C. Yosef,²¹ T. Yoshida,²⁴ D. Yovanovitch,⁷ I. Yu,²² L. Yu,²² J. C. Yun,⁷ A. Zanetti,²⁷ F. Zetti,²⁷ L. Zhang,³⁸ W. Zhang,²⁶ and S. Zucchelli²

(CDF Collaboration)

¹Argonne National Laboratory, Argonne, Illinois 60439

²Istituto Nazionale di Fisica Nucleare, University of Bologna, I-40127 Bologna, Italy

³Brandeis University, Waltham, Massachusetts 02264

⁴University of California at Los Angeles, Los Angeles, California 90024

⁵University of Chicago, Chicago, Illinois 60637

⁶Duke University, Durham, North Carolina 28708

⁷Fermi National Accelerator Laboratory, Batavia, Illinois 60510

⁸University of Florida, Gainesville, Florida 33611

⁹Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy

¹⁰University of Geneva, CH-1211 Geneva 4, Switzerland

¹¹Harvard University, Cambridge, Massachusetts 02138

¹²Hiroshima University, Higashi-Hiroshima 724, Japan

¹³University of Illinois, Urbana, Illinois 61801

¹⁴Institute of Particle Physics, McGill University, Montreal H3A 2T8
and University of Toronto, Toronto M5S 1A7, Canada

¹⁵The Johns Hopkins University, Baltimore, Maryland 21218

¹⁶Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany

¹⁷National Laboratory for High Energy Physics (KEK), Tsukuba, Ibaraki 315, Japan

¹⁸Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720

¹⁹Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

²⁰University of Michigan, Ann Arbor, Michigan 48109

²¹Michigan State University, East Lansing, Michigan 48824

²²University of New Mexico, Albuquerque, New Mexico 87132

²³The Ohio State University, Columbus, Ohio 43220

²⁴Osaka City University, Osaka 588, Japan

²⁵Università di Padova, Istituto Nazionale di Fisica Nucleare, Sezione di Padova, I-36132 Padova, Italy

²⁶University of Pennsylvania, Philadelphia, Pennsylvania 19104

²⁷Istituto Nazionale di Fisica Nucleare, University and Scuola Normale Superiore of Pisa, I-56100 Pisa, Italy

²⁸University of Pittsburgh, Pittsburgh, Pennsylvania 15270

²⁹Purdue University, West Lafayette, Indiana 47907

³⁰University of Rochester, Rochester, New York 14628

³¹Rockefeller University, New York, New York 10021

³²Rutgers University, Piscataway, New Jersey 08854

³³Academia Sinica, Taipei, Taiwan 11530, Republic of China

³⁴Texas A&M University, College Station, Texas 77843

³⁵Texas Tech University, Lubbock, Texas 79409

³⁶University of Tsukuba, Tsukuba, Ibaraki 315, Japan

³⁷Tufts University, Medford, Massachusetts 02155

³⁸University of Wisconsin, Madison, Wisconsin 53806

³⁹Yale University, New Haven, Connecticut 06511

(Received 24 July 1997)

We search for the flavor-changing neutral current decays of the top quark $t \rightarrow q\gamma$ and $t \rightarrow qZ$ (here q represents the c and u quarks) in $\bar{p}p$ collisions at $\sqrt{s} = 1.8$ TeV. We use a dataset ($\int \mathcal{L} dt \sim 110$ pb⁻¹) collected during the 1992–1995 run of the Collider Detector at Fermilab. We set 95% confidence level limits on the branching fractions $B(t \rightarrow q\gamma) < 3.2\%$ and $B(t \rightarrow qZ) < 33\%$, consistent with the standard model. [S0031-9007(98)05627-0]

PACS numbers: 12.15.Mm, 14.65.Ha

Physics beyond the standard model can manifest itself by altering the expected rates of flavor-changing neutral current (FCNC) interactions. As an analogous historical example within the standard model, the presence of the charm quark can be inferred from its effect on the branching fraction $B(K_L^0 \rightarrow \mu^+ \mu^-)$ [1]. FCNC decays of the top quark are of particular interest [2,3]. The large mass of the top quark suggests a strong connection with the electroweak symmetry breaking sector. Evidence for unusual decays of the top quark might provide insights into the mechanism of electroweak symmetry breaking. For the top quark, the FCNC decays $t \rightarrow qZ$ and $t \rightarrow q\gamma$ [4] (where q denotes either a c or a u flavored quark) are expected to be exceedingly rare (branching fractions of 10^{-10} or smaller) [5] and any observation of these decays in the presently available data would indicate new physics.

We present results using data from proton-antiproton collisions at a center of mass energy of 1.8 TeV collected at the Collider Detector at Fermilab (CDF) during the 1992–1995 run of the Fermilab Tevatron ($\int \mathcal{L} dt \sim 110 \text{ pb}^{-1}$). The CDF detector is described in detail elsewhere [6]; a brief discussion follows.

In the CDF detector, a 51 cm long silicon vertex detector (SVX) [7], located immediately outside the beam pipe, provides precise track reconstruction in the plane transverse to the beam, and is used to identify secondary vertices that can be produced by b and c quark decays. Because $p\bar{p}$ interactions are spread along the beam line with a standard deviation of about 30 cm, slightly more than half of the events originate from primary vertices inside the SVX fiducial region. The momentum of charged particles is measured in the central tracking chamber (CTC), which sits inside a 1.4 T superconducting solenoidal magnet. Outside the CTC are electromagnetic and hadronic calorimeters, arranged in a projective tower geometry, covering the pseudorapidity region $|\eta| < 4.2$ [8]. In the central ($|\eta| < 1.0$) electromagnetic calorimeter, finely segmented proportional chambers (CES) used to measure transverse shower profiles are placed at a depth of approximately six radiation lengths. Jet photon and electron candidates are identified in the calorimeters, as is the missing transverse energy \cancel{E}_T [9]. Surrounding the calorimeters, drift chambers in the region $|\eta| < 1.0$ provide muon identification. A three level trigger selects events with high transverse momentum electrons, muons and/or photons for this analysis.

At CDF, a photon is identified as an energy cluster in the electromagnetic calorimeter with no track pointing at it. To improve identification efficiency, we additionally permit one single soft track (presumably from a random overlap) with less than 10% of the energy of the photon to point at the cluster. For the typical photon shower, the electromagnetic cluster consists of two adjacent calorimeter towers. The energy of the photon is measured in the calorimeter, and the direction of the photon is defined by a line between the event vertex and the centroid of the

electromagnetic shower as measured in the CES. To reduce backgrounds from hadronic jets, we require that the total energy (electromagnetic plus hadronic) in a cone of $\Delta R = 0.4$ (where $\Delta R \equiv \sqrt{\Delta\eta^2 + \Delta\phi^2}$) around and excluding the photon candidate be less than 15% of the photon energy. To reduce the background from neutral hadrons, the ratio of energy deposited in the hadronic calorimeter to that in the electromagnetic calorimeter is required to be less than $0.055 + 0.00045E$, where E is the total energy of the candidate in GeV. To suppress π^0 and multiphoton backgrounds, the transverse shower profile in the CES and the energy sharing between the calorimeter towers must be consistent with a single electromagnetic shower.

An electron is also identified as an energy cluster in the electromagnetic calorimeter, this time with a single track pointing to it. The energy of the cluster and the momentum of the track are required to be equal within measurement uncertainties, and the extrapolation of the track to the CES is required to match the measured position of the shower in the CES. Only electrons in the central region ($|\eta| < 1.0$) are used in this analysis.

A muon is identified by requiring a match between the extrapolated track as reconstructed in the CTC and track segments reconstructed in the muon chambers, taking into account multiple scattering of the muon. Furthermore, the energy deposition in the calorimeter must be consistent with a minimum ionizing particle. In this analysis, we restrict ourselves to muons in the central region ($|\eta| < 1.0$). Further details of the electron and muon identification requirements can be found in [10].

A jet is identified by energy deposited in the calorimeters in a cone of $R < 0.4$. Details of the CDF jet reconstruction algorithm can be found in [11].

If the branching fraction of a particle into a particular final state (e.g., a FCNC decay) is x , the branching fraction into another final state (e.g., the decay $t \rightarrow Wb$ [12]) can be no larger than $(1 - x)$. Therefore, the ratio r of the number of events detected in a rare decay mode normalized to a common decay mode is at least $x/(1 - x)$, after corrections for efficiency and acceptance. Measuring r allows us to calculate an upper limit on the branching fraction $x \leq r/(1 + r)$, after corrections for misidentification, efficiency, and acceptance, and for the fact that in $t\bar{t}$ events there are two top quarks that can decay into a particular final state.

In both the $t \rightarrow q\gamma$ and $t \rightarrow qZ$ searches, we calculate limits on branching fractions by comparing the number of candidate events in the FCNC candidate samples to the number of $t\bar{t}$ events observed in the normalization sample. Our normalization sample consists of events consistent with the hypothesis where both top quarks decayed via $t \rightarrow Wb$ decays, one W decayed leptonically, and the other W decayed hadronically. These events are identified by a high p_T ($p_T > 20 \text{ GeV}/c$) electron or muon, at least 20 GeV of missing transverse energy from the undetected neutrino, and three or more jets above 15 GeV, at least

one of which must be identified as containing a b hadron from the presence of a secondary vertex [10]. This procedure is called “ b tagging” the jet. In a sample of 110 pb^{-1} , we observe 34 $t\bar{t}$ candidates over an estimated background of 9 ± 1.5 events.

At Tevatron energies, the dominant source of top quarks is $t\bar{t}$ pair production from $q\bar{q}$ annihilation. In the search for the decay $t \rightarrow q\gamma$, we assume that the other top quark in the pair decays via the decay $t \rightarrow Wb$. We consider two event signatures, depending on whether the W decayed leptonically or hadronically. If the W decayed leptonically, we search for events with an identified W (via a lepton with $p_T > 20 \text{ GeV}/c$ and 20 GeV of missing transverse energy carried by the neutrino), a moderately high E_T photon ($E_T > 20 \text{ GeV}$), and at least two jets with $E_T > 15 \text{ GeV}$. If the W decayed hadronically, we search for events with at least four jets (two from the W decay), and a high E_T ($E_T \geq 50 \text{ GeV}$) photon. One jet must contain a secondary vertex, identifying it as containing a b hadron. The higher E_T requirement on the photon is to reduce the background; events with jets are more common than events with high p_T leptons. In both cases, there must be a photon and jet combination with mass between $140\text{--}210 \text{ GeV}/c^2$, consistent with the mass of the top quark. In the nonleptonic case, the remaining jets must have $\Sigma E_T \geq 140 \text{ GeV}$, consistent with the decay of a second top quark in the event. The b -tagged jet must be associated with the second top combination. 40% of our $t \rightarrow q\gamma$ acceptance is in our photon plus multijet mode, and 60% is in the lepton plus photon mode.

We expect the background in the leptonic mode to be dominated by $W + \gamma + 2$ or more jet events. Barger *et al.* [13] calculate the rate of both $W + \gamma + 2$ jet production and $W + \gamma$ production at Tevatron energies. From that ratio and the number of observed $W + \gamma$ events at CDF, we estimate the number of background events, and find it to be less than half an event. In the hadronic mode, we can estimate the background by studying a sample of events that passes all selection requirements except for the identification of a jet containing a b hadron. From the number of events in this sample, plus the measured probability of misidentifying a jet as containing a b hadron [10,14], we estimate that the expected background is smaller than half an event. To set a conservative limit, we assume any events passing the selection requirements are signal and do not subtract any background.

We have defined two samples, one in which the event topology is consistent with both top quarks decaying via $t \rightarrow Wb$ (the “standard model sample”) and one in which the event topology is consistent with one top quark decaying via $t \rightarrow Wb$ and the other via $t \rightarrow c\gamma$ (the “FCNC sample”). Because we are interested in the relative number of events with these two signatures, we calculate the ratio of acceptances and efficiencies in the two decay modes. To do this, we use the ISAJET [15] Monte

TABLE I. Efficiencies and acceptances for $t \rightarrow q\gamma$ decays, normalized to standard model top decays. The first uncertainty reflects the uncertainty for all sources except the b quark identification efficiency; the second uncertainty reflects the uncertainty from the b quark identification efficiency. W branching fractions are included as part of the efficiency.

Generated $t\bar{t}$ decay	Identified as	Relative efficiency
$WbWb$	standard model	1 (by definition)
$Wb\gamma q$	standard model	$0.266 \pm 0.013 \pm 0.005$
$Wb\gamma q$	FCNC	$3.96 \pm 0.17 \mp 0.20$
$WbWb$	FCNC	negligible
$\gamma q\gamma q$	standard model	negligible
$\gamma q\gamma q$	FCNC	negligible

Carlo event generator and a parametric simulation of the CDF detector. Results are shown (relative to the standard model $WbWb$ signature, and including W branching fractions) in Table I. The first uncertainty is the uncertainty in the relative acceptance; the second is the effect of the b -tagging efficiency of $(45 \pm 5)\%$ per b jet contained in the SVX fiducial region. Positive (negative) uncertainties indicate that the change in relative efficiency is in the same (opposite) direction as the change in b -tagging efficiency. We also calculate how often events are misidentified (standard model events identified as FCNC events and vice versa) and incorporate this into the limit [16]. To be conservative, we do not subtract off the expected number of FCNC candidates from misidentified standard model decays. In 110 pb^{-1} of data, one event is observed in the leptonic channel, and no events are seen in the nonleptonic (i.e., photon plus multijet) channel.

The single event that passes all selection requirements has a 72 GeV muon, an 88 GeV photon, three jets, and a missing transverse energy of 24 GeV. While it enters the FCNC decay candidate sample, this event is also kinematically consistent with the decay $t \rightarrow W^+b$, $\bar{t} \rightarrow W^- \bar{b}\gamma$, followed by $W^+ \rightarrow \mu^+ \nu$ and $W^- \rightarrow$ jets. However, the photon E_T is exceptionally large for this decay.

Observation of one event passing the selection requirements implies a 95% confidence limit of fewer than 6.45 events (including systematic uncertainties) which translates into a branching fraction limit of

$$B(t \rightarrow c\gamma) + B(t \rightarrow u\gamma) < 3.2\%.$$

The statistical uncertainty in the number of events in the normalization sample is the dominant source of systematic uncertainty, increasing the limit by approximately 16%. The uncertainty in b quark identification efficiency also contributes to the overall systematic uncertainty, adding approximately 10%. The uncertainties on acceptance and photon identification are negligible by comparison.

We also search for $t \rightarrow qZ$ events, using the channel where the Z decays to e^+e^- or $\mu^+\mu^-$, and the other t quark decays to three jets. (Allowing the other t quark to decay to a lepton, neutrino, and jet does not substantially increase the acceptance, and so does not substantially

TABLE II. Efficiencies and acceptances for $t \rightarrow Zq$ decays, normalized to standard model top decays. The first uncertainty reflects the uncertainty for all sources except the b quark identification efficiency; the second uncertainty reflects the uncertainty from the b quark identification efficiency. W and Z branching fractions are included as part of the efficiency.

Generated $t\bar{t}$ decay	Identified as	Relative efficiency
$WbWb$	standard model	1 (by definition)
$WbZq$	standard model	$0.693 \pm 0.035 \mp 0.027$
$WbZq$	FCNC	$0.359 \pm 0.018 \mp 0.044$
$WbWb$	FCNC	$0.022 \pm 0.004 \pm 0.003$
$ZqZq$	standard model	$0.062 \pm 0.003 \pm 0.000$
$ZqZq$	FCNC	$0.630 \pm 0.032 \mp 0.077$

improve the limit.) The expected signature is, therefore, an event with four jets and with two leptons with an invariant mass consistent with a Z boson. Because the Z branching fraction to charged leptons is small, this analysis is less sensitive than the $t \rightarrow q\gamma$ search. Candidate $Z \rightarrow l^+l^-$ events were selected from inclusive samples of events with electron and muon candidates selected by the trigger ($\int \mathcal{L} dt = 108 \text{ pb}^{-1}$) using criteria described in detail in [17]. Z bosons are identified as opposite-charge same-flavor lepton pairs inside the range $75 < M_{l^+l^-} < 105 \text{ GeV}/c^2$. We require each of the four jets to have $E_T > 20 \text{ GeV}$ and be contained in the region $|\eta| < 2.4$.

As before, we use the ISAJET Monte Carlo event generator and a parametric simulation of the CDF detector to calculate the efficiencies and acceptances for top quark pairs to be identified as either standard model decay candidates or FCNC decay candidates. Again, we determine the acceptances and efficiencies (including misidentification probabilities) relative to the standard model signal (Table II). The first uncertainty is the uncertainty in the relative acceptance; the second is the effect of the b -tagging efficiency. W and Z branching fractions are included as part of the efficiencies.

There are two comparable sources of background to the $t \rightarrow qZ$ signal. One is ordinary $Z +$ multijet production; 0.5 background $Z + 4$ jet events are expected. The second source of background is from $t\bar{t}$ events where both top quarks decay via $t \rightarrow Wb$, and the event topology is such that it is accepted into this sample. One example would be where both W bosons decay leptonically, the two leptons have an invariant mass that falls within the Z candidate range, and there are two jets produced via gluon radiation. Simulations using the ISAJET Monte Carlo lead us to expect approximately 0.6 events in this category. A smaller source of background is diboson (WZ or ZZ) + 2 or more jet events in which the Z decays leptonically (0.1 events). The total background is approximately 1.2 events. As before, to set a conservative limit, we assume any events passing the selection requirements are signal and do not subtract this background.

A single $Z \rightarrow \mu^+\mu^-$ event passes all the selection requirements. The event kinematics better fit the $Z +$

multijet hypothesis than the FCNC decay hypothesis. Two of the jets have low transverse energies, a feature more common to $Z +$ multijet decays than top quark decays.

Observation of one event passing the selection requirements implies a 95% confidence limit of fewer than 6.4 events (including systematic uncertainties), which translates to a branching fraction limit

$$B(t \rightarrow cZ) + B(t \rightarrow uZ) < 33\%.$$

As before, the systematic uncertainties are dominated by the statistical uncertainty in the number of events in the normalization sample (16%), followed by the uncertainty in b -tagging efficiency (10%). Additional contributions to the uncertainties are caused by uncertainties in initial and final state gluon radiation and uncertainties in the jet energy scale (contributing in total 10%), which affects the efficiency for an event to pass the minimum jet energy requirements.

In summary, we search for the flavor-changing neutral current decays $t \rightarrow q\gamma$ and $t \rightarrow qZ$ in $\bar{p}p$ collisions at $\sqrt{s} = 1.8 \text{ TeV}$. No statistically significant excess of these events is seen, although one candidate event survives all cuts in each search. Both events have interpretations outside of the FCNC decay hypothesis. In order to set conservative limits, we treat these events as FCNC candidates and set 95% confidence level limits on the branching fractions $B(t \rightarrow q\gamma) < 3.2\%$ and $B(t \rightarrow qZ) < 33\%$, consistent with standard model expectations.

The vital contributions of the Fermilab staff and the technical staffs of the participating institutions are gratefully acknowledged. This work is supported by the U.S. Department of Energy, the National Science Foundation, the Natural Sciences and Engineering Research Council of Canada, the Istituto Nazionale di Fisica Nucleare of Italy, the Ministry of Education, Science and Culture of Japan, the National Science Council of the Republic of China, and the A. P. Sloan Foundation.

*Visitor.

- [1] S. L. Glashow, J. Iliopoulos, and L. Maiani, *Phys. Rev. D* **2**, 1285 (1970).
- [2] H. Fritzsch, *Phys. Lett. B* **224**, 423 (1989).
- [3] T. Han, R. D. Peccei, and X. Zhang, *Nucl. Phys.* **B434**, 527 (1995).
- [4] References to a particular charge state imply the charge-conjugate state as well.
- [5] S. Parke, in *Proceedings of the 1994 Meeting of the American Physical Society, Division of Particles and Fields* (World Scientific, Singapore, 1995), pp. 726–730.
- [6] F. Abe *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **271**, 387 (1988); F. Abe *et al.*, Report No. FERMILAB-PUB-94/097-E.
- [7] D. Amidei *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **350**, 73 (1994); P. Azzi *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **360**, 137 (1995).

- [8] We use a coordinate system with z along the proton beam axis, azimuthal angle ϕ , and polar angle θ . The pseudorapidity η is defined as $-\ln \tan(\theta/2)$.
- [9] Here $p_T = p \sin \theta$ and $E_T = E \sin \theta$, where θ is the polar angle with respect to the proton beam direction. The difference between the vector sum of all the transverse energies and zero is the missing transverse energy (\cancel{E}_T).
- [10] F. Abe *et al.*, Phys. Rev. D **50**, 2966 (1994).
- [11] F. Abe *et al.*, Phys. Rev. D **45**, 1448 (1992).
- [12] In the standard model, the decay $t \rightarrow Wb$ dominates (the branching fraction is above 99%). In this paper, the term “standard model decay” refers to this mode.
- [13] V. Barger, T. Han, J. Ohnemus, and D. Zeppenfeld, Phys. Rev. D **41**, 2782 (1990).
- [14] F. Abe *et al.*, Phys. Rev. Lett. **74**, 2626 (1995).
- [15] F. Paige and S.D. Protopopescu, Brookhaven Report No. BNL-38034, 1986.
- [16] One source of misidentification is that photon candidates will also pass jet selection requirements.
- [17] F. Abe *et al.*, Phys. Rev. D **52**, 2624 (1995); F. Abe *et al.*, Phys. Rev. D **44**, 29 (1991); F. Abe *et al.*, Phys. Rev. Lett. **69**, 28 (1992).