

Searching for New Physics in $b \rightarrow ss\bar{d}$ Decays

Katri Huitu,¹ Cai-Dian Lü,² Paul Singer,³ and Da-Xin Zhang¹

¹*Helsinki Institute of Physics, University of Helsinki, P.O. Box 9, FIN-00014 Helsinki, Finland*

²*II. Institut für Theoretische Physik, Universität Hamburg, 22761 Hamburg, Germany*

³*Department of Physics, Technion-Israel Institute of Technology, Haifa 32000, Israel*

(Received 19 February 1998; revised manuscript received 18 May 1998)

For any new physics possibly emerging in the future B experiments, the problem is how to extract the signals from the standard model (SM) background. We consider the decay $b \rightarrow ss\bar{d}$ which is very small in the SM. In the minimal supersymmetric SM this decay is possibly accessible in the future experiments. In supersymmetric models with R -parity violating couplings, this channel is not strictly constrained and thus is useful in obtaining bounds on the lepton-number violating couplings. A typical candidate for the suggested search is the $B^- \rightarrow K^- K^- \pi^+$ mode. [S0031-9007(98)07643-1]

PACS numbers: 13.25.Hw, 12.60.Jv

Rare b decays offer a good opportunity to discover new physics beyond the standard model (SM). Many investigations have been done in the past years on the predictions of processes induced by flavor changing neutral current (FCNC) interactions, both within the SM and beyond [1]. One of these FCNC induced processes, $b \rightarrow s\gamma$, has been measured [2] and the branching ratio is comparable with the SM prediction [3] which, however, still contains significant uncertainties. Thus it is hard to make any definite conclusion of signals of new physics. This is also true in most of the channels like $b \rightarrow sq\bar{q}$ [4] and $b \rightarrow s\bar{l}l$ [5], due to the theoretical uncertainties.

In a recent study, Gabbiani *et al.* [6] considered the nonleptonic processes $s \rightarrow dq\bar{q}$ ($q = u, d, s$) more com-

pletely in the supersymmetric model by including also the box diagrams, in addition to the penguin contribution calculated before. Bounds from $b \rightarrow s\gamma$, $B\bar{B}$, and $K\bar{K}$ mixings are considered and their conclusion is that the box diagrams cannot be neglected in the nonleptonic transitions.

Here we will consider a novel channel $b \rightarrow ss\bar{d}$ which turns out to be exceedingly small in the SM. In the SM, this process can be induced by box diagrams with the up-type quarks and weak bosons inside the loop. Because of the strong Glashow-Iliopoulos-Maiani suppression and the small Cabibbo-Kobayashi-Maskawa (CKM) angles involved, the W -box contribution is found to be very small. We perform a simple estimate and find that within the SM,

$$\Gamma = \frac{m_b^5}{48(2\pi)^3} \left| \frac{G_F^2}{2\pi^2} m_W^2 V_{tb} V_{ts}^* \left[V_{td} V_{ts}^* f\left(\frac{m_W^2}{m_t^2}\right) + V_{cd} V_{cs}^* \frac{m_c^2}{m_W^2} g\left(\frac{m_W^2}{m_t^2}, \frac{m_c^2}{m_W^2}\right) \right] \right|^2, \quad (1)$$

where

$$f(x) = \frac{1 - 11x + 4x^2}{4x(1-x)^2} - \frac{3}{2(1-x)^3} \ln x, \quad (2)$$

$$g(x, y) = \frac{4x - 1}{4(1-x)} + \frac{8x - 4x^2 - 1}{4(1-x)^2} \ln x - \ln y.$$

In Eq. (1) the m_c^2/m_W^2 term is numerically about one-half of the highly CKM-suppressed contribution at the amplitude's level. We have dropped in (1) the kinematics dependent contribution which is smaller than 10% of the term proportional to m_c^2/m_W^2 . Even though the relative phase between the two contributions in (1) is unknown, the branching ratio is always less than 10^{-11} , far beyond the designed ability of B factories. By comparing with the analogous processes $B^0\bar{B}^0$ and $K^0\bar{K}^0$ mixings [7], we suppose that including QCD correction will not change the value greatly. Furthermore, the so-called "dipenguin" [8] is only part of the $\mathcal{O}(\alpha_s)$ corrections to the lowest order W -box diagram and is thus less important. In order to consider new physics, this is a clean and useful channel.

If this process were observed at future experiments, we would be confident that there is new physics involved.

In the minimal supersymmetric standard model (MSSM), this transition can be induced by the squark-gaugino (or higgsino) box diagrams. Since $\cot\beta$ is constrained to be small in the MSSM, there is no large contribution from the charged Higgs box diagrams and we will not consider it further. An alternative mechanism for this channel in the supersymmetric models is through the R -parity violating couplings. These two seem to be the only ones capable in mediating this decay within the supersymmetric models without strong suppression. The nonsupersymmetric models like the two Higgs doublet models, including the so-called model III [9], are worth a separate investigation.

To simplify our discussion, we consider only the squark-gluino box which is generally the dominant contribution. Following the mass-insertion approximation [6,10], universal squark masses are assumed, and the squark mixings are described by the off-diagonal elements in the mass squared matrices. We keep only the left-handed

sector in the squark mixing, following the observation made in [11] that the left-right and the right-right sectors are more strongly constrained. The effective Hamiltonian is then

$$\mathcal{H} = -\frac{\alpha_s^2 \delta_{12}^{d*} \delta_{23}^d}{216 m_{\tilde{d}}^2} [24x f_6(x) + 66 \tilde{f}_6(x)] \times (\bar{s} \gamma^\mu d_L) (\bar{s} \gamma_\mu b_L), \quad (3)$$

where $x = m_{\tilde{g}}^2/m_{\tilde{d}}^2$, and the functions $f_6(x)$ and $\tilde{f}_6(x)$ can be found in [6]. δ_{ij}^d parametrizes the mixing between the down-type left-handed squarks. The decay width is calculated as

$$\Gamma = \frac{\alpha_s^4 |\delta_{12}^{d*} \delta_{23}^d|^2 m_b^5}{48(2\pi)^3 m_{\tilde{d}}^4} \left[\frac{2}{9} x f_6(x) + \frac{11}{18} \tilde{f}_6(x) \right]^2. \quad (4)$$

At present, the strongest bounds on the squark mixing parameter δ_{12}^d come from $K\bar{K}$ mixing, and δ_{23}^d from $b \rightarrow s\gamma$ [6]. These bounds are obtained using $\Delta m_K < 3.521 \times 10^{-15}$ GeV and $BR(b \rightarrow s\gamma) < 4 \times 10^{-4}$. They depend on x . Using these bounds, we plot in Fig. 1 the maximum branching ratio of $b \rightarrow ss\bar{d}$ depending on x . When doing numerical calculations, our parameters are chosen as $m_{\tilde{d}} = 500$ GeV, $\tau_B = 1.59$ ps, $f_K = 160$ MeV, and $m_b = 4.5$ GeV. Note that QCD corrections are less important in the MSSM [6].

The MSSM can be extended by including R -parity violating interactions. The term in the R -parity violating part of the superpotential, which is relevant here is

$$W = \lambda'_{ijk} L_i Q_j d_k, \quad (5)$$

where i, j, k are indices for the families and L, Q, d are, under the SM gauge group, the superfields for the lepton doublet, the quark doublet, and the down-type quark singlet, respectively. λ' is a dimensionless coupling. The transition $b \rightarrow ss\bar{d}$ can be induced also by the lepton number violating interactions in W . Following the notations in

$$\Gamma = \frac{m_b^5 f_{\text{QCD}}^2}{512(2\pi)^3} \left(\left| \sum_n \frac{1}{m_{\tilde{\nu}_n}^2} \lambda'_{n32} \lambda_{n21}^{j*} \right|^2 + \left| \sum_n \frac{1}{m_{\tilde{\nu}_n}^2} \lambda'_{n12} \lambda_{n23}^{j*} \right|^2 \right). \quad (9)$$

Note that the couplings are not strongly constrained by the present experiments [15]:

$$\begin{aligned} |\lambda'_{132} \lambda_{121}^{j*}| &< 0.34 \times 0.035, & |\lambda'_{112} \lambda_{123}^{j*}| &< 0.02 \times 0.20, \\ |\lambda'_{232} \lambda_{221}^{j*}| &< 0.36 \times 0.18, & |\lambda'_{212} \lambda_{223}^{j*}| &< 0.09 \times 0.18, \\ |\lambda'_{332} \lambda_{321}^{j*}| &< 0.48 \times 0.20, & |\lambda'_{312} \lambda_{323}^{j*}| &< 0.10 \times 0.20, \end{aligned} \quad (10)$$

if using $m_{\tilde{\nu}_n} = 100$ GeV, and we have

$$\sum_n \sqrt{|\lambda'_{n32} \lambda_{n21}^{j*}|^2 + |\lambda'_{n12} \lambda_{n23}^{j*}|^2} < 0.1, \quad (11)$$

which is too weak to constrain the present mode. Thus a search for this decay mode will improve our knowledge on these couplings. At present, an analysis of this transition at




FIG. 1. The branching ratio of $b \rightarrow ss\bar{d}$ as a function of $x = m_{\tilde{g}}^2/m_{\tilde{d}}^2$ in MSSM without R -parity violation, when the squark-gluino box diagram is included. Region above the line has been excluded by the present data on $b \rightarrow s\gamma$ and on Δm_K .

[12], the tree level effective Hamiltonian is

$$\mathcal{H} = -\sum_n \frac{f_{\text{QCD}}}{m_{\tilde{\nu}_n}^2} (\lambda'_{n32} \lambda_{n21}^{j*} \bar{s}_R b_L \bar{s}_L d_R + \lambda'_{n12} \lambda_{n23}^{j*} \bar{s}_R d_L \bar{s}_L b_R). \quad (6)$$

The QCD corrections to the left-right operators in Eq. (6) have been found to be important [13]. The next-to-leading order QCD corrections are also available [14]. For simplicity, here we include only the leading order QCD corrections which are given by a scaling factor

$$f_{\text{QCD}} = \left(\frac{\alpha_s(m_b)}{\alpha_s(m_t)} \right)^{24/23} \left(\frac{\alpha_s(m_t)}{\alpha_s(m_{\tilde{\nu}_n})} \right)^{24/21} \quad (7)$$

for $m_{\tilde{\nu}_n} > m_t$, and by

$$f_{\text{QCD}} = \left(\frac{\alpha_s(m_b)}{\alpha_s(m_{\tilde{\nu}_n})} \right)^{24/23} \quad (8)$$

for $m_{\tilde{\nu}_n} < m_t$. Using $m_{\tilde{\nu}_n} = 100$ GeV, we estimate $f_{\text{QCD}} \simeq 2$.

Then the decay rate induced by the R -parity violating couplings is

the level of branching ratio $10^{-4} - 10^{-5}$ is realistic, and a negative result will improve the bound in (11) to 10^{-4} . Note that the stricter constraints on $|\sum_n \frac{1}{m_{\tilde{\nu}_n}^2} \lambda'_{n32} \lambda_{n21}^{j*}|$ from ΔM_B and $|\sum_n \frac{1}{m_{\tilde{\nu}_n}^2} \lambda'_{n12} \lambda_{n21}^{j*}|$ from ΔM_K [12] are independent from the present combination of the couplings.

Next we consider the experimental implications of the discussed channel. In the MSSM, the branching ratio of this decay is smaller than $10^{-7} - 10^{-8}$, which is difficult to reach at the B factories, but hopefully is possible at Hera-B or at LHC. In the MSSM with R -parity violation, there is no strict constraint on the mode, and the branching ratio might be quite large. A search for this mode will help to improve the bounds on these λ' -type R -parity violating couplings.

Typical final exclusive processes of $b \rightarrow ss\bar{d}$ include $B^\pm \rightarrow K^\pm K^0 (\bar{K}^0)$, which are difficult to separate from the standard penguin process $b \rightarrow ds\bar{s}$ through $K^0 \bar{K}^0$ mixing. Although the interference of these two sources of the final states is novel in the study of the phenomena such as CP violation, these channels are not suitable for the direct search for the new physics. However, the three-body mode of the charged B decays like $B^- \rightarrow K^- K^- \pi^+$, either a direct three-body transition or through a \bar{K}^{0*} -like resonance, will be a clear signal for this mode. In the neutral B decays, the channel $\bar{B}^0 \rightarrow K^- K^- \pi^+ \pi^+$ is also a clear signal. Similar consideration also applies in other $K^\mp K^\mp +$ (no strange) final states, which can be searched at the B factories. Thus we suggest to search for the signals of multibody channels in the B decays, which will be useful in bounding the R -parity violating couplings at present, and in discovering physics beyond the SM in the future.

To estimate the semi-inclusive rate of $B \rightarrow K^\mp K^\mp +$ (no strange), we assume that the multibody transitions are dominated by the two-body channels which contain the excited states of the K mesons. Because of the short lifetimes of these excited states the mixing effects between the neutral excited states are totally negligible. We denote an excited K as K^* and the ground state as K and estimate the possibility of K^* decays into a charged K by isospin analyses. In the decay $K^* \rightarrow K +$ (no strange) the isospin of the nonstrange system can be $I = 1$ or $I = 0$. In the $I = 1$ channels of the neutral K^* decays, the possibility for a final charged K is $\frac{2}{3}$. This possibility for the charged K^* is $\frac{1}{3}$. In the $I = 0$ channels, all the charged K^* 's decay into charged final K 's while for the neutral K^* 's there is no charged K in the final states. To avoid model calculations for the individual isospin amplitudes, we simply average over both the charged and neutral K^* 's and over all the channels $K^* \rightarrow K$, and we expect that about half of the decays $K^* \rightarrow K$ have charged K 's in the final states. Thus in the $B \rightarrow K^* K^*$ decays induced by $b \rightarrow ss\bar{d}$, a quarter of these transitions materialize as $B \rightarrow K^\mp K^\mp +$ (no strange). Similar analysis applies for the decays $B \rightarrow K^* K$ with a smaller possibility to have two charged K 's in the final states; however, we can expect that there are less $K^* K$ than $K^* K^*$ channels, and the individual transition $B \rightarrow KK$ is even less dominant. We conclude that the estimated $\frac{1}{4}$ possibility of having two charged (same sign) K 's roughly works, and the semi-inclusive process $B \rightarrow K^\mp K^\mp +$ (no strange) consists of about $\frac{1}{4}$ of all the $b \rightarrow ss\bar{d}$ transitions.

Finally, let us consider a similar process $b \rightarrow dd\bar{s}$ due to the same mechanisms. An interesting exclusive channel of this process is $B^- \rightarrow K^+ \pi^- \pi^-$. In the SM, this process suffers even stronger suppression than $b \rightarrow ss\bar{d}$ (by a factor of $|V_{td}/V_{ts}|$ in the amplitude). In the MSSM, the decay rate is proportional to the more strongly constrained $|\delta_{21}^{d*} \delta_{13}^d|^2$. Thus its upper bound is much smaller (only 10^{-4} of $b \rightarrow ss\bar{d}$). However, the hope that $b \rightarrow dd\bar{s}$ can be induced by the λ' -type R -parity violating couplings is still alive. By comparing the involved λ' combination with that in $b \rightarrow ss\bar{d}$, again, there exists only a very loose bound

$$\sum_n \sqrt{|\lambda'_{n31} \lambda_{n12}^{*}|^2 + |\lambda'_{n21} \lambda_{n13}^{*}|^2} < 0.05, \quad (12)$$

and the available data can be used to improve this bound to the order of 10^{-4} .

We thank Kenneth Österberg for helpful discussions. The work of K.H. and D.X.Z. is partially supported by the Academy of Finland (No. 37599). The work of P.S. is partially supported by the Fund for Promotion of Research at the Technion.

-
- [1] For reviews, see A. Masiero and L. Silvestrini, hep-ph/9709244, in *Proceedings of the 2nd International Conference on B Physics and CP Violation (BCONF 97), Honolulu, HI, 1997*, edited by T.E. Browder, F.A. Harris, and S. Pakvasa (World Scientific, Singapore, 1998), p. 172; J.L. Hewett, in *Proceedings of the 28th International Conference on High-Energy Physics (ICHEP 96), Warsaw, Poland, 1996*, edited by Z. Ajduk and A.K. Wroblewski (World Scientific, Singapore, 1997), Vol. 2, p. 1143.
 - [2] CLEO Collaboration, R. Ammar *et al.*, Phys. Rev. Lett. **71**, 674 (1993); CLEO Collaboration, M.S. Alam *et al.*, Phys. Rev. Lett. **74**, 2885 (1995).
 - [3] K. Chetyrkin, M. Misiak, and M. Munz, Phys. Lett. B **400**, 206 (1997).
 - [4] L.L. Chau, H.Y. Cheng, W.K. Sze, H. Yao, and B. Tseng, Phys. Rev. D **43**, 2176 (1991); C.-D. Lü and D.-X. Zhang, Phys. Lett. B **400**, 188 (1997).
 - [5] C.-D. Lü and D.-X. Zhang, Phys. Lett. B **397**, 279 (1997); G. Buchalla and G. Isidori, Nucl. Phys. **B525**, 333 (1998); see also P. Singer, in *Proceedings of the 4th International Workshop on Progress in Heavy Quark Physics, Rostock, Germany, 1997*, edited by M. Beyer, T. Mannel, and H. Schroeder (University of Rostock, Rostock, Germany, 1998), p. 225.
 - [6] F. Gabbiani, E. Gabrielli, A. Masiero, and L. Silvestrini, Nucl. Phys. **B477**, 321 (1996).
 - [7] A. J. Buras, M. Jamin, and P.H. Weisz, Nucl. Phys. **B347**, 491 (1990).
 - [8] J.F. Donoghue, E. Golowich, and G. Valencia, Phys. Rev. D **33**, 1387 (1986); J.O. Eeg and I. Picek, Z. Phys. C **39**, 521 (1988).
 - [9] T.P. Cheng and M. Sher, Phys. Rev. D **35**, 3484 (1987); M. Sher and Y. Yuan, Phys. Rev. D **44**, 1461 (1991); L. Hall and S. Weinberg, Phys. Rev. D **48**, 979 (1993).

-
- [10] M. Dugan, B. Grinstein, and L. Hall, Nucl. Phys. **B255**, 413 (1985); J. S. Hagelin, S. Kelley, and T. Tanaka, Nucl. Phys. **B415**, 293 (1994).
- [11] S. Bertolini, F. Borzumati, A. Masiero, and G. Ridolfi, Nucl. Phys. **B353**, 591 (1991).
- [12] D. Choudhury and P. Roy, Phys. Lett. B **378**, 153 (1996).
- [13] J.L.A. Bagger, K.T. Matchev, and R.J. Zhang, Phys. Lett. B **412**, 77 (1997).
- [14] M. Ciuchini, E. Franco, V. Lubicz, G. Martinelli, I. Scimemi, and L. Silvestrini, Nucl. Phys. **B523**, 501 (1998).
- [15] G. Bhattacharyya, in *Proceedings of the 1st International Conference on Particle Physics Beyond the Standard Model, Beyond the Desert 1997, Castle Ringberg, Tegernsee, Germany, 1997* (Institute of Physics Publishing, Bristol, 1998), p. 194.