

**Search for Second Generation Leptoquarks in the Dimuon Plus Dijet Channel
of $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV**

- F. Abe,¹⁷ H. Akimoto,³⁹ A. Akopian,³¹ M. G. Albrow,⁷ A. Amadon,⁵ S. R. Amendolia,²⁷ D. Amidei,²⁰ J. Antos,³³ S. Aota,³⁷ G. Apollinari,³¹ T. Arisawa,³⁹ T. Asakawa,³⁷ W. Ashmanskas,¹⁸ M. Atac,⁷ P. Azzi-Bacchetta,²⁵ N. Bacchetta,²⁵ S. Bagdasarov,³¹ M. W. Bailey,²² P. de Barbaro,³⁰ A. Barbaro-Galtieri,¹⁸ V. E. Barnes,²⁹ B. A. Barnett,¹⁵ M. Barone,⁹ G. Bauer,¹⁹ T. Baumann,¹¹ F. Bedeschi,²⁷ S. Behrends,³ S. Belforte,²⁷ G. Bellettini,²⁷ J. Bellinger,⁴⁰ D. Benjamin,³⁵ J. Bensinger,³ A. Beretvas,⁷ J. P. Berge,⁷ J. Berryhill,⁵ S. Bertolucci,⁹ S. Bettelli,²⁷ B. Bevensee,²⁶ A. Bhatti,³¹ K. Biery,⁷ C. Bigongiari,²⁷ M. Binkley,⁷ D. Bisello,²⁵ R. E. Blair,¹ C. Blocker,³ S. Blusk,³⁰ A. Bodek,³⁰ W. Bokhari,²⁶ G. Bolla,²⁹ Y. Bonushkin,⁴ D. Bortoletto,²⁹ J. Boudreau,²⁸ L. Breccia,² C. Bromberg,²¹ N. Bruner,²² R. Brunetti,² E. Buckley-Geer,⁷ H. S. Budd,³⁰ K. Burkett,²⁰ G. Busetto,²⁵ A. Byon-Wagner,⁷ K. L. Byrum,¹ M. Campbell,²⁰ A. Caner,²⁷ W. Carithers,¹⁸ D. Carlsmith,⁴⁰ J. Cassada,³⁰ A. Castro,²⁵ D. Cauz,³⁶ A. Cerri,²⁷ P. S. Chang,³³ P. T. Chang,³³ H. Y. Chao,³³ J. Chapman,²⁰ M.-T. Cheng,³³ M. Chertok,³⁴ G. Chiarelli,²⁷ C. N. Chiou,³³ F. Chlebana,⁷ L. Christofek,¹³ M. L. Chu,³³ S. Cihangir,⁷ A. G. Clark,¹⁰ M. Cobal,²⁷ E. Cocca,²⁷ M. Contreras,⁵ J. Conway,³² J. Cooper,⁷ M. Cordelli,⁹ D. Costanzo,²⁷ C. Couyoumtzelis,¹⁰ D. Cronin-Hennessy,⁶ R. Culbertson,⁵ D. Dagenhart,³⁸ T. Daniels,¹⁹ F. DeJongh,⁷ S. Dell'Agnello,⁹ M. Dell'Orso,²⁷ R. Demina,⁷ L. Demortier,³¹ M. Deninno,² P. F. Derwent,⁷ T. Devlin,³² J. R. Dittmann,⁶ S. Donati,²⁷ J. Done,³⁴ T. Dorigo,²⁵ N. Eddy,²⁰ K. Einsweiler,¹⁸ J. E. Elias,⁷ R. Ely,¹⁸ E. Engels, Jr.,²⁸ W. Erdmann,⁷ D. Errede,¹³ S. Errede,¹³ Q. Fan,³⁰ R. G. Feild,⁴¹ Z. Feng,¹⁵ C. Ferretti,²⁷ I. Fiori,² B. Flaugher,⁷ G. W. Foster,⁷ M. Franklin,¹¹ J. Freeman,⁷ J. Friedman,¹⁹ H. Frisch,⁵ Y. Fukui,¹⁷ S. Gadomski,¹⁴ S. Galeotti,²⁷ M. Gallinaro,²⁶ O. Ganel,³⁵ M. Garcia-Sciveres,¹⁸ A. F. Garfinkel,²⁹ C. Gay,⁴¹ S. Geer,⁷ D. W. Gerdes,¹⁵ P. Giannetti,²⁷ N. Giokaris,³¹ P. Giromini,⁹ G. Giusti,²⁷ M. Gold,²² A. Gordon,¹¹ A. T. Goshaw,⁶ Y. Gotra,²⁸ K. Goulian,³¹ H. Grassmann,³⁶ L. Groer,³² C. Grossi-Pilcher,⁵ G. Guillian,²⁰ J. Guimaraes da Costa,¹⁵ R. Guo,³³ C. Haber,¹⁸ E. Hafen,¹⁹ S. R. Hahn,⁷ R. Hamilton,¹¹ T. Handa,¹² R. Handler,⁴⁰ F. Happacher,⁹ K. Hara,³⁷ A. D. Hardman,²⁹ R. M. Harris,⁷ F. Hartmann,¹⁶ J. Hauser,⁴ E. Hayashi,³⁷ J. Heinrich,²⁶ W. Hao,³⁵ B. Hinrichsen,¹⁴ K. D. Hoffman,²⁹ M. Hohlmann,⁵ C. Holck,²⁶ R. Hollebeek,²⁶ L. Holloway,¹³ Z. Huang,²⁰ B. T. Huffman,²⁸ R. Hughes,²³ J. Huston,²¹ J. Huth,¹¹ H. Ikeda,³⁷ M. Incagli,²⁷ J. Incandela,⁷ G. Introzzi,²⁷ J. Iwai,³⁹ Y. Iwata,¹² E. James,²⁰ H. Jensen,⁷ U. Joshi,⁷ E. Kajfasz,²⁵ H. Kambara,¹⁰ T. Kamon,³⁴ T. Kaneko,³⁷ K. Karr,³⁸ H. Kasha,⁴¹ Y. Kato,²⁴ T. A. Keaffaber,²⁹ K. Kelley,¹⁹ R. D. Kennedy,⁷ R. Kephart,⁷ D. Kestenbaum,¹¹ D. Khazins,⁶ T. Kikuchi,³⁷ B. J. Kim,²⁷ H. S. Kim,¹⁴ S. H. Kim,³⁷ Y. K. Kim,¹⁸ L. Kirsch,³ S. Klimenko,⁸ D. Knoblauch,¹⁶ P. Koehn,²³ A. Konigeter,¹⁶ K. Kondo,³⁷ J. Konigsberg,⁸ K. Kordas,¹⁴ A. Korytov,⁸ E. Kovacs,¹ W. Kowald,⁶ J. Kroll,²⁶ M. Kruse,³⁰ S. E. Kuhlmann,¹ E. Kuns,³² K. Kurino,¹² T. Kuwabara,³⁷ A. T. Laasanen,²⁹ S. Lami,²⁷ S. Lammel,⁷ J. I. Lamoureux,³ M. Lancaster,¹⁸ M. Lanzoni,²⁷ G. Latino,²⁷ T. LeCompte,¹ S. Leone,²⁷ J. D. Lewis,⁷ P. Limon,⁷ M. Lindgren,⁴ T. M. Liss,¹³ J. B. Liu,³⁰ Y. C. Liu,³³ N. Lockyer,²⁶ O. Long,²⁶ C. Loomis,³² M. Loretta,²⁵ D. Lucchesi,²⁷ P. Lukens,⁷ S. Lusin,⁴⁰ J. Lys,¹⁸ K. Maeshima,⁷ P. Maksimovic,¹¹ M. Mangano,²⁷ M. Mariotti,²⁵ J. P. Marriner,⁷ G. Martignon,²⁵ A. Martin,⁴¹ J. A. J. Matthews,²² P. Mazzanti,² P. McIntyre,³⁴ P. Melese,³¹ M. Menguzzato,²⁵ A. Menzione,²⁷ E. Meschi,²⁷ S. Metzler,²⁶ C. Miao,²⁰ T. Miao,⁷ G. Michail,¹¹ R. Miller,²¹ H. Minato,³⁷ S. Miscetti,⁹ M. Mishina,¹⁷ S. Miyashita,³⁷ N. Moggi,²⁷ E. Moore,²² Y. Morita,¹⁷ A. Mukherjee,⁷ T. Muller,¹⁶ P. Murat,²⁷ S. Murgia,²¹ M. Musy,³⁶ H. Nakada,³⁷ I. Nakano,¹² C. Nelson,⁷ D. Neuberger,¹⁶ C. Newman-Holmes,⁷ C.-Y. P. Ngan,¹⁹ L. Nodulman,¹ A. Nomerotski,⁸ S. H. Oh,⁶ T. Ohmoto,¹² T. Ohsugi,¹² R. Oishi,³⁷ M. Okabe,³⁷ T. Okusawa,²⁴ J. Olsen,⁴⁰ C. Pagliarone,²⁷ R. Paoletti,²⁷ V. Papadimitriou,³⁵ S. P. Pappas,⁴¹ N. Parashar,²⁷ A. Parri,⁹ J. Patrick,⁷ G. Paulette,³⁶ M. Paulini,¹⁸ A. Perazzo,²⁷ L. Pescara,²⁵ M. D. Peters,¹⁸ T. J. Phillips,⁶ G. Piacentino,²⁷ M. Pillai,³⁰ K. T. Pitts,⁷ R. Plunkett,⁷ A. Pompos,²⁹ L. Pondrom,⁴⁰ J. Proudfoot,¹ F. Ptohos,¹¹ G. Punzi,²⁷ K. Ragan,¹⁴ D. Reher,¹⁸ M. Reischl,¹⁶ A. Ribon,²⁵ F. Rimondi,² L. Ristori,²⁷ W. J. Robertson,⁶ T. Rodrigo,²⁷ S. Rolli,³⁸ L. Rosenson,¹⁹ R. Roser,¹³ T. Saab,¹⁴ W. K. Sakamoto,³⁰ D. Saltzberg,⁴ A. Sansoni,⁹ L. Santi,³⁶ H. Sato,³⁷ P. Schlabach,⁷ E. E. Schmidt,⁷ M. P. Schmidt,⁴¹ A. Scott,⁴ A. Scribano,²⁷ S. Segler,⁷ S. Seidel,²² Y. Seiya,³⁷ F. Semeria,² T. Shah,¹⁹ M. D. Shapiro,¹⁸ N. M. Shaw,²⁹ P. F. Shepard,²⁸ T. Shibayama,³⁷ M. Shimojima,³⁷ M. Shochet,⁵ J. Siegrist,¹⁸ A. Sill,³⁵ P. Sinervo,¹⁴ P. Singh,¹³ K. Sliwa,³⁸ C. Smith,¹⁵ F. D. Snider,¹⁵ J. Spalding,⁷ T. Speer,¹⁰ P. Sphicas,¹⁹ F. Spinella,²⁷ M. Spiropulu,¹¹ L. Spiegel,⁷ L. Stanco,²⁵ J. Steele,⁴⁰ A. Stefanini,²⁷ R. Ströhmer,^{7,*} J. Strologas,¹³ F. Strumia,¹⁰ D. Stuart,⁷ K. Sumorok,¹⁹ J. Suzuki,³⁷ T. Suzuki,³⁷ T. Takahashi,²⁴ T. Takano,²⁴ R. Takashima,¹² K. Takikawa,³⁷ M. Tanaka,³⁷ B. Tannenbaum,²² F. Tartarelli,²⁷ W. Taylor,¹⁴ M. Tecchio,²⁰ P. K. Teng,³³ Y. Teramoto,²⁴ K. Terashi,³⁷ S. Tether,¹⁹ D. Theriot,⁷ T. L. Thomas,²² R. Thurman-Keup,¹

M. Timko,³⁸ P. Tipton,³⁰ A. Titov,³¹ S. Tkaczyk,⁷ D. Toback,⁵ K. Tollefson,¹⁹ A. Tollestrup,⁷ H. Toyoda,²⁴
 W. Trischuk,¹⁴ J. F. de Troconiz,¹¹ S. Truitt,²⁰ J. Tseng,¹⁹ N. Turini,²⁷ T. Uchida,³⁷ F. Ukegawa,²⁶ J. Valls,³²
 S. C. van den Brink,²⁸ S. Vejcik III,²⁰ G. Velev,²⁷ R. Vidal,⁷ R. Vilar,^{7,*} D. Vucinic,¹⁹ R. G. Wagner,¹ R. L. Wagner,⁷
 J. Wahl,⁵ N. B. Wallace,²⁷ A. M. Walsh,³² C. Wang,⁶ C. H. Wang,³³ M. J. Wang,³³ A. Warburton,¹⁴ T. Watanabe,³⁷
 T. Watts,³² R. Webb,³⁴ C. Wei,⁶ H. Wenzel,¹⁶ W. C. Wester III,⁷ A. B. Wicklund,¹ E. Wicklund,⁷ R. Wilkinson,²⁶
 H. H. Williams,²⁶ P. Wilson,⁵ B. L. Winer,²³ D. Winn,²⁰ D. Wolinski,²⁰ J. Wolinski,²¹ S. Worm,²² X. Wu,¹⁰ J. Wyss,²⁷
 A. Yagil,⁷ W. Yao,¹⁸ K. Yasuoka,³⁷ G. P. Yeh,⁷ P. Yeh,³³ J. Yoh,⁷ C. Yosef,²¹ T. Yoshida,²⁴ I. Yu,⁷ A. Zanetti,³⁶
 F. Zetti,²⁷ and S. Zucchelli²

(CDF Collaboration)

¹Argonne National Laboratory, Argonne, Illinois 60439

²Istituto Nazionale di Fisica Nucleare, University of Bologna, I-40127 Bologna, Italy

³Brandeis University, Waltham, Massachusetts 02254

⁴University of California at Los Angeles, Los Angeles, California 90024

⁵University of Chicago, Chicago, Illinois 60637

⁶Duke University, Durham, North Carolina 27708

⁷Fermi National Accelerator Laboratory, Batavia, Illinois 60510

⁸University of Florida, Gainesville, Florida 32611

⁹Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy

¹⁰University of Geneva, CH-1211, Geneva 4, Switzerland

¹¹Harvard University, Cambridge, Massachusetts 02138

¹²Hiroshima University, Higashi-Hiroshima 724, Japan

¹³University of Illinois, Urbana, Illinois 61801

¹⁴Institute of Particle Physics, McGill University, Montreal, Canada H3A 2T8

and University of Toronto, Toronto, Canada M5S 1A7

¹⁵The Johns Hopkins University, Baltimore, Maryland 21218

¹⁶Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany

¹⁷National Laboratory for High Energy Physics (KEK), Tsukuba, Ibaraki 305, Japan

¹⁸Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720

¹⁹Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

²⁰University of Michigan, Ann Arbor, Michigan 48109

²¹Michigan State University, East Lansing, Michigan 48824

²²University of New Mexico, Albuquerque, New Mexico 87131

²³The Ohio State University, Columbus, Ohio 43210

²⁴Osaka City University, Osaka 588, Japan

²⁵Università di Padova, Istituto Nazionale di Fisica Nucleare, Sezione di Padova, I-35131 Padova, Italy

²⁶University of Pennsylvania, Philadelphia, Pennsylvania 19104

²⁷Istituto Nazionale di Fisica Nucleare, University and Scuola Normale Superiore of Pisa, I-56100 Pisa, Italy

²⁸University of Pittsburgh, Pittsburgh, Pennsylvania 15260

²⁹Purdue University, West Lafayette, Indiana 47907

³⁰University of Rochester, Rochester, New York 14627

³¹Rockefeller University, New York, New York 10021

³²Rutgers University, Piscataway, New Jersey 08855

³³Academia Sinica, Taipei, Taiwan 11530, Republic of China

³⁴Texas A&M University, College Station, Texas 77843

³⁵Texas Tech University, Lubbock, Texas 79409

³⁶Istituto Nazionale di Fisica Nucleare, University of Trieste, Udine, Italy

³⁷University of Tsukuba, Tsukuba, Ibaraki 315, Japan

³⁸Tufts University, Medford, Massachusetts 02155

³⁹Waseda University, Tokyo 169, Japan

⁴⁰University of Wisconsin, Madison, Wisconsin 53706

⁴¹Yale University, New Haven, Connecticut 06520

(Received 9 July 1998)

We report on a search for second generation leptoquarks (Φ_2) using a data sample corresponding to an integrated luminosity of 110 pb^{-1} collected at the Collider Detector at Fermilab. We present upper limits on the production cross section as a function of Φ_2 mass, assuming that the leptoquarks are produced in pairs and decay into a muon and a quark with branching ratio β . Using a next-to-leading order quantum chromodynamics calculation, we extract a lower mass limit of $M_{\Phi_2} > 202(160) \text{ GeV}/c^2$ at 95% confidence level for scalar leptoquarks with $\beta = 1(0.5)$. [S0031-9007(98)07705-9]

Leptoquarks are hypothetical bosons which carry both baryon and lepton quantum numbers and mediate interactions between quarks and leptons. They appear in many extensions to the standard model, e.g., grand unified theory, superstring, horizontal symmetry, compositeness, or technicolor [1]. Leptoquarks which combine quarks and leptons of different generations result in flavor-changing neutral currents, which are known to be highly suppressed [2]. While these flavor-changing neutral-current constraints do not exclude such leptoquarks, they restrict them to very high masses. For example, in the Pati-Salam model [3], the masses are expected in the multi-TeV range, and indirect searches for such leptoquarks have been made [4]. For this search, we assume that the leptoquarks couple only to leptons and quarks of the same generation. This leads to the classification of leptoquarks of three generations, denoted as Φ_i , $i = 1, 2, 3$ in this paper.

In $p\bar{p}$ collisions, leptoquarks can be pair produced by gluon-gluon fusion or $q\bar{q}$ annihilation [5]. The contribution to the production rate from direct Φql coupling is suppressed relative to the dominant quantum chromodynamics mechanisms [5]. The coupling strength to gluons is determined by the color charges of the particles, and is model independent in the case of scalar leptoquarks. The production of vector leptoquark pairs is also possible. However, vector leptoquarks have model-dependent trilinear and quadratic couplings to the gluon field [6]. In typical cases the production cross section is orders of magnitude larger than for scalar leptoquarks. The acceptance for vector and scalar leptoquark detection is similar, resulting in much more stringent limits on the vector leptoquark mass.

In this analysis, we report on a direct search for pair produced second generation scalar leptoquarks. The possible decay channels are

$$\begin{aligned} \Phi_2 &\rightarrow q_2\mu^\pm, & \text{branching ratio } \beta, \\ \Phi_2 &\rightarrow q_2\nu_\mu, & \text{branching ratio } 1 - \beta, \end{aligned} \quad (1)$$

where β is the branching ratio to charged lepton decay, and q_2 is a second generation quark (c, s). Our search for Φ_2 production is based on events having a topology including two muons and at least two jets ($\Phi_2\Phi_2 \rightarrow \mu^+\mu^- jj$). The production rate of this decay mode is proportional to β^2 .

A previous CDF study [7] excluded $M_{\Phi_2} < 131(96) \text{ GeV}/c^2$ for $\beta = 1.0(0.5)$ using an integrated luminosity of 19 pb^{-1} . A limit has also been published by D0 [8], which excludes $M_{\Phi_2} < 119(89) \text{ GeV}/c^2$ for $\beta = 1(0.5)$. Searches at LEP-1 have excluded leptoquarks with masses below $45 \text{ GeV}/c^2$ independent of β [9]. Here we present a new limit using an integrated luminosity of 110 pb^{-1} collected during the 1992–1993 and 1994–1995 Tevatron runs (including the 19 pb^{-1} of the previous CDF study). Searches have also been made for first and third generation leptoquark production at the Tevatron [10], LEP [9], and HERA [11]. The

H1 and ZEUS experiments at HERA have reported the observation of an excess of events at high Q^2 [12]. The interpretation of the excess as the production of a first generation leptoquark has been ruled out for large β by the Tevatron results [10].

The CDF detector is described in full detail elsewhere [13]. Only the detector subsystems that are important in this analysis are mentioned here. The momenta of muons are measured in the central tracking chamber (CTC), a 2.76 m diameter cylindrical drift chamber. It is surrounded by a 1.4 T superconducting solenoidal magnet, covering a pseudorapidity (η) range up to 1.1, which allows precision measurements of the transverse momenta (p_T) of charged particles. Inside the CTC a vertex tracking chamber (VTX) allows event vertex reconstruction using tracks over the range $|\eta| < 3.25$. Jets are detected by the calorimeters, which are divided into a central barrel ($|\eta| < 1.1$), end plugs ($1.1 < |\eta| < 2.4$), and forward/backward modules ($2.4 < |\eta| < 4.2$). Outside the calorimeter, central muon drift chambers (CMU) in the region $|\eta| < 0.6$ provide muon identification. Outside the CMU lie the central muon upgrade chambers (CMP), with additional steel between the CMU and CMP detectors to reduce the background from hadrons in the muon sample. The region of $0.6 < |\eta| < 1.0$ is covered by the central muon extension chambers.

We use the PYTHIA Monte Carlo generator [14] with the CTEQ4M parton distribution functions [15] and the renormalization and factorization scales defined as $Q^2 = p_T^2$, together with the CDF detector simulation package, to study the detailed properties of the signal for Φ_2 masses between 100 and $240 \text{ GeV}/c^2$. The signal selection criteria are set according to the kinematic distribution (e.g., the p_T of the muons and E_T of the jets) of decay products determined by Monte Carlo studies, optimized to eliminate the background with a minimal loss of signal events [16].

We select events from several different central single-muon triggers [13] with p_T thresholds of 9 or $12 \text{ GeV}/c$. From these events an exclusive dimuon sample is selected by requiring events with two muons satisfying $p_T > 30 \text{ GeV}/c$ (μ_1) and $p_T > 20 \text{ GeV}/c$ (μ_2). We do not require the two muons to have opposite charge because at very high p_T the charge determination is not reliable. One of the muons is required to have a track from the CTC that matches with a stub in the fiducial region of the central muon detectors (within 2 cm for CMU, and 5 cm for CMU/CMP). The muon satisfying this criterion is defined as a “tight” muon. The other muon can be either a tight muon or a “loose” muon. A loose muon is defined as a CTC track that deposits less than 2 GeV of electromagnetic energy and less than 6 GeV of hadronic energy in the calorimeter tower that it traverses. To ensure good track quality, the track is required to traverse at least 75% of the CTC in the radial direction, and be matched to an interaction vertex determined by the VTX to better than 5.0 cm in the Z direction. Both muons are

required to be isolated, defined as $I < 2 \text{ GeV}$, where I is the sum of transverse energies of all calorimeter towers (excluding the one traversed by the muon) within a cone of $\Delta R = 0.4$ around the direction of the muon, where $\Delta R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$ and ϕ is the azimuthal angle.

The total dimuon identification efficiency, averaged over the data sample, lies between 79% at $M(\Phi_2) = 100 \text{ GeV}/c^2$ and 74.5% at $M(\Phi_2) = 240 \text{ GeV}/c^2$, with the dependence on mass being due to the efficiency of the minimum ionizing requirement. The combined average identification and trigger efficiency is approximately 70% over the mass range $100 < M(\Phi_2) < 240 \text{ GeV}/c^2$.

From this high- p_T dimuon event sample, we require ≥ 2 jets with $E_T^{(1)} > 30 \text{ GeV}$ and $E_T^{(2)} > 15 \text{ GeV}$, respectively. Jets are reconstructed by an algorithm using a fixed cone in η - ϕ space. A detailed description of the algorithm can be found in Ref. [17]. For this analysis a cone of 0.7 is used. Both jets are required to be reconstructed in the region $|\eta| < 2.4$. Jet energy corrections, due to the calorimeter nonlinearity, energy deposited outside the jet cone, underlying energy from other interactions, and the detector geometrical dependence, are applied to determine the μ -jet invariant mass. The Z^0 and other resonances such as the J/ψ or Υ are removed by rejecting events with a dimuon invariant mass in the regions $76 < M_{\mu\mu} < 106 \text{ GeV}/c^2$ and $M_{\mu\mu} < 11 \text{ GeV}/c^2$. After applying these requirements, we are left with a sample of 11 events.

Cosmic rays can fake high- p_T dimuon events; however such muons take a finite time to traverse the detector, generally entering from the top of the detector and exiting at the bottom. We use the hadronic calorimeter timing information and a measurement of the opening angle of the two muons to reject cosmic ray events. None of the 11 selected events is identified as a cosmic ray event.

The numbers of events surviving each selection criterion are listed in Table I. A total of 11 events passing the final selection are shown in Fig. 1, plotted in the muon-jet invariant mass plane ($M_{\mu j}^1$ vs $M_{\mu j}^2$). From two muons and two jets, there are two possible muon-jet pairings. We choose the combination having the smallest invariant mass difference to determine the leptoquark mass for possible candidate events. The reconstructed leptoquark

TABLE I. The number of events surviving each cut, using an integrated luminosity of 110 pb^{-1} from CDF data. The estimated total background from standard model sources is 1.4 ± 1.8 events.

Type of selection	Number of events remaining
Total number of samples	30934
First muon selection (tight cut)	6844
Second muon selection (loose cut)	4153
2-jet cuts	937
Jet E_T cut	64
Invariant mass cut	11
Cosmic ray cut	11

candidates of a pair should have equal mass, within the experimental mass resolution σ_r .

We therefore search for leptoquark candidates by selecting events in a $3\sigma_r$ mass resolution region of the $M_{\mu j}^1$ vs $M_{\mu j}^2$ plane around any given mass, as shown in Fig. 1. The mass resolution, estimated from Monte Carlo studies, depends on the event geometry and the total event energy. Consequently, it varies with the leptoquark mass. For example, the maximum values for the mass resolutions are $\sigma_r(\Phi_2 = 120 \text{ GeV}/c^2) = 21.2 \text{ GeV}/c^2$ and $\sigma_r(\Phi_2 = 240 \text{ GeV}/c^2) = 46.5 \text{ GeV}/c^2$. The asymmetric mass resolution (oval-shaped regions shown in Fig. 1) results primarily from the detector resolution, but also includes a small probability of misidentifying the jet when additional jets exist in the collision and cases for which a wrong muon-jet pairing combination is chosen.

The background for the Φ_2 search include higher order Drell-Yan processes, heavy flavor decay (from $\bar{b}b$ or $\bar{t}t$ in the dimuon channel), WW , or $Z \rightarrow \tau^+\tau^-$. An additional background results from W plus multijet events with a fake muon from energetic hadrons penetrating the shielding to reach the muon chambers, or with a hadron decay to a muon. These background processes are studied using relevant Monte Carlo event samples and actual data samples where possible. The major background is from Drell-Yan processes (we expect ~ 12 events for 110 pb^{-1}) for which the final state includes a muon pair ($Z^0/\gamma \rightarrow \mu^+ \mu^-$) plus two or more jets from initial or final state radiation. There is a small contribution from $\bar{t}t$ (~ 1.3 events). Other backgrounds are negligible due to large muon p_T and jet E_T requirements ($\bar{b}b$




FIG. 1. Invariant mass $M(\mu j)$ distribution for events before applying the mass requirement. A total of 11 candidate events are displayed on the $M(\mu j)_2$ versus $M(\mu j)_1$ plane. The oval configuration shows the limit of the mass requirement for Φ_2 masses between 100 and $240 \text{ GeV}/c^2$. $M(\mu j)_1$ is the invariant mass with the higher p_T muon, while $M(\mu j)_2$ is that with the lower p_T muon.

TABLE II. Results for different Φ_2 masses for an integrated luminosity of 110 pb^{-1} from CDF data. No background subtraction is made in the cross section evaluation. The candidate satisfying $M(\Phi_2) > 200 \text{ GeV}/c^2$ was previously published [6].

Φ_2 mass (GeV/c^2)	120	160	200	240
Total signal detection				
efficiency, ε_{tot}	0.13	0.17	0.20	0.22
Systematic error on ε_{tot}	0.019	0.023	0.023	0.023
Number of				
candidate events	1	0	0	1
Estimated background	3.8	1.1	0.3	0.1
σ (at 95% C.L.) in pb	0.34	0.16	0.13	0.19

and $Z \rightarrow \tau^+ \tau^-$), muon isolation requirements (W plus jets), and small cross section (WW). The total estimated background is 14 ± 1.8 for an integrated luminosity of 110 pb^{-1} before applying the $3\sigma_r$ mass cut. This mass requirement reduces the background substantially, since in the background events the reconstructed muon-jet invariant masses are not correlated. For example, we have estimated the background contribution to be only 0.3 events for $M(\Phi_2) = 200 \text{ GeV}/c^2$ for the 110 pb^{-1} data sample.

In the final result, we do not apply a background subtraction procedure, giving the most conservative estimate of the cross section limit. The number of expected events, N , is given by

$$N = \mathcal{L} \beta^2 \sigma(M_{\Phi_2}) \varepsilon_{\text{tot}}, \quad (2)$$

where \mathcal{L} is the total integrated luminosity of the sample, β is the decay branching ratio to the charged lepton plus quark channel, $\sigma(M_{\Phi_2})$ is the cross section for a given mass, and ε_{tot} is the overall efficiency. We evaluate the factors entering the overall efficiency as a function of Φ_2 mass using actual event samples where possible and otherwise simulated event samples. As shown in Table II, it increases monotonically with $M(\Phi_2)$, from 9% at $M(\Phi_2) = 100 \text{ GeV}/c^2$ to 22% at $M(\Phi_2) = 240 \text{ GeV}/c^2$.

Possible systematic uncertainties of the measured cross section limit have been studied. The major source comes from a limited understanding of the initial and final state gluon radiation. We have used Monte Carlo samples with and without gluon radiation to determine the cross section uncertainty due to this effect. The uncertainty decreases as the Φ_2 mass increases, and it is estimated to be 10% for $M(\Phi_2) = 160 \text{ GeV}/c^2$. A systematic uncertainty also results from the Q^2 scale and the structure functions used. We compute this effect by varying the Q^2 scale between $\frac{1}{4}$ and 4 of the default value ($Q^2 = p_T^2$), and by using other structure functions (CTEQ2L [18] and MRS(A) [19]). The jet energy scaling uncertainty, which results from detector performance limitations, is determined by including a 10% energy uncertainty in the Monte Carlo reconstruction. Other sources of uncertainty, resulting from the detector simulation and the limitation of Monte




FIG. 2. The 95% C.L. cross section limit for Φ_2 production for an integrated luminosity of 110 pb^{-1} . The theoretical cross section curves [17] for $\beta = 0.5$ and 1.0 are superimposed.

Carlo statistics, are relatively small. The uncertainty on the luminosity measurement is 7.2%. The total systematic uncertainty varies with Φ_2 mass, and is computed to be 15% at $M_{\Phi_2} = 120 \text{ GeV}/c^2$ and 10% at $M_{\Phi_2} = 240 \text{ GeV}/c^2$, as listed in Table II.

We compute the 95% confidence level (C.L.) limits on $\sigma(p\bar{p} \rightarrow \Phi_2 \bar{\Phi}_2)\beta^2$, including systematic uncertainties with no background subtraction, as a function of the leptoquark mass (see Table II and Fig. 2). The cross section limit for a given Φ_2 mass does not depend on the coupling λ , and therefore the mass limit does not depend on the choice of the theoretical model but only on β . A theoretical next-to-leading order cross section calculation [20] is also shown in Fig. 2, where the band represents the main uncertainty of the calculation coming from the Q^2 value. Comparing the cross section limit to this calculation, a limit of $M_{\Phi_2} > 202(160) \text{ GeV}/c^2$ for $\beta = 1.0(0.5)$ is derived.

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy, the National Science Foundation, the Istituto Nazionale di Fisica Nucleare (Italy), the Ministry of Science, Culture and Education of Japan, the Natural Sciences and Engineering Research Council of Canada, the National Science Council of the Republic of China, the Schwarz A. P. Sloan Foundation and the A. von Humboldt-Stiftung, and the Swiss National Science Foundation.

*Visitor.

[1] H. Georgi and S. L. Glashow, Phys. Today **33**, No. 9, 30 (1980); J. Schwarz, Phys. Rep. **89**, 223 (1982); E. Witten, Nucl. Phys. **B258**, 75 (1985); S. Pakvasa, Int. J. Mod.

- Phys. A **2**, 1317 (1987); E. Farhi, Phys. Rep. **74**, 277 (1981); B. Schremp and F. Schremp, Nucl. Phys. **B231**, 109 (1984).
- [2] W. Büchmuller and D. Wyler, Phys. Lett. B **177**, 337 (1986).
- [3] J. Pati and A. Salam, Phys. Rev. D **10**, 275 (1974).
- [4] CDF Collaboration, in Proceedings of the 11th Topical Workshop on Proton-Antiproton Collider Physics (PBARP 96), Abano Terme, Italy, 1996 (to be published).
- [5] J. L. Hewett and S. Pakvasa, Phys. Rev. D **37**, 3165 (1988).
- [6] J. L. Hewett *et al.*, in *Proceedings of the Workshop on Physics at Current Accelerators and Supercolliders* (Argonne National Laboratory, Argonne, Illinois, 1993).
- [7] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **75**, 1012 (1995).
- [8] D0 Collaboration, S. Abachi *et al.*, Phys. Rev. Lett. **75**, 3618 (1995).
- [9] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **79**, 4327 (1997); CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **78**, 2906 (1997); D0 Collaboration, B. Abbott *et al.*, Phys. Rev. Lett. **80**, 2051 (1998); D0 Collaboration, B. Abbott *et al.*, Phys. Rev. Lett. **79**, 4321 (1997).
- [10] ALEPH Collaboration, D. DeCamp *et al.*, Phys. Rep. **216**, 253 (1992); DELPHI Collaboration, P. Abreu *et al.*, Phys. Lett. B **316**, 620 (1993); L3 Collaboration, O. Adriani *et al.*, Phys. Rep. **236**, 1 (1993); OPAL Collaboration, G. Alexander *et al.*, Phys. Lett. B **263**, 123 (1991).
- [11] H1 Collaboration, S. Aid *et al.*, Phys. Lett. B **369**, 173 (1996); ZEUS Collaboration, M. Derrick *et al.*, Phys. Lett. B **306**, 173 (1993).
- [12] H1 Collaboration, C. Adloff *et al.*, Z. Phys. C **74**, 191 (1997); H1 Collaboration, J. Breitweg *et al.*, Z. Phys. C **74**, 207 (1997).
- [13] CDF Collaboration, F. Abe *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **271**, 387 (1988); CDF Collaboration, F. Abe *et al.*, Phys. Rev. D **50**, 2966 (1994).
- [14] T. Sjöstrand, Report No. CERN-TH 7112/93, 1994 (to be published).
- [15] CTEQ Collaboration, H. L. Lai *et al.*, Phys. Rev. D **55**, 1280 (1997).
- [16] H. Kambara, Ph.D. thesis, University of Geneva, 1998 (URL at http://www-cdf.fnal.gov/grads/thesis_complete.html).
- [17] CDF Collaboration, F. Abe *et al.*, Phys. Rev. D **45**, 1448 (1992).
- [18] CTEQ Collaboration, H. L. Lai *et al.*, Phys. Rev. D **51**, 4763 (1995).
- [19] A. D. Martin *et al.*, Phys. Rev. D **50**, 6734 (1994).
- [20] M. Krämer, T. Plehn, M. Spira, and P. Zerwas, Phys. Rev. Lett. **79**, 341 (1997).