

Vortex Stability of Interacting Bose-Einstein Condensates Confined in Anisotropic Harmonic Traps

David L. Feder,^{1,2} Charles W. Clark,² and Barry I. Schneider³

¹University of Oxford, Parks Road, Oxford OX1 3PU, United Kingdom

²Electron and Optical Physics Division, National Institute of Standards and Technology, Technology Administration, U.S. Department of Commerce, Gaithersburg, Maryland 20899

³Physics Division, National Science Foundation, Arlington, Virginia 22230

(Received 2 February 1999)

Vortex states of weakly interacting Bose-Einstein condensates confined in three-dimensional rotating harmonic traps are investigated numerically at zero temperature. The ground state in the rotating frame is obtained by propagating the Gross-Pitaevskii equation for the condensate in imaginary time. The total energies between states with and without a vortex are compared, yielding critical rotation frequencies that depend on the anisotropy of the trap and the number of atoms. Vortices displaced from the center of nonrotating traps are found to have long lifetimes for sufficiently large numbers of atoms. The relationship between vortex stability and bound core states is explored. [S0031-9007(99)09382-5]

PACS numbers: 03.75.Fi, 05.30.Jp, 32.80.Pj

The recent experimental achievement of Bose-Einstein condensation (BEC) in trapped ultracold atomic vapors [1–5] has provided a unique opportunity to investigate the superfluid properties of weakly interacting dilute Bose gases. Mean-field theories, which are usually based on the Bogoliubov approximation [6,7] or its finite-temperature extensions [8], yield an excellent description of both the static and dynamic properties of the confined gases [9]. These theories also predict that a continuum Bose condensate with repulsive interactions should be a superfluid, which can exhibit second sound, quantized vortices, and persistent currents. While there exists some evidence for second sound in trapped condensates [10], vortices in these systems have never been observed despite considerable experimental effort [11]. Numerous techniques for the generation of vortices have been suggested, including stirring the condensate with a blue detuned laser [12,13], adiabatic population transfer via a Raman transition into an angular momentum state [14,15], spontaneous vortex formation during evaporative cooling [16,17], and rotation of anisotropic traps [18].

Several studies of vortex stability recently have been carried out [19–23]. The free energy of a singly quantized vortex attains a local maximum when the vortex is centered in a stationary trap [18]. In the presence of dissipation, such a vortex would migrate to the edge of the trap and eventually disappear [20]. It has been suggested [19,20] that this mechanical instability may be related to a bound state in the vortex core, corresponding to a negative-energy “anomalous” dipole mode found numerically in the vortex state at low densities [24]. It is possible to stabilize singly quantized vortices by rotating the trap at an angular frequency Ω . When Ω is larger than the “metastability” frequency Ω_0 , infinitesimal displacements of the vortex no longer decrease the system’s free energy, and the vortex becomes locally stable; above

the critical frequency $\Omega_c \geq \Omega_0$, the vortex state becomes the ground state of the condensate [18].

While rotation of the confining potential has been proposed as a method to both nucleate and stabilize vortices in trapped Bose gases, the relevant critical angular frequencies are not presently known. In the present work, numerical results are obtained for Bose-condensed atoms confined in three-dimensional rotating anisotropic traps at zero temperature. The critical frequency Ω_c is found to increase with the degree of anisotropy in the plane of rotation. In order to *nucleate* vortices, however, the trapped gas must be rotated either more rapidly than Ω_c , or at temperatures above the BEC transition [16,17]. While vortices in nonrotating traps are found to be always unstable, for large numbers of atoms their lifetimes can be very long compared with a trap period.

The trapped Bose condensate, comprised of N_0 repulsively interacting Rb atoms with mass $M = 1.44 \times 10^{-25}$ kg and scattering length $a \approx 100a_0 = 5.29$ nm [25], obeys the time-dependent Gross-Pitaevskii (GP) equation in the rotating reference frame [26]:

$$i\partial_\tau\psi(\mathbf{r},\tau) = \left[-\frac{1}{2}\vec{\nabla}^2 + V_t + V_H - \Omega L_z\right]\psi(\mathbf{r},\tau), \quad (1)$$

where the trap potential is $V_t = \frac{1}{2}(x^2 + \alpha^2y^2 + \beta^2z^2)$, the Hartree term is $V_H = 4\pi\eta|\psi|^2$, and the condensate is rotated about the z axis at the trap center. The effects of gravity (along \hat{z}) are presumed negligible. The constant rotation at frequency Ω induces angular momentum per particle given by the expectation value of $L_z = i(y\partial_x - x\partial_y)$. The trapping frequencies are $(\omega_x, \omega_y, \omega_z) = \omega_x(1, \alpha, \beta)$, with $\omega_x = 2\pi \times 132$ rad/s, $\alpha \geq 1$, and $\beta = \sqrt{8}$ [1]. Choosing the condensate to be normalized to unity yields the scaling parameter $\eta = N_0a/d_x$. Note that energy, length, and time are given throughout in scaled harmonic oscillator units

$\hbar\omega_x$, $d_x = \sqrt{\hbar/M\omega_x} \approx 0.94 \mu\text{m}$, and $T = 2\pi/\omega_x \approx 7.6 \text{ ms}$, respectively.

The ground state of the GP equation is found within a discrete-variable representation (DVR) [27] by imaginary time propagation using an adaptive stepsize Runge-Kutta integrator. A total of between 40 000 and 130 000 DVR points of a Gauss-Hermite quadrature are used, and all calculations are performed on a standard workstation. The stationary ground state in the rotating frame is found by setting $\tilde{\tau} \equiv i\tau$ and solving the diffusion equation:

$$\partial_{\tilde{\tau}}\psi(\mathbf{r}, \tilde{\tau}) = -(H - \mu)\psi(\mathbf{r}, \tilde{\tau}), \quad (2)$$

where H is the GP operator appearing on the right side of Eq. (1), and μ is the chemical potential. The condensate wave function is assumed to be even under inversion of z , and is initially taken to be the vortex-free Thomas-Fermi result, which is the time-independent solution of Eq. (2), neglecting the kinetic energy operator and L_z . A vortex is generated by imposing one quantum of circulation κ , a 2π winding of the phase around the z axis, on the condensate wave function at $\tilde{\tau} = 0$. At each imaginary time step, the chemical potential μ is readjusted in order to preserve the norm of the wave function (i.e., the value of N_0). The propagation continues until the right side of Eq. (2) is equal to a tolerance $\delta \leq 10^{-10}$ defining the error in the dimensionless chemical potential. Stationary solutions are verified by subsequently integrating in real time; any deviations from self-consistency would be made manifest by collective motion.

The solution to the GP equation for a vortex located at the center of a nonrotating anisotropic trap containing $N_0 = 10^5$ atoms is shown in Fig. 1. In general, the vortex core is found to become decreasingly anisotropic as N_0 increases. Furthermore, the condensate density




FIG. 1. The condensate density with a singly quantized vortex at the origin (solid lines) is shown projected along the x and y axes. The vortex-free case is shown as dashed lines for comparison. Parameters for both cases are $N_0 = 10^5$ and $\alpha = 2$, yielding $\xi \approx 0.12d_x$.

preserves its overall vortex-free shape far from the origin except for a slight overall bulge in order to preserve the norm. The structure of the vortex indicates that the healing length ξ is governed largely by the local density and is only weakly dependent on trap geometry. In the Thomas-Fermi (TF) approximation, which is valid for large N_0 , the healing length scales with the TF \hat{x} -axis radius $R = (15\alpha\beta\eta)^{1/5}d_x$ as $\xi \sim (d_x/R)d_x$ [18]. Indeed, the numerics clearly indicate that the mean vortex core radius (approximately d_x at low densities) shrinks very slowly as the TF limit is approached.

A superfluid subjected to a torque will remain purely irrotational until the critical frequency Ω_c is reached, at which point it becomes globally favorable for the system to contain a vortex with a single quantum of circulation κ . In cylindrically symmetric systems where the Hamiltonian commutes with L_z , the circulation and angular momentum (with quantum m) are identical; in the rotating frame, the free energies of the $m \neq 0$ states are shifted by $m\Omega$, and Ω_c is simply the difference in energy between the $m = 1$ and $m = 0$ states (divided by \hbar). In fully anisotropic traps, however, even the $\kappa = 0$ state is shifted, so the applied Ω in Eq. (1) must be increased until the free energy curves cross. It is straightforward to extend the TF estimate of Ω_c [28] to include a small deviation from cylindrical symmetry [18]; neglecting the shift of the vortex-free chemical potential (valid for $\alpha \sim 1$), one obtains

$$\Omega_c \approx \frac{5\alpha}{2} \left(\frac{d_x^2}{R^2} \right) \ln \left(\frac{R}{\xi} \right) \omega_x. \quad (3)$$

Figure 2 shows the critical frequencies for the global stability of a vortex with $\kappa = 1$ at the trap center. For all geometries, the critical frequency drops monotonically as N_0 is increased. For a given number of atoms, the value of Ω_c increases with in-plane anisotropy, similar to the behavior found for liquid helium in rotating elliptical containers [29]. The energy of vortex formation must compensate for that of the irrotational velocity field, which is finite for a vortex-free condensate in a rotating anisotropic trap. The TF result (3) agrees well with the numerical data in its regime of validity $\alpha \sim 1$, though it tends to slightly overestimate the value of Ω_c .

While Ω_c provides the criterion for the global stability of a vortex, it does not necessarily indicate the critical frequency for vortex nucleation. When initially vortex-free condensates are placed in anisotropic traps rotating at a frequency $\Omega < \omega_x$, the velocity field of the stationary solution is found to be irrotational even for $\Omega \gg \Omega_c$. In a harmonic trap with smooth edges, it is not clear if there exists any suitable locus for vortex formation. The vortices are most likely to originate at the condensate surfaces normal to the axis of weak confinement, where the local critical velocity is small [30] but the tangential superfluid velocity in the laboratory frame is largest [29]. While these issues are beyond the scope of the




FIG. 2. The critical frequency for the global stabilization of a singly quantized vortex at the origin is given as a function of the number of atoms in the condensate N_0 and in-plane anisotropy α . Circles, squares, and triangles show numerical results for $\alpha = 1, 2, 3$; bold solid, short dashed, and long dashed lines are the corresponding TF estimates, respectively. The thin solid line represents the frequency of the anomalous mode (shown as a positive value) in the vortex state for $\alpha = 1$.

present issue of vortex stability, there is evidence that multiple vortices appear at higher frequencies [31]. For smaller N_0 , it would likely be easier to generate a vortex experimentally by rotating the anisotropic trap before the condensate is cooled below the BEC transition [16,17].

When $\alpha > 1$, the angular momentum per particle l_κ is a nontrivial function of N_0 , α , and Ω . In a nonrotating system with unit vorticity, l_κ increases with N_0 . In the absence of a vortex, l_κ is finite for a given Ω , and increases with α ; the superfluid velocity \mathbf{v}_s can be locally appreciable but still remain irrotational $\nabla \times \mathbf{v}_s = 0$. At the critical frequency, the difference between l_1 and l_0 is always less than unity; for the most extreme case considered here, a system with $N_0 = 10^6$ and $\alpha = 3$ rotating at $\Omega_c = 0.14\omega_x$, one obtains $l_1 = 2.63\hbar$ and $l_0 = 1.77\hbar$. As $\alpha \rightarrow \infty$, the angular momentum approaches that of a nonsuperfluid TF cloud $l_0 \approx I_{sb}\Omega$ with “solid-body” moment of inertia $I_{sb} = \frac{1}{7}MR^2$.

An anisotropic harmonic oscillator potential becomes unconfining when it is rotated at a frequency between the smallest and largest trapping frequencies. Since Ω_c exceeds ω_x for sufficiently large α , there exists a critical minimum number of condensed atoms N_c able to support a vortex. The value of N_c increases with α and is given by the intercept of the $\Omega/\omega_x = 1$ line in Fig. 2. In cylindrically symmetric systems $N_c = 1$, since in the rotating frame the free energies for all the m states become degenerate at $\Omega = \omega_x$. In the limit of extreme anisotropy $\alpha \rightarrow \infty$ vortices can never be stabilized.

It should be noted that states with vortices at the center of anisotropic harmonic traps are found to be stationary

solutions of Eq. (2) for all values of N_0 and $\Omega \geq 0$ considered; such configurations do not appear to decay in either real or imaginary time. A vortex at the center of a nonrotating cylindrical trap increases the system’s free energy, but is stationary because both the vorticity and angular momentum commute with the Hamiltonian; in principle, the angular momentum can be eliminated, and the free energy reduced, only if this symmetry is broken by displacing the vortex from the center. Since angular momentum is not conserved in anisotropic traps, the apparent vortex stability is likely due to the free energy maximum at the trap center [18]. In the absence of an external pinning mechanism, any such configuration should be unstable against infinitesimal displacements.

In order to further explore the issue of vortex stability in nonrotating traps, the initial condensate phase is wound by 2π a small distance $x_0 \approx 0.2d_x$ from the origin of a trap with $\alpha = 1$. For all values of $N_0 \leq 10^6$, the condensate wave function rapidly (by $\tilde{\tau} \sim T$) converges to a metastable solution with a vortex, where the fluctuations in μ become smaller than $\delta \approx 10^{-7}$ per time step $\Delta\tilde{\tau} \sim 10^{-3}T$. This wave function subsequently decays to the true ground state, but both the real and imaginary time required to do so is found to increase with N_0 [32]. To an excellent approximation, the total time diverges as $\tilde{\tau} \propto N_0^{2/5}T$; for $N_0 \geq 10^5$, the time required $\sim 30T$ becomes computationally inaccessible and the vortex state becomes numerically indistinguishable from stationary. The numerics suggest that while vortices in nonrotating traps are always unstable against off-center displacements, they may be very long-lived.

The observed $x_0 > 0$ instability of the vortex state is likely due to the existence of an anomalous collective mode ω_a at low densities [18,20,24]. This dipole mode, which has positive norm but negative energy (or vice versa), is associated with a zero angular momentum bound state in the vortex core [20]; its value corresponds to the precession frequency of the vortex relative to the cloud [18]. Previous numerical calculations [24] found $|\omega_a| > 0$ for all $N_0 \leq 10^4$. As the core radius shrinks with larger N_0 , however, the anomalous energy might be pushed to zero, yielding long-lived or even stable vortices in the TF limit.

The low-lying excitation frequencies of a nonrotating condensate in the vortex state are calculated using the Bogoliubov equations [6,7]. For completely anisotropic geometries, however, the Bogoliubov operator is too large to diagonalize explicitly. Calculations are therefore restricted to the cylindrical case $\alpha = 1$, where the vortex condensate is $\psi \equiv \psi_1(\rho, z)e^{i\phi}$ and the quasiparticle amplitudes u and v are labeled by m , the projection of the angular momentum operator L_z . The Bogoliubov equations are then

$$\begin{pmatrix} \hat{O} & -V_H \\ V_H & -\hat{O}^\dagger \end{pmatrix} \begin{pmatrix} u_m \\ v_{2-m} \end{pmatrix} = \epsilon_m \begin{pmatrix} u_m \\ v_{2-m} \end{pmatrix}, \quad (4)$$

where $\hat{O} \equiv -\frac{1}{2\rho} \frac{\partial}{\partial \rho} \rho \frac{\partial}{\partial \rho} - \frac{1}{2} \frac{\partial^2}{\partial z^2} + \frac{m^2}{2\rho^2} + V_t + 2V_H$, $\hat{O}' \equiv \hat{O} + \frac{2(1-m)}{\rho^2}$, and $u_1 = v_1 = \psi_1$ when $\epsilon_1 = 0$. In the $\hat{\rho}$ direction, the points of the DVR grid correspond to those of Gauss-Laguerre quadrature, and the kinetic energy matrix elements are obtained using the prescription of Baye and Heenen [33].

The anomalous mode ω_a , which is labeled by $m = 2$, is shown as a function of N_0 in Fig. 2. The results indicate that $0 < |\omega_a| \leq \Omega_c$ for all $N_0 \leq 10^6$ considered. Our calculations suggest that for large numbers of atoms, ω_a coincides with the metastability rotation frequency Ω_0 discussed above; the numerical value of ω_a is consistent with the TF result $\Omega_0 = \frac{3}{5} \Omega_c$ [18]. Indeed, in the frame of a condensate rotating at $\Omega = \omega_a$, the frequency of the vortex oscillation would be Doppler shifted to zero. Alternatively, it can be shown in both the weakly interacting and TF limits that Ω_0 is also the frequency at which the chemical potentials μ for the vortex and vortex-free states become equal; in the TF limit, ω_a vanishes when the vacua (or energy zero) for quasiparticle excitations for both states coincide.

Since the anomalous mode corresponds to the precession of the vortex about the trap origin, one may make a crude estimate of the vortex lifetime τ . In the presence of dissipation, the vortex will spiral out of the condensate after a few orbit periods ω_a^{-1} . Assuming that $\omega_a = \frac{5}{3} \Omega_c$, then with Eq. (3) one obtains $\tau \sim N_0^{2/5} T$ in the TF limit neglecting logarithmic factors. This result is consistent with the time required to yield the vortex-free ground state in the fully three-dimensional numerical calculations discussed above. Similar decay times have been obtained for solitons and vortices in the presence of a small non-condensate component [34].

In summary, we have obtained numerically the critical frequencies Ω_c for the stabilization of a vortex at the center of a rotating anisotropically trapped Bose condensate. Since Ω_c increases with the in-plane anisotropy $\alpha = \omega_y/\omega_x$ and the condensate becomes unconfined for $\Omega > \omega_x$, there is a minimum number of atoms able to support a vortex state. Vortices in nonrotating traps are found to be unstable against small off-center displacements, but their decay time diverges with the total number of atoms.

The authors are grateful for many stimulating discussions with M. Brewczyk, J. Denschlag, M. Edwards, A.L. Fetter, D. Guery-Odelin, D.A.W. Hutchinson, M. Matthews, W.D. Phillips, W. Reinhardt, and C.J. Williams. This work was supported by the U.S. Office of Naval Research.

- [1] M.H. Anderson *et al.*, *Science* **269**, 198 (1995); D.S. Jin *et al.*, *Phys. Rev. Lett.* **77**, 420 (1996).
 [2] K.B. Davis *et al.*, *Phys. Rev. Lett.* **75**, 3969 (1995).
 [3] C.C. Bradley, C.A. Sackett, and R.G. Hulet, *Phys. Rev. Lett.* **78**, 985 (1997).

- [4] L.V. Hau *et al.*, *Phys. Rev. A* **58**, R54 (1998).
 [5] D.G. Fried *et al.*, *Phys. Rev. Lett.* **81**, 3811 (1998).
 [6] N.N. Bogoliubov, *J. Phys. (Moscow)* **11**, 23 (1947).
 [7] A.L. Fetter, *Ann. Phys. (N.Y.)* **70**, 67 (1972).
 [8] A. Griffin, *Phys. Rev. B* **53**, 9341 (1996).
 [9] F. Dalfovo, S. Giorgini, L.P. Pitaevskii, and S. Stringari, *Rev. Mod. Phys.* **71**, 463 (1999).
 [10] D.M. Stamper-Kurn *et al.*, *Phys. Rev. Lett.* **81**, 500 (1998).
 [11] W. Ketterle and E.A. Cornell (private communication).
 [12] B. Jackson, J.F. McCann, and C.S. Adams, *Phys. Rev. Lett.* **80**, 3903 (1998).
 [13] B.M. Caradoc-Davies, R.J. Ballagh, and K. Burnett, *cond-mat/9902092*.
 [14] K.-P. Marzlin, W. Zhang, and E.M. Wright, *Phys. Rev. Lett.* **79**, 4728 (1997); K.-P. Marzlin and W. Zhang, *Phys. Rev. A* **57**, 3801 (1998); **57**, 4761 (1998).
 [15] R. Dum, J.I. Cirac, M. Lewenstein, and P. Zoller, *Phys. Rev. Lett.* **80**, 2972 (1998).
 [16] P.D. Drummond and J.F. Corney, *cond-mat/9806315*.
 [17] R.J. Marshall, G.H.C. New, K. Burnett, and S. Choi, *Phys. Rev. A* **59**, 2085 (1999).
 [18] A.A. Svidzinsky and A.L. Fetter, *cond-mat/9811348*; A.L. Fetter, *cond-mat/9811366*.
 [19] A.A. Svidzinsky and A.L. Fetter, *Phys. Rev. A* **58**, 3168 (1998).
 [20] D.S. Rokhsar, *Phys. Rev. Lett.* **79**, 2164 (1997); D.S. Rokhsar, *cond-mat/9709212*.
 [21] E.J. Müller, P.M. Goldbart, and Y. Lyanda-Geller, *Phys. Rev. A* **57**, R1505 (1998).
 [22] H. Pu, C.K. Law, J.H. Eberly, and N.P. Bigelow, *Phys. Rev. A* **59**, 1533 (1999).
 [23] T. Isoshima and K. Machida, *Phys. Rev. A* **59**, 2203 (1999).
 [24] R.J. Dodd, K. Burnett, M. Edwards, and C.W. Clark, *Phys. Rev. A* **56**, 587 (1997).
 [25] P.S. Julienne, F.H. Mies, E. Tiesinga, and C.J. Williams, *Phys. Rev. Lett.* **78**, 1880 (1997).
 [26] E.P. Gross, *Nuovo Cimento* **20**, 454 (1961); L.P. Pitaevskii, *Zh. Eksp. Teor. Fiz.* **40**, 646 (1961) [*Sov. Phys. JETP* **13**, 451 (1961)].
 [27] B.I. Schneider and D.L. Feder, *Phys. Rev. A* **59**, 2232 (1999).
 [28] S. Sinha, *Phys. Rev. A* **55**, 4325 (1997).
 [29] A.L. Fetter, *J. Low Temp. Phys.* **16**, 533 (1974).
 [30] G.M. Kavoulakis and C.J. Pethick, *Phys. Rev. A* **58**, 1563 (1998).
 [31] D.L. Feder, C.W. Clark, and B.I. Schneider (unpublished).
 [32] Since the GP equation conserves energy and contains no explicit damping terms, one might expect the displaced vortex to precess about the origin indefinitely. In fact, the vortex gradually spirals toward the low-density surface in real time, inducing collective motion of the condensate in the process.
 [33] D. Baye and P.-H. Heenen, *J. Phys. Math. A* **19**, 2041 (1986).
 [34] A.E. Muryshev, H.B. van Linden van den Heuvell, and G.V. Shlyapnikov, *cond-mat/9811408*; P.O. Fedichev and G.V. Shlyapnikov, *cond-mat/9902204*.